



This is a section from the

**2018**

**Mid-Atlantic**

**Commercial Vegetable**

**Production Recommendations**

The manual, which is published annually, is **NOT** for home gardener use.

The **full manual**, containing recommendations specific to New Jersey, can be found on the Rutgers NJAES website in the Publications section:

*<http://njaes.rutgers.edu/pubs/publication.asp?pid=E001>*.

The **label** is a legally-binding contract between the user and the manufacturer. The user must follow all rates and restrictions as per label directions. The use of any pesticide inconsistent with the label directions is a violation of Federal law.

**Cooperating Agencies:** Rutgers, The State University of New Jersey, U.S. Department of Agriculture, and County Boards of Chosen Freeholders. Rutgers Cooperative Extension, a unit of the Rutgers New Jersey Agricultural Experiment Station, is an equal opportunity program provider and employer.

## F. Commodity Recommendations

### Pesticide Use Disclaimer

#### THE LABEL IS THE LAW

**Before using a pesticide, check the label for up to date rates and restrictions.**

Labels can be downloaded from: <http://www.cdms.net/>, <http://www.greenbook.net/> or <http://www.agrian.com/labelcenter/results.cfm>

#### Guide to the Recommended Pesticide Tables in the Following Crop Chapters:

1. Pesticides are listed by **group or code number based on chemical structure and mode of action**, as classified by the Weed Science Society of America (WSSA) for herbicides, the Insecticide Resistance Action Committee (IRAC) for insecticides, and the Fungicide Resistance Action Committee (FRAC) for fungicides.  
**If the number is in bold font, the product may have resistance concerns.**
2. For **restricted use pesticides**, the restricted active ingredients are labeled with a \*. See the Pesticide Safety chapter for more information.
3. **In addition to the pesticides listed below, other formulations or brands with the same active ingredient(s) may be available. ALWAYS CHECK THE LABEL:**
  - a) to ensure a pesticide is labeled for the same use,
  - b) to ensure the pesticide is labeled for the desired crop, and
  - c) for additional restrictions.
4. All pesticide recommendations are made for spraying a **broadcast area of 1 acre** (43,560 square feet). **Adjust the rate for banded applications** (for more information, see the Pest Management chapter, Calibrating Granular Applicators section).
5. Check the label for the maximum amount of pesticide per application and the maximum number of applications per year.
6. **Bee Toxicity Rating (Bee TR):** N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing and method of application are correct, but should NOT be applied directly to crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.

# Cucumbers

For earlier cucumber production and higher, more concentrated yields, use gynoecious varieties. A gynoecious plant produces a high percentage of female flowers and fruit. To produce pollen, 1 to 15% of pollinator must be planted and seed companies add this seed to the gynoecious variety. Both pickling and slicing gynoecious varieties are available. Parthenocarpic cucumbers that produce fruit without pollination are also available for protected culture and field production.

## Recommended Varieties<sup>1</sup>

	Variety	Days	F1 <sup>2</sup>	Type <sup>3</sup>	Use <sup>4</sup>	Reported Disease Resistance <sup>5</sup>								
						Scab (Ccu)	PM (Px)	AN (Co)	DM <sup>5</sup> (Pcu)	ALS (Psl)	Cmv	Wmv	Zmv	Prsv
<b>Standard Slicing Varieties</b>	Bristol	54	Yes	Gyn	F	X	X	X		X	X	X	X	X
	Dasher II	58	Yes	Gyn	F	X	X	X		X	X			
	Dominator	55	Yes	Gyn	F	X	X	X		X	X			
	General Lee	66	Yes	Gyn	F	X	X				X			
	Indy	59	Yes	Gyn	F	X	X	X		X	X	X	X	X
	Intimidator	53	Yes	Gyn	F	X	X	X		X	X			
	Mongoose	55	Yes	Gyn	F	X	X	X		X	X	X	X	X
	Python	55	Yes	Gyn	F	X	X	X		X	X			
	Speedway	56	Yes	Gyn	F	X	X	X		X	X			
	Stonewall	53	Yes	Gyn	F	X	X	X		X	X			
	SV3462CS	56	Yes	Gyn	F	X	X	X	X	X			X	
	SV4719CS	56	Yes	Gyn	F	X	X	X	X	X			X	
	Talladega	61	Yes	Gyn	F	X	X	X		X	X	X		
Thunder	58	Yes	Gyn	F	X	X	X		X	X		X		
<b>Slicers Long Types</b>	Suyo Long	61	No	Mon	F		X							
	Tasty Green	52	Yes	Mon	F		X							
<b>Pickels</b>	Bowie	51	Yes	Parth	MP	X	X							
	Calypso	51	Yes	Gyn	HF	X	X	X		X	X			
	Citadel	52	Yes	Gyn	HMP	X	X	X	X	X	X			
	Eureka	57	Yes	Mon	HF	X	X	X		X	X	X		X
	Expedition	50	Yes	Gyn	MP	X	X	X		X	X			
	Fanci Pak	53	Yes	Gyn	HF	X	X	X		X	X			
	Feisty	57	Yes	Gyn	MP	X	X	X		X	X			
	Jackson Supr.	52	Yes	Gyn	HMFP	X	X	X		X	X			
	Lafayette	52	Yes	Gyn	MP	X	X	X		X	X			
	Logan	51	Yes	Gyn	MP	X	X	X		X	X			
	Max Pack	57	Yes	Mon	FH	X	X	X		X	X	X	X	X
	NQ5007	50	Yes	Parth	MP	X	X	X		X	X			
	NQ5543	49	Yes	Parth	MP	X	X	X		X	X			
	Peacemaker	52	Yes	Gyn	MHP	X	X	X	X	X	X			
	Puccini	50	Yes	Parth	HMFP	X	X	X		X	X			
SV7140CN	50	Yes	Gyn	MP	X	X	X		X	X				
Vlaspik	51	Yes	Gyn	MP	X	X	X		X	X				
<b>Protected Culture / High Tunnels</b>	Corinto	48	Yes	Parth	F	X					X			
	Cucapa	48	Yes	Parth	F		X				X			
	Lisboa	60	Yes	Parth	F	X								
	Picolino	45	Yes	Parth	F		X				X			
	Rocky	46	Yes	Parth	F	X	X							
Socrates	52	Yes	Parth	F	X	X								

<sup>1</sup>Varieties are listed alphabetically. <sup>2</sup>Hybrid. <sup>3</sup>Gyn=Gynoecious or mostly female flowers; 5-15% of a monoecious pollinizer variety added; Mon=Monoecious type with female and male flowers; Parth=Parthenocarpic type that sets fruit without pollination. <sup>4</sup>F=Fresh Market, P=Processing (pickling), H=Hand harvest multiple times, M=Machine harvest once. <sup>5</sup>X=high or intermediate level of resistance to Scab, PM=Powdery Mildew, AN=Anthracnose, DM=Downy Mildew, ALS=Angular Leaf Spot, Cmv=Cucumber mosaic virus, Wmv=Watermelon Mosaic Virus, Zmv=Zucchini yellows mosaic virus, Prsv=Papaya ring spot virus. <sup>5</sup>Only varieties with some resistance to the current strain of downy mildew are noted with an X.

## Recommended Nutrients Based on Soil Tests

In addition to using the table below, check the suggestions on rate, timing, and placement of nutrients in your soil test report and the Soil and Nutrient Management chapter. Your state's soil test report recommendations and/or your farm's nutrient management plan supersede recommendations found below.

Cucumbers <sup>1</sup>	N (lb/A)	Soil Phosphorus Level				Soil Potassium Level				Nutrient Timing and Method
		Low	Med	High (Opt)	Very High	Low	Med	High (Opt)	Very High	
	80-150	150	100	50	0 <sup>2</sup>	200	150	100	0 <sup>2</sup>	Total nutrient recommended
	25-50	125	75	25	0 <sup>2</sup>	175	125	75	0 <sup>2</sup>	Broadcast and disk-in
	25	25	25	25	0	25	25	25	0	Band place with planter
	25-75	0	0	0	0	0	0	0	0	Sidedress when vines begin to run

<sup>1</sup>For plasticulture, fertilization rates are based on a standard row spacing of 6 ft.

<sup>2</sup>In VA, crop replacement values of 25 lb/A of P<sub>2</sub>O<sub>5</sub> and 50 lb/A of K<sub>2</sub>O are recommended on soils testing Very High.

## Fertigation Schedule Examples

This table provides examples of fertigation schedules based on two common scenarios - sandy coastal plain soils and heavier upland soils. It should be modified according to specific soil tests and base fertility.

Fertigation recommendations for 125 lb N and 125 lb K <sub>2</sub> O <sup>1,2</sup>								
For soils with organic matter content less than 2% or coarse texture and low to medium or deficient K								
Preplant (lb/A) <sup>3</sup>			Nitrogen			Potash		
			25			50		
Stage and Description	Weeks	Days	lb/day	lb/week	lb/stage	lb/day	lb/week	lb/stage
1 Early vegetative	1	1-7	0.5	3.5	3.5	0.4	2.8	2.8
2 Late vegetative	2-3	8-14	0.9	6.3	12.6	0.7	4.9	9.8
3 Fruiting and harvest	4-7	15-42	1.4	9.8	39.2	0.9	6.3	25.2
4 Later harvest <sup>4</sup>	8-10	43-70	0.9	6.3	18.9	0.6	4.2	12.6

  

Fertigation recommendations for 75 lb N and 50 lb K <sub>2</sub> O <sup>1,2</sup>								
For soils with organic matter content greater than 2% or fine texture and high or optimum K								
Preplant (lb/A) <sup>3</sup>			Nitrogen			Potash		
			50			50		
Stage and Description	Weeks	Days	lb/day	lb/week	lb/stage	lb/day	lb/week	lb/stage
1 Early vegetative	1	1-7	1	7	7	1	7	7
2 Late vegetative	2-3	8-14	1.5	10.5	21	1.6	11.2	22.4
3 Fruiting and harvest	4-7	15-42	2.2	15.4	61.6	2.2	15.4	61.6
4 Later harvest <sup>4</sup>	8-10	43-70	1.7	11.9	35.7	1.6	11.2	33.6

<sup>1</sup>Rates are based on 7,260 linear bed ft/A (6 ft bed spacing). If beds are closer or wider, fertilizer rates should be adjusted proportionally. Drive rows should not be used in acreage calculations (see the Fertigation section in the Irrigation Management chapter). <sup>2</sup>Base overall application rate on soil test recommendations. <sup>3</sup>Applied under plastic mulch to effective bed area using modified broadcast method. <sup>4</sup>For extended harvest after 10 weeks continue fertigation at this rate.

## Plant Tissue Testing

Plant tissue testing can be a valuable tool to assess crop nutrient status during the growing season, to aid with in-season fertility programs or to evaluate potential deficiencies or toxicities.

Critical cucumber tissue test values for most recently matured leaves up to first bloom: N 3.5-6 %, P 0.3-0.6 %, K 1.6-3.0 %, Ca 2-4 %, Mg 0.5-0.7% and S 0.3-0.8%. For additional nutrients and other growth stages consult with a tissue testing laboratory or this web link at the University of Florida: <http://edis.ifas.ufl.edu/ep081>

**Seed Treatment** Seed should be treated; check with your seed company and see Disease Control below.

## Planting Dates

Start seeding for transplanting in mid-April in warmer, southern areas and May 10 in PA and other cool areas. Successive plantings can be made through early August. Container-grown plug plants are planted through the plastic when daily mean temperatures have reached 60°F (16°C). Planting dates vary from April 20 in southern regions to June 20 in northern areas. Early plantings should be protected from winds with hot caps, tents, row covers or rye windbreaks.

## Spacing

**Slicers:** Space rows 3-4 ft apart with plants 9-12 inches apart. Seeding rate: 1.5 lb/A.

**Machine Harvest Pickles:** Research and field data have shown that 55,000-65,000 plants/A is the optimum population for yield and quality. To accommodate a harvester with an 84-inch head, 3 rows 26-28 inches apart should be planted on each bed with plants 4-5 inches apart in the row. If the harvester has a 90-inch head, space rows 30 inches apart and space plants 3-4 inches apart in the row. For machine harvest of pickles, high plant populations concentrate pickle maturity. Parthenocarpic pickles are being trialed in the region. These are planted to achieve 22,000 to 28,000 plants/A.

**Hand Harvest Pickles:** Space rows 3-4 ft apart with plants 6-8 inches apart. Seeding rate: 1.5-2 lb/A.

## Mulching and Fumigation

Plastic mulch laid on moist soil before field planting conserves moisture, and increases soil temperature and early and total yield. Various widths of plastic are available; choose one that works with your production system and equipment. Fumigation will be necessary when there is a history of soil-borne diseases in the field; several fumigants can be used on cucumber depending on what the predominant pests are (see the Soil Fumigation section in the Pest Management chapter). Fumigation also aids in the control of weeds. Fumigant and mulch should be applied to well-prepared planting beds; check the fumigant label for the plant-back period that must be adhered to for crop safety. Plastic should be laid immediately over the fumigated soil. Fumigation alone may not provide satisfactory weed control under plastic. Black plastic can be used without a herbicide to provide control of most weeds.

Fertilizer must be applied during bed preparation. At least 50% of the N should be in the nitrate ( $\text{NO}_3^-$ ) form. Drip (trickle) irrigation is recommended for plastic mulch systems and tape is laid at the same time as mulch. Foil and highly reflective mulches can be used to repel aphids that transmit viruses in fall-planted (after July 1) cucurbits. Direct seeding through the mulch is recommended for maximum virus protection; transplants should not be used with foil mulches. Also, an herbicide is not necessary.

Cucumbers also have been successfully grown in no-till systems on cover crop mulch.

## Irrigation

Cucumbers require irrigation for best yield and quality. During flowering and fruiting water use can be over 0.25 inches/day and water deficit during this period will have the greatest negative impact on yield and fruit quality. A balance must be struck, however, between maintaining adequate moisture for fruiting while minimizing wetness in the canopy and on the soil surface which promotes fruit rots and downy mildew.

## Trellising

Fresh market slicer cucumbers and pickles may be produced on trellises which may result in 2-3 times greater average yield than in non-trellised fields. Trellising is the preferred system in high tunnels. Trellising incurs a higher cost than growing cucumbers on the ground, but it has the following benefits:

1. Improved fruit quality, particularly with respect to color and shape (no yellow “ground spot”).
2. More effective control of many diseases and insects.
3. Less damage to vines resulting in a longer harvest season.
4. More consistent and thorough harvesting resulting in fewer jumbos and culls.
5. Easier harvesting than ground grown cucumbers.

Erect the trellis so that it is 6 ft high with a top (No. 8) and bottom (No. 12) wire and plastic twine or netting tied between the two wires at each plant. Posts or poles should be no more than 15 ft apart and the top wire should be very taut. An additional brace between posts may be required when the fruit load becomes heavy. In high tunnels, wires are stretched at the height desired and plastic twine is used to train plants. Training the main stem is required until it reaches and extends over the top wire. Pruning lateral runners near the base of the plant will result in higher

yields. The first 4-6 lateral runners that appear should be removed. Other runners above this point should be allowed to run. Single stem systems are often used in high tunnels.

## Pollination

Honeybees, squash bees, bumblebees and other wild bees are important for proper cucumber pollination and fruit set. In high tunnels bumblebees are particularly effective. Populations of pollinating insects may be adversely affected by insecticides applied to flowers or weeds in bloom. Apply insecticides only in the evening hours or wait until bloom is completed before application. See the section on Pollination in the General Production Recommendations chapter. Bee Toxicity ratings are available in the insecticide tables below.

## Parthenocarpic Cucumbers

Parthenocarpic cucumbers do not require pollination to set fruit. They will be nearly seedless or have unformed seeds. They should be isolated from seeded cucumber types to increase productivity and maintain the seedless nature. Parthenocarpic types should be considered when bee activity is limited such as in high tunnels, under row covers, or in very early plantings.

## Season Extension

**Low Tunnel Cucumber Production** Cucumbers for early production may be successfully grown in high tunnels, in low tunnels with perforated clear plastic row covers, or using floating row covers. Use plastic mulch and trickle irrigation as discussed above. The following field system - similar to that used for early sweet corn - is also successful: A modified bedshaper is used to form a ridge on each side of the plant row, leaving a suitable area for planting. A 36-inch wide piece of embossed clear plastic is then used to cover the plant row, leaving a 5-6 inch high space between the planted row and the plastic cover. It is estimated that temperatures may be increased 10-20°F depending on time of planting and sunlight availability and intensity.

**High Tunnel Cucumber Production** Cucumbers are a potentially profitable crop for spring and fall production within a high tunnel. Cucumbers mature in approximately half the length of time required for tomato ripening. Cucumbers are also amenable to vertical trellising which increases production and quality. High tunnel cucumber varieties are often parthenocarpic (requiring no pollenizers) although gynocious varieties can also be used (with pollenizers). Cucumbers can be established by direct seeding or transplanting. Space plants 12-18 inches apart in-row on 42-48 inch bed centers. High tunnel varieties can remain unpruned, though pruning can reduce pest infestation and improve marketable yield. If pruning is done, the lower laterals (suckers) should be pruned on the bottom 2 ft leaving 1 or 2 stems per plant to trellis. More information on relative planting and harvesting dates is available under "High Tunnels" in the General Production Recommendations chapter.

**Greenhouse Production** Varieties are usually parthenocarpic varieties bred specifically for the lower light conditions of fall, winter, and early spring. European "English" or "Dutch" types and Asian types are available. Hydroponic nutrient solution systems are commonly used and cucumbers are trellised with single or double stems trained onto twine; see also Greenhouse Production in the General Production Recommendations chapter.

## Harvest and Storage

Cucumbers should be harvested when they have reached full size for the variety but while seeds are still soft. For slicers and manually harvested pickles, multiple harvests at 2-3 day intervals will be necessary. Machine-harvested pickles are harvested once when less than 5% have become oversized, as this produces the highest bushel yields. Size requirements of processors will also dictate schedules for machine and hand harvesting pickles.

Cucumbers can be held for 10-14 days at 50-55°F (10-13°C) with a relative humidity of 90-95%. At 50°F and above, cucumbers ripen rapidly, with the green color changing to yellow, starting after about 10 days. The color change is accelerated if cucumbers are stored in the same room as apples, tomatoes, or other ethylene-producing crops. Cucumbers for fresh market are usually waxed to reduce moisture loss. Cucumbers are subject to chilling injury if held below 50°F for longer than about 2 days.

## Weed Control

**THE LABEL IS THE LAW - See the Pesticide Use Disclaimer on page F 1.**

### Recommended Herbicides

1. Identify the weeds in each field and select recommended herbicides. More information is available in the “Herbicide Effectiveness on Common Weeds in Vegetables” Table (E-2) in the Pest Management chapter.
2. Minimize herbicide resistance development. Identify the herbicide site mode of action group and follow recommended good management practices. Include non-chemical weed control whenever possible.

### Weed Control Under Plastic Mulch

Black plastic mulch effectively controls most annual weeds by blocking light from germinated seeds. Herbicides control weeds under the mulch and around the planting hole. Irrigation tubing on the soil surface may cause weed problems by leaching herbicide away at the emitters, especially when clear plastic is used. Bury tubing several inches deep in the bed to reduce this problem.

1. Complete soil tillage, and form raised beds. **Do not** apply residual herbicides before forming beds, or herbicide rate and depth of incorporation may be increased, raising the risk of crop injury. Apply herbicide(s) in a band as wide as the mulch after the bed is formed, as a part of the same operation.
2. Lay the plastic mulch and trickle irrigation tubing. Irrigate to activate the herbicide if necessary. **Condensation that forms on the underside of the mulch will activate the herbicide.** Delay punching the planting holes until seeding or transplanting.

### Weed Control For Soil Strips Between Rows Of Plastic Mulch

Follow the instructions below or crop injury and/or poor weed control may result.

1. Spray preemergence herbicide(s) in bands onto the soil and the shoulders of the plastic mulch before planting and weed germination. **OR** apply after planting as a shielded spray combined with a postemergence herbicide to control emerged weeds. **Do not broadcast spray over the plastic mulch unless specifically stated on the label!**
2. Incorporate preemergence herbicide into the soil with 0.5 to 0.75 inch of rainfall or overhead irrigation within 48 hr of application.

### Weed Control For Seeding Into Soil Without Plastic Mulch

Use the following instructions or crop injury and/or poor weed control may result:

1. Complete soil tillage, apply preplant herbicide(s), and incorporate. Use a finishing disk or field cultivator that sweeps at least 100% of the soil surface twice, at right angles, operated at a minimum of 7 miles per hour (mph), **OR** a PTO driven implement once, operated at less than 2 mph.
2. Seed and apply preemergence herbicide(s) immediately after completing soil tillage, and mechanical incorporation of preplant herbicides. Irrigate if rainfall does not occur, to move the herbicide into the soil and improve availability to germinating weed seeds within 2 days of when the field was last tilled, or plan to control escaped weeds by other methods.

Labeled Applications Sites for Cucumbers									
Herbicides	WSSA group number	Plastic mulch production					Bare-ground production		
		Soil-Applied		Postemergence			Soil-applied	POST	Post-harvest
		Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest			
Sandea	2	YES	YES	YES	YES		YES	YES	
Treflan	3		YES						
Curbit	3		YES				YES		
Prefar	8	YES	YES				YES		
Command	13		YES				YES		
Strategy	3+13		YES				YES		
Select	1			YES	YES			YES	
SelectMax	1			YES	YES			YES	
Poast	1			YES	YES			YES	
Gramoxone	22				YES	YES			YES

1. Soil-Applied						
Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	14	12
<p><b>-Plasticulture:</b> can be applied in a band under the plastic, immediately before laying the mulch; delay seeding or transplanting for 7 days after application. Row row middles: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v or include a non-selective herbicide.</p> <p><b>-Bareground:</b> apply broadcast after seeding but before crop emergence or no sooner than 7 days before transplanting.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. <b>Do not</b> use Group 2 herbicides repeatedly in the same field. <b>Do not</b> apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum number of Sandea applications per year is 2 and <b>do not</b> exceed 2 oz/A during the crop season.</p>						
3	Curbit 3EC	1 to 3 pt/A	ethalfluralin	0.38 to 1.13 lb/A	--	24
<p><b>-Plasticulture</b> row middles only: apply as a banded spray after crop emergence or transplanting. <b>Do not</b> soil incorporate.</p> <p><b>-Bareground:</b> apply broadcast after direct-seeding but prior to crop emergence; <b>do not</b> use on transplanted cucumbers.</p> <p>-Controls annual grasses and certain annual broadleaf weeds, including carpetweed and pigweed sp.</p> <p>-Use lower rate for coarse-textured soils or soils with low organic matter.</p> <p>-Where overhead irrigation is available, activate Curbit with 0.5 inch of irrigation within 2 days after application; if no irrigation or rainfall occurs within 5 days of application, activity of Curbit can be reduced.</p> <p>-Available as a pre-mix herbicide Strategy. Strategy at 3 pt/A= Curbit at 26 fl oz (0.6 lb ai) and Command at 8 fl oz (0.188 lb ai)</p> <p>-Maximum applications per season: not specified</p>						
3	Treflan 4EC	1 to 2 pt/A	trifluralin	0.5 to 1 lb/A	30	12
<p><b>-Plasticulture</b> row middles only: apply as a directed spray after emergence when plants have reached the 3 to 4 true leaf stage of growth.</p> <p><b>-Not labeled for bareground production.</b></p> <p>-Primarily controls annual grasses with a few broadleaf weeds. <b>Do not</b> use (or reduce the rate) when cold, wet soil conditions are expected, or crop injury may result. Maximum applications per season: not specified.</p>						
3 + 13	Strategy 2.ISC	1.5 to 6 pt/A	ethalfluralin plus clomazone	0.39 to 1.58 lb/A	45	24
<p><b>-Plasticulture:</b> row middles application. <b>Bareground:</b> apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Strategy is a prepackage mixture of Curbit 3EC and Command 3ME. Refer to individual products for comments.</p> <p><b>-Do not</b> apply prior to planting crop. <b>Do not</b> soil incorporate. Maximum applications per season: not specified.</p>						
8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	--	12
<p><b>-Plasticulture:</b> under plastic: apply in a band under the plastic, immediately before laying the mulch. Plasticulture: row middles application is labeled.</p> <p><b>-Bareground:</b> apply preemergence or preplant incorporated. Preemergence applications should be followed by irrigation within 36 hrs (apply enough water to wet the soil at least 2 to 4 inches deep). Preplant incorporated applications should be incorporated 1 to 2 inches deep (deeper than 2 inches will result in reduced weed control).</p> <p>-Provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters.</p> <p>-Maximum applications per season: not specified.</p>						
13	Command 3ME	4 to 8 fl oz/A	clomazone	0.094 to 0.19 lb/A	45	12
<p><b>-Plasticulture:</b> row middles application only.</p> <p><b>-Bareground:</b> apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Use the lower rate when used on coarse-textured soils low in organic matter, when weed pressure is light, or to minimize herbicide carryover that could affect subsequent crops.</p> <p>-Controls annual grasses and many broadleaf weeds including common lambsquarters, velvetleaf, spurred anoda, and jimsonweed. Carpetweed, morningglory sp., pigweed sp., and yellow nutsedge will not be controlled. Higher rates will improve control (or expand number of species controlled) such as common cocklebur, common ragweed, or jimsonweed (refer to label for specific weeds and rates).</p> <p><b>-WARNING:</b> Command spray or vapor drift may injure sensitive crops and other vegetation up to several hundred yards from the point of application (refer to label for restrictions).</p> <p>-Available as a pre-mix herbicide Strategy: Strategy at 3 pt/A= Command at 8 fl oz (0.188 lb ai) and Curbit at 26 fl oz (0.6 lb ai)</p> <p>-Maximum number of Command applications per year: 1.</p>						

2. Postemergence						
Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC	6 to 8 fl oz/A	clethodim	0.094 to 0.13 lb/A	14	24
	Select Max 0.97EC	12 to 16 fl oz/A				
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.19 to 0.28 lb/A	3	12

2. Postemergence (Select, Poast) continued on next page

2. Postemergence (Select, Poast) continued on next page



## F Cucumbers

<p>-Postemergence as broadcast spray with both plasticulture and bareground</p> <p>-<b>Select 2EC</b>: use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). <b>Select Max</b>: use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). <b>Poast</b>: Apply with COC at 1.0% v/v. <b>The use of COC may increase the risk of crop injury when hot or humid conditions prevail.</b> To reduce the risk of crop injury, omit additives or switch to nonionic surfactant when grasses are small and soil moisture is adequate.</p> <p>-Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.</p> <p>-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled. Controls many annual and certain perennial grasses, including annual bluegrass, but will not consistently control goosegrass. Control may be reduced if grasses are large or under hot or dry weather conditions. If repeat applications are necessary, allow 14 days between applications.</p> <p>-<b>Do not</b> tank-mix with or apply within 2 to 3 days of any other pesticide - unless labeled - as this may increase the risk of crop injury or reduce the control of grasses.</p> <p>-Rainfastness is 1 hr. <b>Do not</b> apply more than 8 fl oz of Select 2EC in a single application and <b>do not</b> exceed 32 fl oz/A for the season; <b>do not</b> apply more than 16 fl oz of Select Max in a single application and <b>do not</b> exceed 64 fl oz/A for the season.</p> <p>-<b>Do not</b> apply more than 1.5 pt/A Poast in single application and <b>do not</b> exceed 3 pt/A for the season.</p>						
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	14	12
<p>-<b>Plasticulture</b>: broadcast (over the top) or directed to row middles; broadcast for bareground.</p> <p>-<b>Bareground</b>: apply Sandea after the crop has at least 3 to 5 true leaves but before first female flowers appear and no sooner than 14 days after transplanting. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf; control of weeds taller than 3 inches may not be adequate. Sandea will not control common lambsquarters or eastern black nightshade if applied postemergence; for row middle application, tankmix with a non-selective herbicide to increase spectrum of control.</p> <p>-Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. <b>Do not</b> use Group 2 herbicides repeatedly in the same field.</p> <p>-<b>Do not</b> apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Rainfastness is 4 hrs. Maximum number of Sandea applications per year is 2 and <b>do not</b> exceed 2 oz/A during the crop season</p>						
22	Gramoxone 2SL	1.95 pt/A	paraquat*	0.49 lb/A	14	24
<p>-<b>A Supplemental Label has been approved for the use of Gramoxone 2SL for postemergence weed control in DE, MD, NJ, PA, and VA.</b> Row middles as a shielded application. Apply as a directed spray in a minimum of 20 gal spray mix/A to control emerged weeds between the rows after crop establishment. Include a nonionic surfactant at 0.25% v/v.</p> <p>-Use shields or hoods to prevent spray contact with the crop and low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. See the label for additional information and warnings.</p> <p>-Rainfastness is 30 minutes. A maximum of 3 applications per year are allowed.</p>						

### 3. Postharvest

Group	Product Name	Product Rate	Active Ingredient (* = Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 2.0	2.25 to 3 pt/A	paraquat*	0.56 to 0.75 lb/A	--	24
<p>-<b>For postharvest desiccation of vegetable vines.</b> A Special Local-Needs 24(c) label has been approved for the use of Gramoxone SL 2.0 for postharvest desiccation of the crop in DE, NJ and VA.</p> <p>-Apply after the last harvest for bareground or plasticulture. Always include an adjuvant.</p> <p>-Spray coverage is essential for optimum effectiveness. See the label for additional information and warnings.</p> <p>-Rainfastness 30 minutes. A maximum of 2 applications for crop dessication are allowed.</p>						

**4. Other Labeled Herbicides** These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name	Active Ingredient (* = Restricted Use)
9	Roundup (various)	glyphosate
14	Aim EC	carfentrazone

## Insect Control

**THE LABEL IS THE LAW - See the Pesticide Use Disclaimer on page F 1.**  
**Recommended Insecticides**

## Seed Treatments for Seedcorn Maggot

See also the Pest Management chapter, Insect Management section. Control may be achieved by using commercially applied seed treatments containing chlorpyrifos (Lorsban 50W) or thiamethoxam (Farmore DI-400). **Note:** The use of neonicotinoid insecticides (Group 4A) at planting may help reduce seedcorn maggot populations.

**Aphids** **Note:** Aphids transmit multiple viruses.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV (melon aphid)	1.5 to 3.0 pt/A	methomyl*	3	48	H
3A + 4A	Endigo ZC	4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
3A + 6	Gladiator	19.0 fl oz/A	zeta-cypermethrin* + avermectin B1	7	12	H
4A	Admire PRO	7.0 to 10.5 fl oz/A	imidacloprid - <b>soil</b>	21	12	H
4A	Assail 30G	2.5 to 4.0 oz/A	acetamidprid	0	12	M
4A	Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - <b>soil</b>	21	12	H
4A	Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - <b>foliar</b>	7	12	H
4A	Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam - <b>soil</b>	30	12	H
4A	Actara 25WDG	1.5 to 3.0 oz/A	thiamethoxam - <b>foliar</b>	0	12	H
4A + 28	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole - <b>soil/drip</b>	30	12	H
4A + 28	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole - <b>foliar</b>	1	12	H
9B	Fulfill 50WP	2.75 oz/A	pymetrozine	0	12	L
9C	Beleaf 50SG	2.0 to 2.8 oz/A	flonicamid	0	12	L
28 + 6	Minecto Pro	10 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Cabbage Loopers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	3	48	H
3A	Baythroid XL	1.6 to 2.4 fl oz/A	beta-cyfluthrin*	0	12	H
3A	Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H
3A	Bifenture 2EC, Sniper	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H
3A	Danitol 2.4EC	10.67 to 16.0 fl oz/A	fenpropathrin*	7	24	H
3A	Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H
3A	Lambda-Cy, Lambda T CS	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
3A	Mustang Maxx	2.8 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H
3A	Perm-Up 3.2 EC	4.0 to 8.0 fl oz/A	permethrin*	0	12	H
3A	Tombstone	1.6 to 2.4 fl oz/A	cyfluthrin*	0	12	H
3A	Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
3A + 6	Gladiator	14.0 to 19.0 fl oz/A	zeta-cypermethrin* + avermectin B1	7	12	H
3A + 28	Voliam Xpress	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole	1	24	H
4A + 28	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole - <b>foliar</b>	1	12	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant EC	5.0 to 10.0 fl oz/A	spinetoram	1	4	H
11A	Dipel (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22A	Avaunt 30WDG	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - <b>soil/drip/foliar</b>	1	4	L
28 + 6	Minecto Pro	7.5 to 10 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Cucumber Beetles

Cucumber beetles can transmit bacterial wilt; however, losses from this disease vary greatly between fields and varieties. Pickling cucumbers grown in high-density rows for once-over harvesting can compensate for at least 10% stand losses. On farms with a history of bacterial wilt and where susceptible varieties are used, insecticides should be used to control adult beetles before they feed extensively on the cotyledons and first true leaves. If foliar insecticides are used, begin spraying shortly after plant emergence and repeat applications weekly if new beetles

## F Cucumbers

continue to invade fields. Treatments may be required until vines begin to run (usually about 3 weeks after plant emergence). Seeds pretreated with a neonicotinoid seed treatment such as Farmore DI-400 should provide up to 14 days of control of cucumber beetle, otherwise, apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	3	48	H
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Baythroid XL	2.4 to 2.8 fl oz/A	beta-cyfluthrin*	0	12	H
3A	Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H
3A	Bifenture 2EC, Sniper	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H
3A	Danitol 2.4EC	10.67 to 16.0 fl oz/A	fenprothrin*	7	24	H
3A	Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H
3A	Lambda-Cy, Lambda T	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
3A	Mustang Maxx	2.8 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H
3A	Perm-Up 3.2 EC	4.0 to 8.0 fl oz/A	permethrin*	0	12	H
3A	Tombstone	2.4 to 2.8 fl oz/A	cyfluthrin*	0	12	H
3A	Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
3A + 6	Gladiator	19.0 fl oz/A	zetacypermethrin* + avermectin B1	7	12	H
3A + 28	Voliam Xpress	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole	1	24	H
4A	Admire PRO	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
4A	Assail 30SG	2.5 to 5.3 oz/A	acetamiprid	0	12	M
4A	Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - soil	21	12	H
4A	Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - foliar	7	12	H
4A	Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil	21	12	H
4A	Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
4A	Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
4A	Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam - soil	30	12	H
4A	Actara 25WDG	3.0 to 5.5 oz/A	thiamethoxam - foliar	0	12	H
4A + 28	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole - soil	30	12	H
4A + 28	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole - foliar	1	12	H

## Cutworms - See also the Pest Management chapter, Insect Management section.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	3	48	H
3A	Baythroid XL	0.8 to 1.6 fl oz/A	beta-cyfluthrin*	0	12	H
3A	Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H
3A	Bifenture 2EC, Sniper	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H
3A	Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H
3A	Lambda-Cy, LambdaT	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
3A	Mustang Maxx	1.28 to 4.00 fl oz/A	zeta-cypermethrin*	1	12	H
3A	Perm-Up 3.2 EC	4.0 to 8.0 fl oz/A	permethrin*	0	12	H
3A	Tombstone	0.8 to 1.6 fl oz/A	cyfluthrin*	0	12	H
3A	Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
3A + 6	Gladiator	19.0 fl oz/A	zeta-cypermethrin* + avermectin B1	7	12	H
3A + 28	Voliam Xpress	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole	1	24	H

## Leafminers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate 2L	2.0 to 4.0 pt/A	oxamyl*	1	48	H
3A + 4A	Endigo ZC	4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
3A + 6	Gladiator	19.0 fl oz/A	zeta-cypermethrin* + avermectin B1	7	12	H

Leafminers continued on next page

## Leafminers - continued

3A + 28	Voliam Xpress	9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole	1	24	H
4A	Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - <b>soil</b>	21	12	H
4A	Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam - <b>soil</b>	30	12	H
4A	Actara	3.0 to 5.5 oz/A	thiamethoxam - <b>foliar</b>	0	12	H
4A	Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - <b>soil</b>	21	12	H
4A	Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - <b>foliar</b>	1	12	H
4A	Venom 70SG	5.0 to 6.0 oz/A	dinotefuran - <b>soil</b>	21	12	H
4A	Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - <b>foliar</b>	1	12	H
4A + 28	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole - <b>soil</b>	30	12	H
4A + 28	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole - <b>foliar</b>	1	12	H
5	Entrust 2SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant 1SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	H
6	Agri-Mek 0.7 SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	L
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - <b>soil/drip/foliar</b>	1	4	L
28 + 6	Minecto Pro	5.5 to 10 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Melonworms (MW), Pickleworms (PW)

When using foliar materials, make 1 treatment prior to fruit set, and then treat weekly. Check the label for additional instructions when using soil or drip applications. Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	3	48	H
1A	Sevin XLR Plus	0.5 to 1.0 qt/A	carbaryl	3	12	M
3A	Baythroid XL	1.6 to 2.4 fl oz/A	beta-cyfluthrin*	0	12	H
3A	Asana XL (pickleworm)	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H
3A	Bifenture 2EC, Sniper	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H
3A	Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H
3A	Lambda-Cy, Lambda T	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
3A	Mustang Maxx	2.8 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H
3A	Perm-Up 3.2 EC	4.0 to 8.0 fl oz/A	permethrin*	0	12	H
3A	Tombstone	1.6 to 2.4 fl oz/A	cyfluthrin*	0	12	H
3A	Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
3A + 6	Gladiator	19.0 fl oz/A	zeta-cypermethrin* + avermectin B1	7	12	H
3A + 28	Voliam Xpress	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole	1	24	H
4A + 28	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	H
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22A	Avaunt 30WDG	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	2.0 to 3.5 fl oz/A MW 3.0 to 7.5 fl oz/A PW	chlorantraniliprole - <b>drip/foliar</b>	1	4	L
28 + 6	Minecto Pro	5.5 to 10 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Mites

Mite infestations generally begin around field margins and grassy areas. **Do not mow or maintain field margins and grassy areas after midsummer since this forces mites into the crop.** Local infestations can be spot-treated. Begin treatment when 10-15% of the crown leaves are infested early in the season, or when 50% of the terminal leaves are infested later in the season. **Note:** Continuous use of carbaryl or a pyrethroid may result in mite outbreaks.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
3A + 6	Gladiator	19.0 fl oz/A	zeta-cypermethrin* + avermectin B1	7	12	H
6	Agri-Mek 0.7 SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L

Mites continued on next page

## F Cucumbers

Mites - continued

20D	Acramite 50WS	0.75 to 1.00 lb/A	bifenazate	3	12	M
21A	Portal	2.0 pt/A	fenpyroximate	1	12	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28 + 6	Minecto Pro	5.5-10 fl oz/A	cyantraniliprole + abamectin*	7	12	H

## Thrips

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate L	2.0 to 4.0 pt/A	oxamyl*	1	48	H
3A	Lambda-Cy, LambdaT	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
3A	Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
4A	Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam - soil	30	12	H
4A	Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil	21	12	H
4A	Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
4A	Venom 70SG	5.0 to 6.0 oz/A	dinotefuran - soil	21	12	H
4A	Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
4A + 28	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole - soil/drip	30	12	H
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	H

## Disease Control

**THE LABEL IS THE LAW - See the Pesticide Use Disclaimer on page F 1.**

### Recommended Pesticides

## Nematode Control

See also the Pest Management Chapter (Soil Fumigation and Nematodes sections), or apply one of the following:

Code	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate L	0.5 to 1.0 gal/A Incorporate into top 2-4 inches of soil, <b>OR</b> 2.0 to 4.0 pt/A apply 2 w after planting and repeat 2-3 w later.	oxamyl*	1	48	H
7	Velum Prime	6.5 to 6.84 fl oz/A	fluopyram	0	12	--
--	Nimitz 4EC	3.5 to 5.0 pt/A Incorporate or drip-apply 7 d before planting	fluensulfone	n/a	12	N

**Seed Treatment** Check if seed has been treated with an insecticide and fungicide. If it has not been treated, use a mixture of thiram 480DP (4.5 fl oz/100 lb seed) and an approved commercially available insecticide.

## Damping-Off caused by *Pythium*, *Phytophthora*, and *Rhizoctonia*

Code	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Apply one of the following at-planting (see label for application methods and restrictions):</b>						
<b>Phytophthora and Pythium root rot</b>						
4	Ridomil Gold 4SL	0.5 to 1.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4	MetaStar 2E	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
<b>Phytophthora, Pythium, and Rhizoctonia root rot</b>						
4 + 11	Uniform 3.66SE	0.34 fl oz/1000 ft row. Avoid direct seed contact, which may cause delayed emergence.	mefenoxam + azoxystrobin	AP	0	N
<b>Rhizoctonia root rot</b>						
11	azoxystrobin 2.08F	0.40 to 0.80 fl oz/1000 ft row	azoxystrobin	AP	4	N
<b>Pythium root rot only</b>						
28	Previcur Flex 6F	1.2 pt/A in transplant water, drip irrigation, or direct spray at base of plant and soil	Propamocarb HCL	2	12	N

## **Bacterial and Fungal Diseases**

### **Angular Leaf Spot**

At first sign of disease, apply the labeled rates of fixed copper plus mancozeb. Some coppers are OMRI-approved and can be used in organic production systems to help suppress Angular leaf spot and other fungal diseases. Repeat every 7 days. To minimize the spread of disease, avoid working in field while foliage is wet.

### **Anthraco**

Resistant varieties should be used when possible (see table Recommended Varieties). Begin fungicide applications when vines begin to run, or earlier if symptoms are detected. Alternate chlorothalonil or mancozeb with other effective fungicides every 7 days. Fungicides with a high risk for resistance development such as FRAC code 11 fungicides that do not come in a mix with another fungicide active ingredient that is effective on anthracnose, should be tank-mixed with a protectant fungicide. Use at least the minimum labeled rate of each fungicide in the tank-mix.

**Do not** apply FRAC code 11 fungicides more than 4 times total per season. **Do not** apply FRAC code 11 fungicides if resistance exists in the area; use fungicides with a different FRAC code instead.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Under LIGHT or MODERATE disease pressure ALTERNATE:</b>						
M5	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	L
<b>WITH a TANK MIX the following fungicide PLUS mancozeb 75DG 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A:</b>						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N
<b>Under HIGH disease pressure, TANK-MIX one of the following fungicides WITH chlorothalonil 6F 2.0 to 3.0 pt/A:</b>						
3 + 11	Quadris Top 2.7F	12.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
7 + 11	Merivon 500SC	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	0	12	N
7 + 11	Pristine 38WG	18.5 oz/A	pyraclostrobin + boscalid	0	12	--
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG	12.0 to 16.0 fl oz/A	pyraclostrobin	0	12	N
<b>AND ROTATE with a TANK-MIX of the following fungicide PLUS mancozeb 75DG 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A every 7 days</b>						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N

### **Bacterial Wilt**

Controlling striped and spotted cucumber beetles is essential for preventing bacterial wilt. See preceding "Cucumber Beetle" section under Insect Control for specific recommendations. Insecticide applications made at seeding may not prevent beetle damage all season; additional foliar insecticide applications may be necessary.

### **Belly Rot (*Rhizoctonia*)**

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Apply at the 1 to 3 leaf stage. Make a 2<sup>nd</sup> application 10-14 d later or just prior to vine tip-over (whichever occurs first):</b>						
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	1	4	N

### **Cottony Leak (*Pythium*)** - See also Damping off

At planting, apply mfenoxam (Ridomil Gold 4SL, Ultra Flourish 2E) or metalaxyl (MetaStar 2E).

### **Downy Mildew**

The pathogen does not overwinter, but introduction to the region can occur early in the year. Newly developed cultivars with resistance or tolerance should be planted where available (see table Recommended Varieties). Even when using resistant cultivars, a good fungicide program is important. However, fungicide efficacy may vary, as strains of the pathogen may vary between seasons.

Scout fields beginning at plant emergence. Begin sprays when vines run or earlier if disease occurrence is predicted for the region (check the Cucurbit Downy Mildew Forecasting website at <http://cdm.ipmpipe.org>). Once the disease has become established in an area, new plantings should receive an application of Ranman, or Previcur Flex at the 1-3 leaf stage. **Preventative applications are much more effective than applications made after disease is detected. In addition, spray programs that include fungicides with several different modes of action**

## F Cucumbers

**(FRAC codes) are more effective than programs with few modes of action.** For example, alternate Ranman (Code 21) *PLUS* Gavel (Codes M3 + 22), with Orondis Ultra (Codes U15 + 40) *PLUS* chlorothalonil (Code M5). Follow all fungicide label precautions in order to reduce the chance of resistance development.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>The following are the most effective products. Sprays should be applied on a 7-day schedule.</b>						
<b>Under severe disease conditions spray interval may be reduced IF the label allows.</b>						
<b>ALWAYS tank mix these products with a protectant fungicide (listed below):</b>						
U15+40	Orondis Ultra	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
21	Ranman 400SC	2.10 to 2.75 fl oz/A ( <b>do not</b> apply with copper; see label)	cyazofamid	0	12	L
28	Previcur Flex 6F	1.2 pt/A	propamocarb	2	12	N
43	Presidio 4SC	4.0 fl oz/A (caution: pathogen is now less sensitive to Presidio)	fluopicolide	2	12	L
M5 + 22	Zing! 4.9SC	36.0 fl oz/A	chlorothalonil + zoxamide	0	12	N
M5 + 27	Ariston 42SC	3.0 pt/A	chlorothalonil +cymoxanil	3	12	--
11 + 27	Tanos 50DF	8.0 oz/A	famoxadone + cymoxanil	3	12	--
27	Curzate 60DF	3.2 oz/A	cymoxanil	3	12	N
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N
40 + 45	Zampro 525SC	14.0 fl oz/A	acetetradin + dimethomorph	0	12	--
<b>TANK-MIX WITH protectant fungicides:</b>						
M3	mancozeb 75DG	3.0 lb/A	mancozeb	5	24	N
M3 + 22	Gavel 75DF	1.5 to 2.0 lb/A	mancozeb + zoxamide	5	48	--
M5	chlorothalonil 6F	1.5 to 3.0 pt/A	chlorothalonil	0	12	L

## Gummy Stem Blight

Gummy stem blight occurs primarily in the late summer. Fungicides with a high-risk for resistance development such as Pristine (FRAC code 11) should be tank-mixed with a protectant fungicide to reduce the chances for resistance development. Use at least the minimum labeled rate for each fungicide in the tank mix. **Do not** apply FRAC code 11 fungicides more than 4 times total per season. Apply fungicides from a different FRAC code if resistance to FRAC code 11 fungicides exists in the area. Begin sprays when vines begin to run.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>ALTERNATE one of the following formulations:</b>						
M3	mancozeb 75DG	2.0 to 3.0 lb/A	mancozeb	5	24	N
M5	chlorothalonil 6F	2.0 pt/A	chlorothalonil	0	12	L
<b>WITH A TANK-MIX containing either chlorothalonil or mancozeb PLUS one of the following fungicides:</b>						
3	Proline 480SC	5.7 fl oz/A	prothioconazole	7	12	L
3	tebuconazole 3.6 F	8.0 fl oz/A	tebuconazole	7	12	N
3 + 9	Inspire Super 2.8F	16.0 to 20.0 fl oz/A	difenoconazole + cyprodinil	0	12	--
3 + 7	Luna Experience 3.34SC	10.0 to 17.0 fl oz/A	fluopyram + tebuconazole	7	12	--
7	Fontelis 1.67SC	12.0 to 16.0 fl oz/A	penthiopyrad	1	12	L
9 + 12	Switch 62.5WG	11.0 to 14.0 oz/A	cyprodinil + fludioxonil	1	12	N
3 + 11	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	N
3 + 11	TopGuard EQ	5.0 to 8.0 fl oz/A	azoxystrobin + flutriafol	1	12	--
7 + 11	Merivon 500SC	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	7	12	N
7 + 11	Pristine 38WG	12.5 to 18.5 oz/A	pyraclostrobin + boscalid	0	12	--
11	azoxystrobin 2.08F <sup>1</sup>	11.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG <sup>1</sup>	12.0 to 16.0 oz/A	pyraclostrobin	0	12	N

<sup>1</sup>azoxystrobin 2.08F and Cabrio 20EG are not recommended in MD, DE and VA due to resistance development.

## Phytophthora Crown and Fruit Rot

Different strategies should be used to minimize the occurrence of this disease. Rotate away from susceptible crops (such as cucurbits, peppers, lima and snap beans, eggplants, and tomatoes) for as long as possible, improve field drainage, and apply preplant fumigants. When conditions favor disease development apply fungicides following excellent resistance management practices. Fungicides provide suppression only.

*Phytophthora Crown and Fruit Rot continued on next page*

*Phytophthora* Crown and Fruit Rot - continued

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Apply one of the following fungicides. Rotate fungicides with different FRAC codes and tank mix with a fixed copper (exception: do not tank mix Ranman 400SC with copper).</b>						
U15 +40	Orondis Ultra	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
40	Revus 2.08F	8.0 fl oz/A	mandipropamid	0	4	--
40 + 45	Zampro 525SC	14.0 fl oz/A	acetochradin + dimethomorph	0	12	--
43	Presidio 4SC <sup>1</sup>	3.0 to 4.0 fl oz/A	fluopicolide	2	12	L
M3 + 22	Gavel 75DF	1.5 to 2.0 lb/A	mancozeb + zoxamide	5	48	--
11 + 27	Tanos 50DF	8.0 to 10.0 oz/A	famoxadone + cymoxanil	3	12	--
21	Ranman 400SC	2.75 fl oz/A (plus a non-ionic or organosilicon surfactant; see label for additional precautions)	cyazofamid	0	12	L
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N

<sup>1</sup>Presidio may also be applied through the drip irrigation (see supplemental label). Soil drench followed by drip application has given good results in some trials on crown rot caused by *Phytophthora capsici*.

**Powdery Mildew**

Excellent resistance is available (see table Recommended Varieties). The fungus that causes cucurbit powdery mildew has developed resistance to high-risk fungicides. In the Eastern US, resistance to strobilurin (FRAC code 11), DMI (FRAC code 3), and SDHI (FRAC code 7) fungicides has been reported. Proper fungicide resistance management should be followed to help delay the development of resistance and minimize control failures.

Powdery mildew generally occurs from mid-July until the end of the season. Observe plants for the presence of powdery mildew. If one lesion is found on the underside of 45 old leaves/A, begin the following fungicide program:

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>TANK MIX one of these products with a protectant such as chlorothalonil 6F at 2.0 to 3.0 pt/A:</b>						
U6	Torino 0.85SC	3.4 fl oz/A	cyflufenamid	0	4	--
U8	Vivando 2.5SC	15.4 fl oz/A	metafenone	0	12	--
3 + 7	Luna Experience 3.34SC	10.0 to 17.0 fl oz/A	tebuconazole + fluopyram	7	12	--
<b>AND ALTERNATE with a TANK MIX of one of the following and a protectant such as chlorothalonil 6F at 2.0 to 3.0 pt/A</b>						
3	Proline 480SC	5.7 fl oz/A	prothioconazole	7	12	--
3	Procore 480SC	4.0 to 8.0 fl oz/A	triflumizole	0	12	N
3	Rally 40WSP	5.0 oz/A <i>PLUS</i>	myclobutanil	0	24	N
3	tebuconazole 3.6F	4.0 to 6.0 fl oz/A	tebuconazole	7	12	N
3 + 9	Inspire Super 2.8F	16.0 to 20.0 fl oz/A	difenoconazole + cyprodinil	0	12	--
3 + <b>11</b>	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	N
3 + <b>11</b>	TopGuard EQ	5.0 to 8.0 fl oz/A	azoxystrobin + flutriafol	1	12	--
7	Fontelis 1.67SC	12.0 to 16.0 fl oz/A	penthiopyrad	1	12	L
7 + <b>11</b>	Pristine 38WG	12.5 to 18.5 oz/A	pyraclostrobin + boscalid	0	12	--

**Scab**

Scab typically occurs during cool periods. Excellent resistance is available in some varieties and they should be used when possible.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
<b>Apply one of the following as true leaves form and repeat every 5-7 days:</b>						
M3	mancozeb 75DG	2.0 to 3.0 lb/A	mancozeb	5	24	N
M5	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	L

**Viruses**

The most prevalent virus in the mid-Atlantic region is WMV2, followed by PRSV, ZYMV and CMV. Use varieties with multiple virus resistance when possible (see table Recommended Varieties). Plant fields far away from existing cucurbit plantings to help reduce aphid transmission of viruses into new fields.



**For Immediate Medical Attention**

**Call 911**

**For a Pesticide Exposure Poisoning  
Emergency Call**



**For All States**

This number will automatically connect you to the poison center nearest you.

**Anyone with a poisoning emergency can call the toll-free telephone number for help.** Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

### **For Pesticide Spills**

**Small Spills:** See the product label for cleanup advice.

**Large spills:** Call the National Response Center at 1-800-424-8802 or CHEMTREC at 800-424-9300 (24 hours) - Industry assistance with emergency response cleanup procedures for large, dangerous spills.

**Be aware of your responsibility to report spills to the proper state agency.**