



This is a section from the

2020/2021

Mid-Atlantic

Commercial Vegetable

Production Recommendations

The recommendations are **NOT** for home gardener use.

The **full manual**, containing recommendations specific to New Jersey, can be found on the Rutgers NJAES website in the Publications section: <http://njaes.rutgers.edu/pubs/publication.asp?pid=E001>.

This manual will be revised biennially. In January 2021, a **critical update** with important updates to the 2020/2021 manual will be communicated through local Extension Agents and Vegetable Specialists.

The **label** is a legally-binding contract between the user and the manufacturer. The user must follow all rates and restrictions as per label directions. The use of any pesticide inconsistent with the label directions is a violation of Federal law.

Cooperating Agencies: Rutgers, The State University of New Jersey, U.S. Department of Agriculture, and County Boards of Chosen Freeholders. Rutgers Cooperative Extension, a unit of the Rutgers New Jersey Agricultural Experiment Station, is an equal opportunity program provider and employer.

F. Commodity Recommendations

Pesticide Use Disclaimer

THE LABEL IS THE LAW

Before using a pesticide, check the label for up to date rates and restrictions.

Labels can be downloaded from: <http://www.cdms.net/>, <https://www.greenbook.net/>
or <http://www.agrian.com/labelcenter/results.cfm>

For more information on Pesticide Safety and the Pesticide Label see chapter D.

Guide to the Recommended Pesticide Tables in the Following Crop Sections:

- 1. Pesticides are listed by group or code number based on chemical structure and mechanism of action**, as classified by the Weed Science Society of America (WSSA) for herbicides, the Insecticide Resistance Action Committee (IRAC) for insecticides, and the Fungicide Resistance Action Committee (FRAC) for fungicides.
If the number is in bold font, the product may have resistance concerns.
- 2. For restricted use pesticides**, the restricted active ingredients are labeled with a *.
(See section D 3.2.1 “Restricted Use Classification Statement” for more information).
- 3. In addition to the pesticides listed below, other formulations or brands with the same active ingredient(s) may be available. ALWAYS CHECK THE LABEL:**
 - a) to ensure a pesticide is labeled for the same use,**
 - b) to ensure the pesticide is labeled for the desired crop, and**
 - c) for additional restrictions.**
- 4. All pesticide recommendations are made for spraying a broadcast area of 1 acre** (43,560 square feet). **Adjust the rate for banded applications** (for more information, see section E 1.3 Calibrating Granular Applicators).
- 5. Check the label for the maximum amount of pesticide per application and the maximum number of applications per year.**
- 6. Bee Toxicity Rating (Bee TR):** N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.

F Cucumbers

Labeled Applications Sites for Cucumbers									
		Plastic mulch production					Bare-ground production		
		Soil-Applied		Postemergence			Soil-applied	POST	Post-harvest
Herbicides	WSSA group number	Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest			
Sandea	2	YES	YES	YES	YES		YES	YES	
Treflan	3		YES						
Curbit	3		YES				YES		
Prefar	8	YES	YES				YES		
Command	13		YES				YES		
Strategy	3+13		YES				YES		
Select	1			YES	YES			YES	
SelectMax	1			YES	YES			YES	
Poast	1			YES	YES			YES	
Gramoxone*	22					YES		YES	

*Special Local Needs Label 24(c), be sure it is registered for the specific state and for the intended use.

1. Soil-Applied						
Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	14	12
<p>-Plasticulture: can be applied in a band under the plastic, immediately before laying the mulch; delay seeding or transplanting for 7 days after application. Row row middles: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v or include a non-selective herbicide.</p> <p>-Bareground: apply broadcast after seeding but before crop emergence or no sooner than 7 days before transplanting.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field. Do not apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum Sandea applications per year is 2 and do not exceed 2 oz/A during the crop season.</p>						
3	Curbit 3EC	1 to 3 pt/A	ethalfuralin	0.38 to 1.13 lb/A	--	24
<p>-Plasticulture row middles only: apply as a banded spray after crop emergence or transplanting. Do not soil incorporate.</p> <p>-Bareground: apply broadcast after direct-seeding but prior to crop emergence; do not use on transplanted cucumbers.</p> <p>-Controls annual grasses and certain annual broadleaf weeds, including carpetweed and pigweed sp.</p> <p>-Use lower rate for coarse-textured soils or soils with low organic matter.</p> <p>-Where overhead irrigation is available, activate Curbit with 0.5 inch of irrigation within 2 days after application; if no irrigation or rainfall occurs within 5 days of application, activity of Curbit can be reduced.</p> <p>-Available as a pre-mix herbicide Strategy. Strategy at 3 pt/A= Curbit at 26 fl oz (0.6 lb ai) and Command at 8 fl oz (0.188 lb ai)</p> <p>-Maximum applications per season: not specified</p>						
3	Treflan 4EC	1 to 2 pt/A	trifluralin	0.5 to 1 lb/A	30	12
<p>-Plasticulture row middles only: apply as a directed spray after emergence when plants have reached the 3 to 4 true leaf stage of growth.</p> <p>-Not labeled for bareground production. Primarily controls annual grasses with a few broadleaf weeds. -Do not use (or reduce the rate) when cold, wet soil conditions are expected, or crop injury may result. -Maximum applications per season: not specified.</p>						
3 + 13	Strategy 2.1SC	1.5 to 6 pt/A	ethalfuralin plus clomazone	0.39 to 1.58 lb/A	45	24
<p>-Plasticulture: row middles application. Bareground: apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Strategy is a prepackage mixture of Curbit 3EC and Command 3ME. Refer to individual products for comments.</p> <p>-Clomazone spray or vapor drift may injure susceptible crops and other vegetation, refer to Command 3ME for comments.</p> <p>-Do not apply prior to planting crop. Do not soil incorporate. Maximum applications per season: not specified.</p>						
8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	45	12
<p>-Plasticulture: under plastic: apply in a band under the plastic, immediately before laying the mulch. Allow 7 day before making transplant holes to allow condensation to incorporate the herbicide. Plasticulture: row middles application is labeled.</p> <p>-Bareground: apply preemergence or preplant incorporated.</p> <p>-Preemergence applications should be followed by irrigation within 36 h (apply enough water to wet the soil at least 2 to 4 inches deep). Preplant incorporated applications should be incorporated 1 to 2 inches deep (deeper than 2 inches will result in reduced weed control).</p> <p>-Provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters.</p> <p>-Do not apply more than 6 lb ai/A per season.</p>						
13	Command 3ME	0.4 to 1 pt/A	clomazone	0.015 to 0.375 lb/A	45	12
<p>Plasticulture: row middles application only. -Bareground: apply broadcast just before planting or after planting but before crop emergence. -Supplemental labeling reduces PHI to 30 days (label expires 12/10/2021).</p>						

1. Soil-Applied, Command - continued on next page

1. Soil-Applied, Command - continued

-Use the lower rate when used on coarse-textured soils low in organic matter, when weed pressure is light, or to minimize herbicide carryover that could affect subsequent crops.

-Controls annual grasses and many broadleaf weeds including common lambsquarters, velvetleaf, spurred anoda, and jimsonweed. Carpetweed, morningglory sp., pigweed sp., and yellow nutsedge will not be controlled. Higher rates will improve control (or expand number of species controlled) such as common cocklebur, common ragweed, or jimsonweed (refer to label for specific weeds and rates).

-**WARNINGS:** Command spray or vapor drift may injure sensitive crops and other vegetation up to several hundred yards from the point of application. **Do not** apply adjacent to sensitive crops (see label) or vegetation, or under unfavorable wind or weather conditions. Command may limit subsequent cropping options, see the label. -Available as a pre-mix herbicide Strategy: Strategy at 3 pt/A= Command at 8 fl oz (0.188 lb ai) and Curbit at 26 fl oz (0.6 lb ai) -Maximum number of Command applications per year: 1.

2. Postemergence

Group	Product Name	Product Rate	Active Ingredient (* = Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC Select Max 0.97EC	6 to 8 fl oz/A 9 to 16 fl oz/A	clethodim	0.094 to 0.13 lb/A	14	24
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.19 to 0.28 lb/A	3	12

-**Select 2EC:** use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). **Select Max:** use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). **Poast:** Apply with COC at 1.0% v/v. -**The use of COC may increase the risk of crop injury when hot or humid conditions prevail.** To reduce the risk of crop injury, omit additives or switch to NIS when grasses are small and soil moisture is adequate. -Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.

-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled.

-Controls many annual and certain perennial grasses, including annual bluegrass, but Poast is preferred for goosegrass control. For best results, treat annual grasses when they are actively growing and before tillers are present. Control may be reduced if grasses are large or under hot or dry weather conditions.

-Repeated applications may be necessary to control certain perennial grasses. If repeat applications are necessary, allow 14 days between applications.

-**Do not** tank-mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses.

-**Do not** apply more than 8 fl oz of Select 2EC in a single application and **do not** exceed 32 fl oz/A for the season; **do not** apply more than 16 fl oz of Select Max in a single application and **do not** exceed 64 fl oz/A for the season.

-**Do not** apply more than 1.5 pt/A Poast in single application and **do not** exceed 3 pt/A for the season. -Rainfastness is 1 h.

2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	14	12
---	-------------	---------------	--------------	---------------------	----	----

-**Plasticulture:** broadcast (over the top) or directed to row middles; broadcast for bareground.

-**Bareground:** apply Sandea after the crop has at least 3 to 5 true leaves but before first female flowers appear and no sooner than 14 days after transplanting. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v.

-Suppresses or controls yellow nutsedge and certain broadleaf; control of weeds taller than 3 inches may not be adequate. Sandea will not control common lambsquarters or eastern black nightshade if applied postemergence; for row middle application, tankmix with a non-selective herbicide to increase spectrum of control.

-Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. **Do not** use Group 2 herbicides repeatedly in the same field.

-**Do not** apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.

-Rainfastness is 4 h. Sandea applications per year is 2 and **do not** exceed 2 oz/A during the crop season

22	Gramoxone 2SL	1.95 pt/A	paraquat*	0.49 lb/A	14	24
----	---------------	-----------	-----------	-----------	----	----

-**A Supplemental Label has been approved for the use of Gramoxone 2SL for postemergence weed control in DE, MD, NJ, PA, and VA.** Row middles as a shielded application. Apply as a directed spray in a minimum of 20 gal spray mix/A to control emerged weeds between the rows after crop establishment. Include a nonionic surfactant at 0.25% v/v.

-Use shields or hoods to prevent spray contact with the crop and low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. See the label for additional information and warnings.

-Rainfastness is 30 min. A maximum of 3 applications per year are allowed.

-**Restricted-use pesticide.** Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<http://usparaquattraining.com>); certified applicators must repeat training every three years.

3. Postharvest

Group	Product Name	Product Rate	Active Ingredient (* = Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 2.0	2.25 to 3 pt/A	paraquat*	0.56 to 0.75 lb/A	--	24

-**A Special Local Needs Label 24(c) has been approved in VA (expires 12/31/2022) and a Supplemental Label in DE for the use of Gramoxone SL 2.0 for postharvest application to desiccate the crop.**

-Apply after the last harvest for bareground or plasticulture. Always include an adjuvant.

3. Postharvest, Gramoxone, continued on next page

F Cucumbers

3. Postharvest, Gramoxone, continued

-Spray coverage is essential for optimum effectiveness. See the label for additional information and warnings.
 -Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed.
 -**Restricted-use pesticide.** Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (<http://usparaquattraining.com>); certified applicators must repeat training every three years.

4. Other Labeled Herbicides These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name	Active Ingredient (*=Restricted Use)
14	Aim	carfentrazone

Insect Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Insecticides

Seed Treatments for Seedcorn Maggot

Control may be achieved by using commercially applied seed treatments containing chlorpyrifos (Lorsban 50W) or thiamethoxam (Farmore DI-400). **Note:** The use of neonicotinoid insecticides (Group 4A) at planting may help reduce seedcorn maggot populations. See also Maggots in section E 3.1 Soil Pests - Detection and Control.

Aphids Note: Aphids transmit multiple viruses.

Apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl* - melon aphid only	1-3	48	H
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil	21	4	M
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
29	Beleaf 50SG	2.0 to 2.8 oz/A	flonicamid	0	12	L

Armyworms and Cabbage Loopers

Apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
11A	Dipel DF, others (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
11A	XenTari (OMRI) (armyworms)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
11A	XenTari (OMRI) (cabbage loopers)	0.5 to 1.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel (armyworms)	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H

Armyworms and Cabbage Loopers - continued on next page

Armyworms and Cabbage Loopers - continued

28	Exirel (cabbage loopers)	10.0 to 17.0 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Cucumber Beetles

Cucumber beetles can transmit bacterial wilt; however, losses from this disease vary greatly between fields and varieties. Pickling cucumbers grown in high-density rows for once-over harvesting can compensate for at least 10% stand losses. On farms with a history of bacterial wilt control adult beetles before they feed extensively on the cotyledons and first true leaves. If foliar insecticides are used, begin spraying shortly after plant emergence and repeat weekly if new beetles continue to invade fields. Treatments may be required until vines begin to run (usually about 3 weeks after plant emergence). Seeds pretreated with a neonicotinoid seed treatment such as Farmore DI-400 should provide up to 14 days of control of cucumber beetle, otherwise, apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
28	Exirel	20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Cutworms See also section E 3.1. Soil Pests - Detection and Control.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV (variegated cutworm)	1.5 pt/A	methomyl*	1	48	H
1A	Lannate LV (granulate cutworm)	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					

Leafminers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	M
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Melonworms and Pickleworms

Apply one of the following formulations. When using foliar materials, make one treatment prior to fruit set, and then treat weekly. Check the label for additional instructions when using soil or drip applications.						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
1A	Sevin XLR Plus	0.5 to 1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	1	4	M

Melonworms and Pickleworms - continued on next page

F Cucumbers

Melonworms and Pickleworms - continued

Group	Product Name	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	1	4	M
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	2.0 to 3.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole	30	12	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Mites

Mite infestations generally begin around field margins and grassy areas. **Do not mow or maintain field margins and grassy areas after midsummer since this forces mites into the crop.** Local infestations can be spot-treated. Begin treatment when 10-15% of the crown leaves are infested early in the season, or when 50% of the terminal leaves are infested later in the season. **Note:** Continuous use of carbaryl or a pyrethroid may result in mite outbreaks.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L
20B	Kanemite 15SC	31.0 fl oz/A	acequinocyl	1	12	L
21 A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
21A	Portal XLO	2.0 pt/A	fenpyroximate	1	12	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M

Thrips

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Cucumbers: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	1	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	1	4	M
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Group 3A Pyrethroid Insecticides Registered for Use on Cucumbers

Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):						
Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR	
Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H	
Baythroid XL	0.8 to 2.8 fl oz/A	beta-cyfluthrin*	0	12	H	
Bifenthrin 2EC, others	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H	
Danitol 2.4EC	10.67 to 16.0 fl oz/A	fenpropathrin*	7	24	H	
Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H	
Lambda-Cy 1EC, others	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H	
Mustang Maxx	1.28 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H	
Permethrin 3.2EC, others	4.0 to 8.0 fl oz/A	permethrin*	0	12	H	
Tombstone, others	0.8 to 2.8 fl oz/A	cyfluthrin*	0	12	H	
Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H	
Combo products containing a pyrethroid						
Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam (Group 4A)	1	24	H	
Gladiator	19.0 fl oz/A	zeta-cypermethrin* + abamectin* (Group 6)	7	12	H	
Besiege	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole (Group 28)	1	24	H	

Group 4A Neonicotinoid Insecticides Registered for Use on Cucumbers					
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):					
Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Actara 25WDG	1.5 to 5.5 oz/A	thiamethoxam	0	12	H
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H
Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
Assail 30SG	2.5 to 5.3 oz/A	acetamiprid	0	12	M
Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - soil/drip	21	12	H
Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - foliar (note: PHI: do not make application after 4 th true leaf has unfolded)	see note	12	H
Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil/drip	21	12	H
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
Venom 70SG	5.0 to 7.5 oz/A	dinotefuran - soil/drip	21	12	H
Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
Combo products containing a neonicotinoid					
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H
Endigo ZC	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin* (Group 3A)	1	24	H
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole (Group 28)	1	12	H

Disease Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F.
Recommended Fungicides

Nematode Control

See also the chapter E Pest Management (sections E 1.5 Soil Fumigation and E 1.6 Nematode Control), or apply one of the following:

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate L	0.5 to 1.0 gal/A Incorporate into top 2-4 inches of soil, OR 2.0 to 4.0 pt/A apply 2 w after planting and repeat 2-3 w later.	oxamyl*	1	48	H
7	Velum Prime 4.16SC	6.5 to 6.84 fl oz/A	fluopyram	0	12	--
--	Nimitz 4EC	3.5 to 5.0 pt/A Incorporate or drip-apply 7 d before planting	fluensulfone	n/a	12	N

Seed Treatment Check if seed has been treated with an insecticide and fungicide. If it has not been treated, use a mixture of thiram 480DP (4.5 fl oz/100 lb seed) and an approved commercially available insecticide.

Damping-Off caused by *Pythium*, *Phytophthora*, and *Rhizoctonia*

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply one of the following at-planting (see label for application methods and restrictions):						
Phytophthora and Pythium root rot						
4	Ridomil Gold 4SL	0.5 to 1.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4	MetaStar 2E AG	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
Phytophthora, Pythium, and Rhizoctonia root rot						
4 + 11	Uniform 3.66SE	0.34 fl oz/1000 ft row. Avoid direct seed contact, which may cause delayed emergence.	mefenoxam + azoxystrobin	AP	0	N
Rhizoctonia root rot						
11	azoxystrobin 2.08F	0.40 to 0.80 fl oz/1000 ft row	azoxystrobin	AP	4	N
Pythium root rot only						
28	Previcur Flex 6F	1.2 pt/A in transplant water, drip irrigation, or direct spray at base of plant and soil	propamocarb HCl	2	12	N

Bacterial and Fungal Diseases

Angular Leaf Spot

Resistant varieties should be used when possible (see table Recommended Varieties). At first sign of disease, apply the labeled rates of fixed copper plus mancozeb. Some coppers are OMRI-approved and can be used in organic production systems to help suppress Angular leaf spot and some fungal diseases. Repeat every 7 days. To minimize the spread of disease, avoid working in field while foliage is wet.

Anthracnose

Resistant varieties should be used when possible (see table Recommended Varieties). Begin fungicide applications when vines begin to run, or earlier if symptoms are detected. Alternate chlorothalonil or mancozeb with other effective fungicides every 7 days. Fungicides with a high risk for resistance development such as FRAC code 11 fungicides that do not come in a mix with another fungicide active ingredient that is effective on anthracnose, should be tank-mixed with a protectant fungicide. Use at least the minimum labeled rate of each fungicide in the tank-mix. **Do not** apply FRAC code 11 fungicides more than 4 times total per season. **Do not** apply FRAC code 11 fungicides if resistance exists in the area; use fungicides with a different FRAC code instead.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Under LIGHT or MODERATE disease pressure ALTERNATE:						
M05	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	N
WITH a TANK MIX the following fungicide PLUS mancozeb 75DF 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A:						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N
Under HIGH disease pressure, TANK-MIX one of the following fungicides WITH chlorothalonil 6F 2.0 to 3.0 pt/A:						
3 + 11	Quadris Top 1.67SC	12.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
7 + 11	Merivon 2.09SC	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	0	12	N
7 + 11	Pristine 38WG	18.5 oz/A	boscalid + pyraclostrobin	0	12	--
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG	12.0 to 16.0 fl oz/A	pyraclostrobin	0	12	N
AND ROTATE with a TANK-MIX of the following fungicide PLUS mancozeb 75DF 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A every 7 days						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N

Bacterial Wilt

Controlling striped and spotted cucumber beetles is essential for preventing bacterial wilt. See preceding "Cucumber Beetle" section under Insect Control for specific recommendations. Insecticide applications made at seeding may not prevent beetle damage all season; additional foliar insecticide applications may be necessary.

Belly Rot (*Rhizoctonia*)

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply at the 1 to 3 leaf stage. Make a 2nd application 10-14 d later or just prior to vine tip-over (whichever occurs first):						
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	1	4	N

Cottony Leak (*Pythium*) - See also Damping off

At planting, apply mefenoxam (Ridomil Gold 4SL, Ultra Flourish 2E) or metalaxyl (MetaStar 2E AG).

Downy Mildew

The pathogen does not overwinter, but introduction to the region can occur early in the year. Newly developed cultivars with resistance or tolerance should be planted where available (see table Recommended Varieties). Even when using resistant cultivars, a good fungicide program is important. However, fungicide efficacy may vary, as strains of the pathogen may vary between seasons.

Scout fields beginning at plant emergence. Strains of downy mildew that infect one cucurbit crop may not affect cucumber. Unnecessary fungicide application can be avoided by not spraying until disease is predicted in the region on cucumber. Begin sprays when vines run or earlier if disease occurrence is predicted for the region (check the Cucurbit Downy Mildew Forecasting website at <http://cdm.ipmpipe.org>). Once the disease has become established in an area, new plantings should receive an application of Ranman, or Previcur Flex at the 1-3 leaf stage. **Preventative applications are much more effective than applications made after disease is detected. In**

For Immediate Medical Attention

Call 911

**For a Pesticide Exposure Poisoning
Emergency Call**



For All States

This number will automatically connect you to the poison center nearest to you.

Anyone with a poisoning emergency can call the toll-free telephone number for help. Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

For Pesticide Spills

Small Spills: See the product label for cleanup advice.

Large spills: Call the National Response Center at 1-800-424-8802 or CHEMTREC at 800-424-9300 (24 hours) - Industry assistance with emergency response cleanup procedures for large, dangerous spills.

Be aware of your responsibility to report spills to the proper state agency.