



This is a section from the

**2026/2027**

# **Mid-Atlantic Commercial Vegetable Production Recommendations**

The recommendations are **NOT** for home gardener use.

The **full manual**, containing recommendations specific to New Jersey, can be found on the Rutgers NJAES website in the Publications section at: <https://njaes.rutgers.edu/pubs/publication.php?pid=e001>.

This manual will be revised biennially. **In January 2027, a Critical Update** with important updates to the 2026/2027 manual will be communicated through local Extension Agents and Vegetable Specialists.

The **label** is a legally-binding contract between the user and the manufacturer. The user must follow all rates and restrictions as per label directions. The use of any pesticide inconsistent with the label directions is a violation of federal law.

**Cooperating Agencies:** Rutgers, The State University of New Jersey, U.S. Department of Agriculture, and County Boards of Commissioners. Rutgers Cooperative Extension, a unit of the Rutgers New Jersey Agricultural Experiment Station, is an equal opportunity program provider and employer.

**2026/2027**  
**Mid-Atlantic Commercial Vegetable**  
**Production Recommendations**



**If you are having a medical emergency after using pesticides, always **call 911** immediately.**



**In Case of an Accident**

- Remove the person from exposure
- Get away from the treated or contaminated area immediately
- Remove contaminated clothing
- Wash with soap and clean water
- Call a physician and/or the National Poison Control Center (1-800-222-1222).  
Your call will be routed to your State Poison Control Center.
- **Have the pesticide label with you!**
- Be prepared to give the EPA registration number to the responding center/agency

# Preface

## NOT TO BE USED BY HOME GARDENERS

**This copy of the 2026/2027 Mid-Atlantic Commercial Vegetable Production Recommendations replaces all previous editions of the Commercial Vegetable Production Recommendations published individually for Delaware, Maryland, New Jersey, Pennsylvania, Virginia, and West Virginia.** Information presented in this publication is based on research results from the University of Delaware, the University of Maryland, Rutgers - The State University of New Jersey, The Pennsylvania State University, Virginia Polytechnic Institute and State University, West Virginia University, Delaware State University, Cornell University, the U.S. Department of Agriculture, and other institutions, combined with industry and grower knowledge and experience.

**This publication will be revised biennially. In January 2027, a Critical Update with important updates for this publication will be communicated through local Extension Agents and Vegetable Specialists.** The editors welcome constructive criticism and suggestions from growers and industry personnel who may wish to help improve future editions of this publication.

**These recommendations are intended for the commercial vegetable grower** who has to make numerous managerial decisions. Although the proper choices of variety, pesticides, equipment, irrigation, fertilizer, and cultural practices are the individual vegetable grower's responsibility, it is intended that these recommendations will facilitate decision-making. Recommended planting dates will vary across the six-state region. Local weather conditions, grower experience, and variety may facilitate successful harvest on crops planted outside the planting dates listed in this guide. This can be evaluated in consultation with the local agents and state specialists. Government agencies and other organizations administering crop insurance programs or other support programs should contact local Extension agents and/or vegetable specialists for guidance.

## Disclaimer

- The label is a legally-binding contract between the user and the manufacturer.
- The user **MUST** follow all rates and restrictions as per label directions.
- The use of any pesticide inconsistent with the label directions is a violation of Federal law.

## Pesticide User Responsibility

Always follow the label and use pesticides safely. For Special Local Needs Label 24(c) registrations or Section 18 exemptions, do not use the material without a copy of the special label or written instructions from your Extension Agent or another recognized authority. **The user is always responsible for the proper use of pesticides, residues on crops, storage, and disposal, as well as for damage caused by drift.**

State and federal pesticide regulations are constantly under revision. Be sure to determine if such changes apply to your situation. Using pesticides inconsistent with label directions is illegal.

## Days Between Last Application and Harvest

The minimum number of days between the last application and harvest (**PHI**, Pre-Harvest Interval, in days) and reentry information (**REI**, Restricted Entry Interval, in hours) are listed in the herbicide, insecticide and fungicide recommendation tables in Chapter F Commodity Recommendations. Always follow the label to avoid the occurrence of deleterious chemical residues on harvested crops.

## Trade or Brand Names

The trade or brand names given herein are supplied with the understanding that no discrimination is intended, and no endorsement is implied. Furthermore, in some instances the same compound may be sold under different trade names, which may vary as to label clearances. For the convenience of our users, both product names and active ingredients are provided, and any product name omissions are unintended.

# Coordinators and Editors

## 2026/2027 Mid-Atlantic Commercial Vegetable Production Recommendations

### Coordinators

**C.A. Wyenandt, Ph.D.**

Extension Specialist in Vegetable Pathology (Rutgers University)

**M.M.I. van Vuuren, Ph.D.** (Rutgers University)

### Discipline Editors

#### **Entomology**

**D. Owens, Ph.D.** (University of Delaware)

#### **Horticulture**

**E. Sánchez, Ph.D.** (Penn State)

#### **Pesticide Safety**

**D.L. Frank, Ph.D.** (Virginia Tech)

#### **Plant Pathology**

**C.A. Wyenandt, Ph.D.** (Rutgers University)

#### **Weed Science**

**M.J. VanGessel, Ph.D.** (University of Delaware)

### Front Cover Illustration

**K.E. Chasalow**

# Contributors

These recommendations were prepared and reviewed by the following individuals from respective institutions with the purpose of providing up to date information for commercial vegetable growers in the Mid-Atlantic states of Delaware, Maryland, New Jersey, Pennsylvania, Virginia, and West Virginia.

## Cornell University

### **Weed Science**

L. Sosnoskie

## University of Delaware

### **Entomology**

D. Owens

### **Horticulture**

E. Ernest

### **Pathology**

A.M. Betts

### **Weed Science**

M.J. VanGessel

## University of Maryland

### **Entomology**

K.A. Hamby

V. Yurchak

E. Zobel

### **Pathology**

M.J. Hu

### **Weed Science**

K. Vollmer

## University of Massachusetts

### Amherst

### **Weed Science**

M. Gannett

## Penn State University

### **Entomology**

H.L. Grabb

### **Horticulture**

K. Demchak

F. Di Gioia

L. Duque

T.E. Elkner

L. Fronk

E. Sánchez

K. Snyder

### **Weed Science**

D. Lingenfelter

J.M. Wallace

## Rutgers University

### **Entomology**

M. Cramer

A. Quadrel

### **Horticulture**

M. Infante-Casella

W. Kline

M. Melendez

P. Nitzsche

### **Pathology**

C.A. Wyenandt

### **Soil Science**

J. R. Heckman

### **Weed Science**

T. Besançon

## Virginia Tech

### **Entomology**

T.P. Kuhar

K.L. Sutton

J.M. Wilson

### **Horticulture**

J.B. Samtani

### **Pathology**

D. Higgins

D.B. Langston

### **Pesticide Safety**

D.L. Frank

### **Soil Science**

M.S. Reiter

### **Weed Science**

V. Singh

### **Wildlife Conservation**

J.A. Parkhurst

## West Virginia University

### **Entomology**

C.R. Quesada

### **Horticulture**

L.W. Jett

### **Pathology**

M.M. Rahman

### **Weed Science**

R.S. Chandran

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

# State Extension Information

## DELAWARE

### University of Delaware

University of Delaware Cooperative Extension: <https://www.udel.edu/canr/cooperative-extension/>

Mid-Atlantic Commercial Vegetable Production Recommendations: publication EB137

Vegetable Program Trial Reports, Publications, and Budgets:

<https://www.udel.edu/academics/colleges/canr/cooperative-extension/sustainable-production/commercial-crops/vegetable-crops/>

Weekly Crop Update Newsletter: <https://sites.udel.edu/weeklycropupdate/>

Delaware Insect Pest Management, Trial Reports, Insect Trapping Program:

<https://www.udel.edu/academics/colleges/canr/cooperative-extension/sustainable-production/pest-management/>

UD Plant Diagnostic Clinic: <https://www.udel.edu/academics/colleges/canr/cooperative-extension/sustainable-production/plant-diagnostic-clinic/>

University of Delaware Weed Science:

<https://www.udel.edu/academics/colleges/canr/cooperative-extension/sustainable-production/weed-science/>

### Statewide

Emmalea Ernest, Extension Specialist, Vegetables & Fruit, 302-856-7303, [emmalea@udel.edu](mailto:emmalea@udel.edu)

Mark VanGessel, Extension Specialist, Weed Science, 302-856-7303, [mjv@udel.edu](mailto:mjv@udel.edu)

Alyssa Betts, Extension Specialist, Plant Pathology, 302-856-7303, [akoehler@udel.edu](mailto:akoehler@udel.edu)

David Owens, Extension Specialist, Entomology, 302-698-7125, [owensd@udel.edu](mailto:owensd@udel.edu)

Jill Pollok Plant Diagnostician, 302-831-1390, [jillp@udel.edu](mailto:jillp@udel.edu)

### County Offices

New Castle County: Carrie Murphy, 302-831-2506, [cjmurphy@udel.edu](mailto:cjmurphy@udel.edu)

Kent County: Drew Harris, 302-730-4000, [raharris@udel.edu](mailto:raharris@udel.edu)

Sussex County: Tracy Wootten, 302-856-7303, [wootten@udel.edu](mailto:wootten@udel.edu)

Sussex County: Cory Whaley, 302-856-7303, [whaley@udel.edu](mailto:whaley@udel.edu)

## MARYLAND

### University of Maryland

Maryland Vegetables: <https://extension.umd.edu/mdvegetables>

University of Maryland Plant Diagnostic Lab: <https://extension.umd.edu/plantdiagnosticlab>

University of Maryland Fruit and Vegetable News: <https://extension.umd.edu/programs/agriculture-food-systems/program-areas/fruit-vegetable-production/vegetable-fruit-news>

Extensión en Español: <https://extensionesp.umd.edu/>

University of Maryland Extension Agriculture & Food Systems Directory:

<https://agmr.umd.edu/about/directory/agriculture-food-systems>

### Statewide

Kurt Vollmer, Weed Management, [kvollmer@umd.edu](mailto:kvollmer@umd.edu)

Veronica Yurchak, Vegetable Production and Entomology, [vjohnso4@umd.edu](mailto:vjohnso4@umd.edu)

Mengjun Hu, Small Fruit Disease Management, [mjhu@umd.edu](mailto:mjhu@umd.edu)

Ana Cristina Fulladolsa, Plant Diagnostician, 301-405-0730, [acfulla@umd.edu](mailto:acfulla@umd.edu)

# State Extension Information

## NEW JERSEY

### Rutgers, The State University of New Jersey

New Jersey Agricultural Experiment Station: <https://njaes.rutgers.edu/ag/>

Fact Sheets and Bulletins: <https://extension.rutgers.edu/fact-sheets>

Plant & Pest Advisory: <https://plant-pest-advisory.rutgers.edu>

Mid-Atlantic Commercial Vegetable Production Recommendations: publication E001  
<https://njaes.rutgers.edu/pubs/publication.php?pid=e001>

Rutgers Pest Management Office: <https://pestmanagement.rutgers.edu/>

### Rutgers NJAES Extension Specialists and IPM Program Personnel

For a complete listing see: <https://njaes.rutgers.edu/extension-specialists/>

Thierry Besançon (Weed Science), A.J. Both (Controlled Environmental Engineering),

Joseph Heckman (Soil Fertility), Amanda Quadrel (Vegetable

IPM Senior Program Coordinator), Maria Cramer (Vegetable IPM Senior Program Coordinator),

Jim Simon (Plant Biology), C.A. Wyenandt (Vegetable Pathology)

### Rutgers Cooperative Extension County Agricultural Agents/Program Associates\* with Vegetable Responsibilities

For a complete listing of county offices and agents see: <https://njaes.rutgers.edu/county/>

Atlantic County, Patrick McMullen, 609-625-0056, [pwm38@njaes.rutgers.edu](mailto:pwm38@njaes.rutgers.edu)

Burlington County, David Hlubik, 609-265-5050, [djh222@njaes.rutgers.edu](mailto:djh222@njaes.rutgers.edu)

Cape May County, Jennifer Sawyer\*, 609-465-5115, [sawyer@njaes.rutgers.edu](mailto:sawyer@njaes.rutgers.edu)

Cumberland County, Wesley Kline, 856-451-2800, [wkline@njaes.rutgers.edu](mailto:wkline@njaes.rutgers.edu)

Gloucester County, Michelle Infante-Casella, 856-224-8040, [minfante@njaes.rutgers.edu](mailto:minfante@njaes.rutgers.edu)

Mercer County, Meredith Melendez, 609-989-6830, [melendez@njaes.rutgers.edu](mailto:melendez@njaes.rutgers.edu)

Monmouth County, William Errickson, 732-431-7260, [william.errickson@njaes.rutgers.edu](mailto:william.errickson@njaes.rutgers.edu)

Morris County, Peter Nitzsche, 973-285-8304, [nitzsche@njaes.rutgers.edu](mailto:nitzsche@njaes.rutgers.edu)

Somerset County, Kate Brown, 908-526-6293, [kbrown@njaes.rutgers.edu](mailto:kbrown@njaes.rutgers.edu)

### Rutgers Vegetable Working Group Directory

For complete listing and contact information see:

<https://nj-vegetable-crops-online-resources.rutgers.edu/> and go to “members” section.

### Rutgers NJAES Plant Diagnostic Lab and Nematode Detection Service

<https://njaes.rutgers.edu/plantdiagnosticlab/>; [rutgerspdl@njaes.rutgers.edu](mailto:rutgerspdl@njaes.rutgers.edu)

20 Indyk-Engel Way, North Brunswick, NJ 08902, phone: 732-932-9140, fax: 732-932-1270

### Rutgers Soil Testing Lab <https://extension.rutgers.edu/soil-testing-lab>

57 US Highway 1, New Brunswick, NJ 08901, phone: 848-932-9295, [soiltest@njaes.rutgers.edu](mailto:soiltest@njaes.rutgers.edu)

### Rutgers NJAES Pesticide Safety Education Program

<https://pestmanagement.rutgers.edu/PAT> (pesticide manuals, record forms, regulations, and safety)

<https://pestmanagement.rutgers.edu/worker-protection/> (WPS training videos and regulations),

phone: 848-932-9802

### Rutgers NJAES OCPE Pesticide Applicator Certification Exam Registration (PACER)

<https://pacer.rutgers.edu/index.php>, phone: 848-932-9271 Option 7; email: [pacer@njaes.rutgers.edu](mailto:pacer@njaes.rutgers.edu)

# State Extension Information

## PENNSYLVANIA

### The Pennsylvania State University

Penn State Extension, including publications, fact sheets, and more: <https://extension.psu.edu>

Mid-Atlantic Commercial Vegetable Production Recommendations: publication AGRS-028:

<https://extension.psu.edu/mid-atlantic-commercial-vegetable-production-recommendations-sections> and

<https://extension.psu.edu/catalog/product/view/id/21333/s/commercial-vegetable-production-recommendations/>

Penn State Vegetable Production: <https://extension.psu.edu/forage-and-food-crops/vegetables> (click on the “News” button to see recent articles related to vegetable and small fruit production)

### Penn State Vegetable Team Directory

For a complete listing and contact information of Extension offices throughout Pennsylvania, see:

<https://extension.psu.edu/county-offices>

**Plant Disease Clinic:** <https://plantpath.psu.edu/about/facilities/plant-disease-clinic>

220 Buckhout Laboratory, University Park, PA 16802, phone: 814-865-2204 [PlantClinic@psu.edu](mailto:PlantClinic@psu.edu)

### Insect Identification Laboratory:

<https://ento.psu.edu/about/contact/instructions-for-insect-identification-requests>, 501 Agricultural Sciences and Industries Building, University Park, PA 16802, phone: 814-865-1895 [entomology@psu.edu](mailto:entomology@psu.edu)

**Agricultural Analytical Services Laboratory:** <https://agsci.psu.edu/aasl>,

111 Ag Analytical Services Lab, University Park, PA 1682, phone: 814-863-0841, [aaslab@psu.edu](mailto:aaslab@psu.edu)

## VIRGINIA

### Virginia Tech & Virginia State University

Virginia Cooperative Extension (VCE): <https://ext.vt.edu/>

VCE Publications and Educational Resources: <https://ext.vt.edu/agriculture.html> and Mid-Atlantic Commercial

Virginia Tech Pesticide Programs (VTPP): <https://vtpp.ento.vt.edu/>

### Virginia Extension Specialists

Douglas S. Higgins, Assistant Professor, Plant Pathology, 757-807-6584, [doughiggins@vt.edu](mailto:doughiggins@vt.edu)

Thomas P. Kuhar, Professor, Entomology, 540-231-6129, [tkuhar@vt.edu](mailto:tkuhar@vt.edu)

David B. Langston, Jr., Professor, Plant Pathology, 757-657-6450, [dblankston@vt.edu](mailto:dblankston@vt.edu)

James A. Parkhurst, Associate Professor, Wildlife, 540-231-9283, [jbparkhur@vt.edu](mailto:jbparkhur@vt.edu)

Mark S. Reiter, Professor, Soils and Nutrient Management, 757-807-6576, [mreiter@vt.edu](mailto:mreiter@vt.edu)

Steve L. Rideout, Professor, Plant Pathology, 540-231-3197, [srideout@vt.edu](mailto:srideout@vt.edu)

Jayesh B. Samtani, Associate Professor, Small Fruit, 757-363-3901, [jsamtani@vt.edu](mailto:jsamtani@vt.edu)

Vijay Singh, Associate Professor, Weed Science and Precision Agriculture, 757-807-6579, [vijaysingh@vt.edu](mailto:vijaysingh@vt.edu)

Laura K. Strawn, Professor, Food Science, 540-315-4925, [lstrawn@vt.edu](mailto:lstrawn@vt.edu)

Kemper L. Sutton, Assistant Professor, Entomology, 757-807-6586, [klsutton@vt.edu](mailto:klsutton@vt.edu)

James Wilson, Apiculturist, 540-231-2168, [keepbees@vt.edu](mailto:keepbees@vt.edu)

**Plant Disease Clinic** <https://spes.vt.edu/affiliated/plant-disease-clinic.html>

Virginia Tech, 106 Price Hall, 170 Drillfield Drive, Blacksburg, VA 24061-0331

Phone: 540-231-6758, [clinic@vt.edu](mailto:clinic@vt.edu)

Or contact the local VCE: <https://ext.vt.edu/offices.html>

**Insect Identification Lab** <https://www.ento.vt.edu/idlab.html>

205A Price Hall, 170 Drillfeld Drive, Virginia Tech Blacksburg, VA 24061-0319

Phone: 540-231-4899, [insectidlab@vt.edu](mailto:insectidlab@vt.edu)

Or contact the local VCE: <https://ext.vt.edu/offices.html>

# State Extension Information

## WEST VIRGINIA

### West Virginia University Extension Service:

<https://extension.wvu.edu/agriculture/horticulture>

### West Virginia University Extension Specialists

Lewis W. Jett, Ph.D., Extension Horticulture Specialist, 304-293-2634, [Lewis.Jett@mail.wvu.edu](mailto:Lewis.Jett@mail.wvu.edu)

Rakesh S. Chandran, Ph.D., Extension Weed Specialist and IPM Coordinator, 304-293-2603,

[RSChandran@mail.wvu.edu](mailto:RSChandran@mail.wvu.edu)

Mahfuz M. Rahman, Ph.D., Extension Plant Pathology Specialist, 304-293-8838, [MM.Rahman@mail.wvu.edu](mailto:MM.Rahman@mail.wvu.edu)

Carlos R. Quesada, Ph.D., Extension Entomology Specialist, 304-293-8835, [carlos.quesada@mail.wvu.edu](mailto:carlos.quesada@mail.wvu.edu)

### Plant Diagnostic Clinic

<https://extension.wvu.edu/lawn-gardening-pests/plant-disease/plant-diagnostic-clinic>

G102 South Ag. Sciences Building, PO Box 6108, Morgantown, WV 26506-6108

Phone: 304-293-8838/288-9541, [mm.rahman@mail.wvu.edu](mailto:mm.rahman@mail.wvu.edu)

### Soil Testing Lab

<https://soiltesting.wvu.edu>

1309-B Agricultural Sciences Bldg., PO Box 6108, Morgantown, WV 26506-6108,

Phone: 304-293-6023, [infoplantsoil@mail.wvu.edu](mailto:infoplantsoil@mail.wvu.edu)

# Table of Contents

<b>Preface</b>	<b>i</b>
<b>Coordinators and Editors</b>	<b>ii</b>
<b>Contributors</b>	<b>iii</b>
<b>State Extension Information</b>	<b>iv</b>
<b>Table of Contents</b>	<b>viii</b>
<b>Tables and Figures</b>	<b>x</b>
<b>Abbreviations and Acronyms</b>	<b>xii</b>

<b>Chapter</b>	<b>Section</b>	<b>Title</b>	<b>Page</b>
<b>A</b> <b>General Production Recommendations</b>	<b>1</b>	Varieties	1
	<b>2</b>	Seed Storage and Handling	2
	<b>3</b>	Specialty Vegetables	3
	<b>4</b>	Organic Production	7
	<b>5</b>	Transplant Production	8
	<b>6</b>	Conservation Tillage Crop Production (No-Till, Strip-Till)	13
	<b>7</b>	Mulches and Row Covers	14
	<b>8</b>	Staking and Trellising	17
	<b>9</b>	High Tunnels	17
	<b>10</b>	Greenhouse Production	20
	<b>11</b>	Wildlife Damage Prevention	21
	<b>12</b>	Pollination	30
	<b>13</b>	Food Safety Concerns	36
<b>B</b> <b>Soil and Nutrient Management</b>	<b>1</b>	Soils	39
	<b>2</b>	Liming Soils	39
	<b>3</b>	Plant Nutrients	44
	<b>4</b>	Nutrient Management	47
	<b>5</b>	Soil Improvement and Organic Nutrient Sources	55
<b>C</b> <b>Irrigation Management</b>	<b>1</b>	Basic Principles	60
	<b>2</b>	Drip (Trickle) Irrigation	62
	<b>3</b>	Fertigation	67
	<b>4</b>	Subsurface Drip Irrigation Systems	68
	<b>5</b>	Chemigation	69
<b>D</b> <b>Pesticide Safety</b>	<b>1</b>	General Information	70
	<b>1.1</b>	Pesticide Registration	70
	<b>1.2</b>	Pesticides and Food Safety	70
	<b>2</b>	Certification of Pesticide Applicators	71
	<b>3</b>	The Pesticide Label	72
	<b>3.1</b>	Labels and Labeling	72
	<b>3.2</b>	Label Statements	73
	<b>3.2.1</b>	Restricted Use Pesticide Classification Statement	73
	<b>3.2.2</b>	Signal Words	73
	<b>3.2.3</b>	First Aid Statements	75
	<b>3.2.4</b>	Other Label Statements	76
	<b>3.3</b>	Significant Labeling Changes	77
	<b>3.3.1</b>	Soil Fumigants	77
	<b>3.3.2</b>	Paraquat Dichloride (Paraquat)	77
<b>3.3.3</b>	Chlorpyrifos	78	

Table of Contents - continued next page

*Table of Contents - continued*

<b>Chapter</b>	<b>Section</b>	<b>Title</b>	<b>Page</b>
<b>D</b> <b>Pesticide Safety</b> <i>- continued</i>	<b>4</b>	Handling Pesticides	78
	<b>4.1</b>	Prior to Pesticide Application	78
	<b>4.2</b>	Pesticide Application and Recordkeeping	79
	<b>4.3</b>	Pesticide Transport	81
	<b>4.4</b>	Pesticide Storage	81
	<b>4.5</b>	Disposal of Pesticides	82
	<b>4.6</b>	Disposal of Pesticide Containers	83
	<b>4.7</b>	Pesticide Spills	85
	<b>5</b>	Reducing Risks to Handlers and Workers	86
	<b>5.1</b>	EPA's Worker Protection Standard	86
	<b>5.2</b>	Personal Protective Equipment for Pesticides	89
	<b>5.2.1</b>	Body Protection for Early Entry Workers and Pesticide Handlers	90
	<b>5.2.2</b>	Respiratory Protection for Pesticide Handlers	91
	<b>6</b>	Protection of the Environment	92
	<b>6.1</b>	Minimize Off-Target Drift	92
	<b>6.2</b>	Protection of Surface and Ground Water	93
	<b>6.3</b>	Protection of Non-Target Organisms	95
	<b>6.3.1</b>	Protection of Pollinators	95
	<b>6.3.2</b>	Protection of Endangered Species	99
	<b>7</b>	Mid-Atlantic Region Pesticide Program Contacts	102
	<b>E</b> <b>Pest Management</b>	<b>1</b>	How to Improve Pest Management
<b>1.1</b>		Recommendations for More Effective Pest Control	104
<b>1.2</b>		Calibrating Field Sprayers	108
<b>1.3</b>		Calibrating Granular Applicators	109
<b>1.4</b>		Pesticide Drift and Misapplication	110
<b>1.5</b>		Soil Fumigation	111
<b>1.6</b>		Nematode Control	111
<b>2</b>		Weed Control	115
<b>2.1</b>		Postharvest Perennial Weed Control	115
<b>2.2</b>		Herbicide Mode of Action	115
<b>2.3</b>		Herbicide Effectiveness on Common Weeds in Vegetables	118
<b>2.4</b>		Weed Species Prone to Resistance and of Concern in the Mid-Atlantic Region	121
<b>2.5</b>		Crop Rotation Planting Restrictions	122
<b>2.6</b>		Prepackaged Herbicide Mixtures	133
<b>3</b>		Insect Control	135
<b>3.1</b>		Soil Pests - Detection and Control	135
<b>3.2</b>		Insecticide Mode of Action	138
<b>3.3</b>		Insect Pest and Mite Control for Greenhouse Production	138
<b>3.4</b>		Insect Pest and Mite Control for Chemigation	142
<b>3.5</b>		Other Labeled Formulations for Insecticides	142
<b>3.6</b>		Biological or Organic Insecticides	144
<b>4</b>		Disease Control	146
<b>4.1</b>		Fungicide Mode of Action	146
<b>4.2</b>		How Biofungicides Work	146
<b>4.3</b>		Disease Control in Seeds, Plant Growing Mix, and Plant Beds	147
<b>4.4</b>		Fungicides Registered for Vegetables	148
<b>4.5</b>		Disease Control for Greenhouse Production	155

*Table of Contents - continued next page*

*Table of Contents - continued*

Chapter	Section	Title	Page
<b>F</b> <b>Commodity Recommendations</b>		Pesticide Use Disclaimer	160
		Asparagus	161
		Beans (Snap and Lima)	171
		Beets (Garden)	186
		Carrots	191
		Celery	198
		Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, and Kohlrabi	206
		Cucumbers	223
		Edamame	238
		Eggplant	245
		Garlic	255
		Greens (Mustard, Turnip)	262
		Horseradish	269
		Leeks	275
		Lettuce, Endive and Escarole	280
		Muskmelons and Mixed Melons	289
		Okra	304
		Onions	310
		Parsley	319
		Parsnips	325
		Peas (Succulent)	329
		Peppers	337
		Potatoes	354
		Pumpkins and Winter Squash	368
		Radishes, Rutabagas and Turnips	383
		Spinach	390
	Strawberries	397	
	Summer Squash	416	
	Sweet Corn	429	
	Sweet Potatoes	446	
	Tomatoes	453	
	Watermelons	476	
<b>G</b> <b>Resources</b>	<b>1</b>	Vegetable Seed Sizes	494
	<b>2</b>	Plant Spacing and Populations	494
	<b>3</b>	Frequently Used Weights and Measures	495
	<b>4</b>	Making a Plant-Growing Mix	495

## Tables and Figures

Figure	Title	Page
B-1	Nutrient Application Rates Vary in Relation to Soil Test Category	44
B-2	Petiole Delineation for Several Plant Species	52
B-3	Changes in Soil Test Levels over Time under Different Nutrient Management Scenarios	54
E-1	Droplet Size and Surface Coverage, and Trade-off Between Spray Coverage and Drift Reduction	106

*Tables and Figures - continued next page*

*Tables and Figures - continued*

<b>Table</b>	<b>Title</b>	<b>Page</b>
A-1	World Vegetable Crops for Mid-Atlantic Growers	4
A-2	Baby and Miniature Vegetable Varieties and Harvest Stage	5
A-3	Potherbs and Salad Greens	6
A-4	Temperature and Planting Recommendations for Transplant Production	9
A-5	Planting and Harvesting Schedule for Freestanding High Tunnel Vegetable Crop Production	20
A-6	State Apiarist Contact Information	31
B-1	Target Soil pH Values for Vegetable Crops	40
B-2	Pounds of Calcium Carbonate Equivalent (CCE) Recommended per Acre	43
B-3	Conversion of Recommended Calcium Carbonate Equivalent to Recommended Limestone	43
B-4a,b	Soil Test Categories for Nutrients Extracted by Mehlich 3 and 1	45
B-5	Composition of Principal Macronutrient Fertilizer Materials	49
B-6	Chemical Sources of Secondary and Micronutrients	49
B-7	Boron Recommendations Based on Soil Tests for Vegetable Crops	51
B-8	Recommendations for Correction of Vegetable Crop Nutrient Deficiencies	53
B-9	Sufficiency Ranges for Fresh Petiole Sap Concentrations in Vegetable Crops	53
B-10	Plant Nutrient Value Credits to Be Allowed for Manure Applications and Crop Residues	58
B-11	Status for Organic Production, Mineral Nutrient Value, and Relative Availability of Various Materials	58
C-1	Most Critical Periods of Water Needs by Crops	60
C-2	Available Water-Holding Capacity Based on Soil Texture	61
C-3	Soil Infiltration Rates Based on Soil Texture	61
C-4	Irrigation Applied per Hour per Cropped Acre (Inches)	63
C-5	Maximum Number of Hours per Application for Drip Irrigated Vegetables	63
C-6	Irrigation Guidelines for Tensiometers	64
C-7	Equivalent Injection Proportions	67
D-1	EPA Signal Words According to Toxicity Categories (I, II, III, IV) of Pesticide Products	74
E-1	Ground Speed Conversion	108
E-2	Important Herbicide Groups for Commercial Vegetables	116
E-3	Herbicide Effectiveness on Common Weeds in Vegetables	118
E-4	Weed Species Prone to Resistance and of Concern in the Mid-Atlantic Region	121
E-5	Crop Rotation Planting Restrictions: Months After Herbicide Application Until Planting New Crop	122
E-6	Prepackaged Herbicide Mixtures	133
E-7	Insecticides and Miticides Labeled for Use on Greenhouse Vegetables	138
E-8	Insecticides with Labels for Chemigation	142
E-9	Other Labeled Formulations for Insecticides	142
E-10	Biological or Organic Insecticides	144
E-11	FRAC Codes and Corresponding Chemical Groups for Commonly Used Fungicides	146
E-12	Effective Seed Treatment Temperature Protocols (2 <sup>nd</sup> Bath) for Pathogen Eradication	148
E-13	Commonly Used Fungicides Registered for Vegetables	149
E-14	Selected Fungicides Labeled for Greenhouse Use	155
G-1	Vegetable Seed Sizes	494
G-2	Plant Spacing and Populations	494
G-3	Frequently Used Weights and Measures	495
G-4	Simple Plant-Growing Mix	496
G-5	Preferred Plant-Growing Mix	496

# Abbreviations and Acronyms

## Units of Measurement

/A	per acre
bu	bushel(s)
°C	degrees Celsius
cc	cubic centimeter(s)
cu ft	cubic foot (feet)
cu yd	cubic yard(s)
cwt	hundredweight
d	day(s)
°F	degrees Fahrenheit
ft	foot (feet)
fl oz	fluid ounce(s)
g	gram(s)
gal	gallon(s)
gpm	gallons per minute
h	hour(s)
in	inch(es)
lb	pound(s)
min	minute(s)
mph	miles per hour
oz	ounce(s)
ppm	parts per million
psi	pounds per square inch
pt	pint(s)
qt	quart(s)
sq ft	square foot (feet)
tbs	tablespoon(s)
tsp	teaspoon(s)
wk	week(s)
yr	year(s)

## Product Formulations

A	acid
CS	capsulated suspension
D	dust
DF	dry flowable
DP	dry prill
DS	dry salt
E	emulsion
EC	emulsifiable concentrate
ES	emulsifiable suspension
EW	emulsion in water
F	flowable
FC	flowable concentrate
FL	fluid
FM	flowable micro-encapsulated
G	granule

## Product Formulations (*continued*)

L	liquid
LC	liquid concentrate
LF	liquid flowable
LFR	liquid fertilizer ready
LV	low volatile
ME	micro-encapsulated
OF	oil formulation
SC	spray concentrate, soluble concentrate
SE	suspoemulsion
SG	soluble granules
SL	soluble liquid
SP	soluble powder
W	wettable
WBE	water-based emulsion
WDG	water-dispersible granules
WDL	water-dispersible liquid
WP	wettable powder
WS	water soluble
WSB	water-soluble bag
WSP	water-soluble packet
ZC	CS and SC mixture

## Other

ae	acid equivalent
ai	active ingredient
AMS	ammonium sulfate
AP	at planting
COC	crop oil concentrate
FRAC	Fungicide Resistance Action Committee
HRAC	Herbicide Resistance Action Committee
IRAC	Insecticide Resistance Action Committee
K	potassium
K <sub>2</sub> O	available potash
MoA	mode of action
MSO	methylated seed oil
N	nitrogen
NIS	non-ionic surfactant
OLF	other labeled formulations
OMRI	Organic Materials Research Institute
P	phosphorus
P <sub>2</sub> O <sub>5</sub>	available phosphoric acid
PHI	Pre-Harvest Interval (in days)
REI	Restricted Entry Interval (in hours)
TR	Toxicity Rating
WSSA	Weed Science Society of America

# A. General Production Recommendations

## 1. Varieties

New varieties are constantly being developed throughout the world, and it is impossible to list and describe them all; only those that are available and adapted to the Mid-Atlantic region are listed in this publication (see crop sections in Chapter F). These lists of varieties are based on research trials at universities in the Mid-Atlantic, our observations of what farmers are growing, and conversations with seed company representatives. They are not comprehensive lists. Check with your state Extension organization for science-based recommendations when available. Farmers also select varieties by talking with other farmers and seed company representatives and going to trade shows. We recommend evaluating new varieties on a small scale before planting them in a large area. The ultimate value of a variety for a particular purpose is determined by the farmer based on performance under their management, adaptation to site-specific environmental conditions, as well as having desired horticultural characteristics for markets.

### Some Variety Selection Criteria:

**Yield:** The variety should have the potential to produce crops at the same or better yield and quality as those already grown. Harvested yield may be much less than potential yield depending on markets and quality factors.

**Days to Harvest:** Choose varieties that meet market requirements based on days to harvest. Earliness is a major selection factor for first spring plantings, and days to harvest are a critical selection factor for late summer and fall maturing crops, especially in shorter-season areas of the region. Days to harvest in seed guides are based on the most common planting date and may be considerably longer in cooler periods or shorter in warmer periods. A more accurate guide to maturity will be Growing Degree Days (GDD), which are calculated for a specific crop using daily highs and lows and a base temperature. GDD information for different vegetable crops, such as peas and sweet corn, is available from seed suppliers and breeders.

**Disease and Insect Resistance, Herbicide Resistance:** The most economical and effective means of pest management is through the use of varieties that are resistant or tolerant to diseases caused by fungi, bacteria, viruses, or nematodes. When all other factors are equal, select a variety with needed disease resistance or tolerance. In some vegetables, such as sweet corn, insect resistant varieties are also available and should be considered where they fit your requirements. Herbicide resistant varieties of sweet corn are also available to allow for the use of post emergence non-selective herbicides for weed control. The continuous or intense production of herbicide or pest-resistant varieties can potentially lead to herbicide-tolerant weeds and new, more virulent pest strains. Adherence to venter or Extension recommendations and a long-term crop rotation plan should minimize this risk.

**Resistance to Adverse Environmental Conditions:** Choose varieties that are resistant to environmental conditions that are likely to be encountered. This includes heat or cold tolerance (bolting or disorders such as tuber heat necrosis or frost tolerance); drought tolerance; resistance to wet weather (disorders such as cracking and edema); and resistance to nutrient-related disorders such as blossom end rot, leaf tip burn, and hollow stem.

**Horticultural Quality:** Choose varieties that meet produce quality requirements for your market. Quality attributes such as taste, texture, size, shape, color, uniformity, and defects often dictate variety selection. Grades, percentages by grade, or pack-outs are key quality attributes for some markets. Variety test data such as soluble solids (sugars or sweetness), acidity, pungency, fiber content, and consumer preference information can assist in variety selection where available. Processing performance is a major concern for frozen, canned, or pickled vegetables. Other considerations include the ability to handle mechanical harvest or the ability to be packed and shipped distances with minimum damage in contrast to vegetables that are adapted only to hand harvest and local sales or short-distance shipping. Other quality characteristics to consider include holding or storage ability, ripening characteristics, nutritional content, and culinary qualities.

## A. General Production Recommendations

**Plant Characteristics:** Plant characteristics that may be considered in variety selection include plant form such as bush, upright, or vining; plant height; plant size; location of harvested part on the plant (such as top set in beans); and ease of harvest.

**Adaptability:** Successful varieties must perform well under the range of environmental conditions and production practices commonly encountered on individual farms. Seasonal adaptation is another selection consideration. Note that varieties listed under the “Recommended Varieties” section for individual crops in Chapter F may not be adapted to all areas of the region (*e.g.*, mountain vs. coastal growing areas).

**Market Acceptability:** The harvested plant product must have characteristics desired by both you and your buyers. Consider the requirements or desires of consumers, packers, shippers, wholesalers, retailers, or processors. Included among these qualities are flavor, pack out, size, shape, color, culinary qualities, nutritional quality, or processing quality. Specialty markets such as ethnic markets, restaurants, or gourmet sales will have very specific variety requirements. Many vegetable seed companies offer varieties that are transgenic or “GMO” (genetically modified organisms). GMO varieties feature a small amount of DNA from a source outside of the crop species gene pool: another plant species, bacterium, virus, or even an animal. This foreign DNA is either the direct source of a new trait such as herbicide, disease, or insect resistance or is needed to assist the gene insertion process. GMO products in the food chain are of concern to a portion of consumers and buyers. Be aware of potential adverse public sentiment before growing and marketing GMO varieties of vegetable crops.

**Variety selection is a very dynamic process.** Some varieties retain favor for many years, whereas others might be used only a few seasons. Companies frequently replace varieties with newer ones. In the Mid-Atlantic, variety selection often requires special regional consideration due to the wide range of climatic variations. **There are many sources of information for farmers to aid in choosing a variety.** University trials offer unbiased comparisons of varieties from multiple sources. Commercial trials from seed distributors also offer multiple source comparisons. Seed company test results offer information about that company’s varieties. Check results from replicated trials and multiple sites, if available. Trials conducted in similar soils and environments and local trials are the most reliable indicators of what will have the potential to perform well on your farm. Visits to local trials can provide good visual information for making decisions. Where quality is a prime concern, look for trials with quality data. Small-scale trial plantings for 2 to 3 years on your farm are suggested for any variety or strain not previously grown. For a true comparison, always include a standard variety, one with proven consistent performance in the same field or planting.

### Plant Resistance or Tolerance Listed in Tables

If a specific disease (or insect) is a serious threat to a vegetable crop, genetic resistance is an effective and often low-cost strategy for disease avoidance. Pathogens are highly changeable, and a resistant variety that performs well in one year may not necessarily continue to do so. There are cases where purported resistance to pathogens breaks down. This may be due to genetic shifts in the pathogen, the development of different strains and races of disease-causing organisms, or environmental conditions that favor the organism or reduce natural plant resistance. In Chapter F, variety tables, disease and insect resistances and tolerances are listed in the tables and footnotes. The disease, insect, or insect reactions listed in this book are from seed companies or university trials, as noted, and are not necessarily verified by Cooperative Extension.

## 2. Seed Storage and Handling

Both high temperature and high relative humidity will reduce seed germination and vigor over time. Do not store seeds in areas that have a combined temperature and humidity value greater than 110, for example 50°F + 60% relative humidity. Ideal storage conditions for most seeds are at a temperature of 35°F (2°C) and less than 40% relative humidity. In addition, primed seeds pretreated with salt or another osmoticum do not usually store well after shipment to the buyer. Seed coating/pelleting may or may not reduce germination rate. When storing coated/pelleted seed, perform a germination test to assess viability before using in subsequent seasons.

Corn, pea, and bean seeds are especially susceptible to mechanical damage due to rough handling. Seed containers of these crops should not be subjected to rough handling since the seed coats and embryos can be damaged, resulting in nonviable seeds. If you plan to treat seeds of these crops with a fungicide, inoculum, or other chemical application, apply the materials gently to avoid seed damage.

## 3. Specialty Vegetables

### Niche Marketing

The term ‘specialty vegetables’ refers to a broad range of crops that are sometimes called ‘exotic’ as they represent a class of vegetables unlike standard tomatoes, peppers, beans, peas, sweet corn, etc.; ‘alternative’ because they represent new enterprises that traditional vegetable growers might try; or ‘designer veggies’ that allow the consumer to be creative with their presentation. Recently, the term ‘ultra-niche crops’ was created to describe very high value specialty crops that provide opportunities to help beginner/small farmers get established. Most fresh-market specialty vegetables and herbs fit this description.

Specialty vegetables can be described as **new or unusual manner in which they are grown** (aeroponic, hydroponic); by the **color, shape or flavor** of the varieties grown (red and oakleaf lettuces, pear tomatoes, heirloom varieties, or unusual greens like radicchio); by their **size** (baby, miniature, micro); or by their **diverse origins and demand** (Asian crucifers and cucurbits, Hispanic peppers, African greens and eggplants).

### Specialty Vegetable Markets

Developing a marketing plan is important for any business and is essential to success with specialty vegetables. Important points to consider include:

- Before planting, know where you will be selling your crop.
- Understand all the quality, grading and packaging requirements, and costs for various market outlets (similar ethnic groups may want different varieties/types of the same crop, use the same/similar names for different types of crops, or different names for the same crop).
- Determine consumers will want it *when* you can produce it.
- Assess the costs of production, especially the time and labor required. Conduct on-farm trials to help determine varieties and production systems. Small plantings can help work out problems that can be resolved easily. Accurate records of small plantings can be used to estimate costs and returns for full-scale plantings.
- Increase production as demand grows but be aware of competitors entering the market (prospective buyers, state and federal crop reporting agencies, and local Extension workers can be good sources of information).
- Project the impact that various levels of competitive supply will have on price to determine if returns will pay for any required capital costs over a specified time period.
- Understand that a specialty crop enterprise may not be limited to a single vegetable but may include a group of complimentary crops that fill a market niche. Several different crops may be required to gain a foothold in the market.

A successful specialty produce business requires knowledge and experience. It is advisable to start small and build the business gradually. Understanding marketing for specialty crops is the first step toward making profitable production decisions. Once a ‘niche’ crop becomes mainstream the profit margin may be too narrow which means it is time to move to different crops. Continually looking for new crops keeps ahead of the competition.

**The following sections describe production practices for specialty vegetables grouped by the general market outlets for the specific crops directing the producer’s attention to that critical part of the decision process.**

**Organic and Hydroponic Production:** Organic and hydroponic production which, in and of themselves, create niche-market specialty crops are not the focus of this guide, but most, if not all of the crops described, can be grown using ‘organic’ practices, *i.e.*, those approved under the USDA National Organic Program. Where appropriate, organic practices and pest control options are provided under each crop throughout this guide (see also section A 4. Organic Production). Using ‘hydroponic’ techniques to grow crops in a nutrient solution, usually within a controlled environment such as a greenhouse, is also suitable for many vegetable crops where there is sufficient market demand to justify the capital investment. Both production systems require selling to specific niche markets where demand provides the greatest return. ‘Hydroponic’ specialties should be marketed on their own unique qualities. If producing crops organically check with the certifying organization for requirements.

**World Crops:** Unfamiliar vegetables that are not conventional in North American markets but are culturally appropriate to many new immigrant populations that have settled throughout the Mid-Atlantic and Northeast provide opportunities for specialty produce farmers to cater to these ethnically diverse consumers. Major retailers are responding to these population shifts as well creating sales opportunities for both retail and wholesale growers.

## A. General Production Recommendations

It is critical to understand the ethnic community for which you will be growing to make the correct crop and variety selections, harvest at the correct stage, and to package in appropriate containers. The [worldcrops.org](http://worldcrops.org) website is designed to help growers exploring ethnic crop markets understand the nuances of marketing to such diverse groups (Table A-1). For example, peppers/chilis are popular Hispanic vegetables, but Hispanic consumers from different countries, even different regions of the same country, may prefer very different types of peppers.

Likewise, eggplant is very popular in many Asian and African cuisines, but in India, a small, egg size, pink 'brinjal' eggplant is preferred, while in China and Japan, consumers look for long, slender fruit. People in various African countries consume a white or pale green, medium-sized eggplant (a little smaller than the traditional Italian/Sicilian eggplants) that many call 'Bitter Ball' or 'Garden Egg'. A similar eggplant consumed in Brazil is called 'Gilo'. West Africans also use a pea-sized, red eggplant for medicinal purposes, known as the 'Ghanan pea' in most countries. That unusual eggplant is called 'Kiteley' in Liberia, while 'Kitley' is the term used for the 'Bitter Ball' eggplant in Ghana.

**Table A-1. World Vegetable Crops for Mid-Atlantic Growers**

(See <https://worldcrops.org/> for more information)

Vegetable Types	Community	Culturally Appropriate Crop Name
Solanaceous Eggplant	Brazil	Gilo
	West Africa	Bitter Ball, Kiteley, Ghanan Pea
	India	Brinjal
	France	Aubergine
Pepper/Chili	Mexico	Habanero
	Dominican Republic	Aji Dulce
Husk Tomato	Mexico	Tomatillo
<b>Cruciferous</b>	China, Southeast Asia	Napa/Chinese Cabbages, Pak Choys, Mustards, Flowering Broccoli
<b>Other Greens</b>	West Africa	Jute
	India	Fenugreek (Methi)
	Mexico	Purslane (Verdolaga)
	Universal	Amaranth, Roselle, Malabar Spinach

**“Designer Veggies:”** Coined to describe unusual produce used by creative chefs to decorate gourmet plates with more than a traditional garnish, “designer veggies” can be any crop grown for its size, shape, color, texture, or flavor. Types of “designer veggies” may include, but are not limited to, any/all the crops described in the following sections. They are usually ‘trendy’ crops that help celebrity chefs stand out from the crowd, so one year’s hot item may be a slow mover a year or two later, especially if several growers add more plantings. Radicchio can be considered one of the original “designer veggies”. When it appeared in produce aisles in the mid-1980s there was nothing like its bright red leaves with contrasting white veins and strong bitter flavor. Radicchio leaves are now common ingredients in many salad mixes, and studies have shown that it qualifies as a nutrient-dense ‘super food’. Coupling nutritional qualities with its ability to stand up to cooking in a variety of ways, along with increasing attention by food marketers, radicchio may once again be propelled into “designer veggie” status.

Success in the “designer veggie” business requires working closely with chefs and gourmand customers, paying close attention to food and trade publications and TV, attending produce and gourmet food shows, and being able to grow and deliver small quantities of labor-intensive produce.

## Baby, Miniature and Micro Vegetables

### Variety Selection

Demand continues for smaller vegetables among gourmet, restaurant, and specialty food outlets. **Micro-greens** are a trendy niche crop that are popular among small/beginner farmers. **Micro-greens** are seedling plants consumed at a stage (stem and cotyledon as 1 or 2 true leaves appear - smaller than transplant seedlings) between **sprouts** (roots and unopened cotyledons) and **baby** sizes (immature root vegetables or the first 2-4 true leaves of many greens). **Micro-greens** are cut above the soil line, so no roots or seed coats typically found in sprouted crops are included. Many types of vegetables can be harvested at these immature stages and sold as **baby** or **micro-vegetables** (Table A-2). There are other cultivars of vegetables which mature smaller than standard types of the same vegetable. These are referred to as **miniatures** and are grown to full maturity. Most seed companies offering specialty vegetables now also list specific varieties for **micro-greens** or **baby** production, in addition to miniature varieties.

**Table A-2. Baby and Miniature Vegetable Varieties and Harvest Stage**

Vegetable Type	Harvest Stage <sup>1</sup>	Baby Harvest Varieties	Miniature Varieties
<b>Beans</b>	IF	Filet bean types	Maxibel
<b>Beets</b>	IR, G	Any beet harvested at a small stage	Baby Beet
<b>Carrots</b>	IR	Round types, short types	Adelaide, Atlas, Aranka, Mokum
<b>Corn</b>	IF	Any sweet corn variety harvested within 3 days of silk emergence - supersweet varieties with tendencies to produce multiple ears/plant will increase yields	Baby Asian Corn
<b>Eggplant</b>	M	Fingerling eggplant	
<b>Greens</b>	G	Most greens, including mustards, cabbages (European and Oriental), chicories, amaranth, etc. can be harvested at the 4-6" stage. A mixture of baby greens and lettuces can be sold as "Mesclun" salad mix.	
<b>Lettuce</b>	G	Leaf types harvested before heading, Salanova varieties, one cut Types	Little Gem types, Baby bibb types, Mini romaine types
<b>Onions</b>	IR	High density plantings of white, yellow, and red onions for pearl onions	
<b>Peppers</b>	M		Miniature Baby Bell peppers, "snack" peppers, lunchbox peppers
<b>Radish</b>	IR, G	Early harvested radishes	
<b>Squash</b>	IF	Scallop or patty pan types, ball types, early harvested zucchini or yellow squash with blossom	
<b>Tomatoes</b>	M		Cherry, grape, and mini-pear tomatoes
<b>Turnips</b>	IR, G	White Asian types	

<sup>1</sup>IF=immature fruit, IR=immature roots (usually ½ to 1-inch diameter), G=greens (usually 4-6 inches or before head formation), M=miniature mature fruit.

### Culture

**Micro-greens** can be grown in protected culture for year-round, continuous harvesting. Typically, microgreens are planted in the containers in which they will be shipped and/or sold, leaving the 'harvesting' to the end user (chef or consumer). For wholesale customers, plastic transplant trays are lined with rock-wool/coir mats or a thin layer of soilless mix upon which the seeds are spread. Similarly, consumer packages such as clamshells or lidded trays can be used. Germination and 1 or 2 days of growth without light will cause the seedlings to stretch taller for easier harvest. This is followed by 2-3 days in full light which will allow the plant to produce chlorophyll and a dark green color. Un-cut micro-greens can then be delivered directly to chefs/consumers within a week to 10 days. Wheatgrass is grown in a similar manner using untreated wheat seed.

**Baby and miniature vegetables** are planted and grown much the same as standard varieties. Plant spacing is one major exception because miniatures are physically smaller and baby leaf and root crops are often harvested at the stage a standard variety would be thinned. Higher plant densities are desirable to maximize production. Baby leaf and some root crops can be grown in a solid bed by broadcast seeding since they will be harvested before crowding becomes a factor, or they may be drilled in rows 4-6 inches apart and as many across a bed as will fit. Spacing of miniature varieties will depend on the final size of the dwarfed plant. On the other hand, vegetables grown for their fruit (seeds or pods) such as beans, corn and squash should be grown at standard plant spacing to maximize output per plant. Crowding can affect the production of fruit, reducing yields, even if those fruit are to be harvested at an immature stage.

Field fertility may be modified depending on the crop and harvest stage. Immature, baby vegetables are harvested before they begin drawing significant amounts of nutrients from the soil. Most will perform well with little additional fertilizer beyond the reserves left from previous crops. Baby and miniature vegetables production can be scheduled to provide continual year-round harvests by using high tunnels or greenhouses.

Harvesting baby and miniature fruiting vegetables is time-consuming as many are hand harvested. Conversely, specialty equipment manufacturers, especially in Europe, have developed efficient mechanical harvesters for baby greens. These tools may need to be used in conjunction with matched bed shapers and other implements, so careful analysis of the market and size of production is required to justify the added expense. Smaller scale manual and semi-mechanical harvest tools have been developed for smaller operations.

## A. General Production Recommendations

### **Post-Harvest Handling**

Baby vegetables are immature crops at harvest-time and as such, both fruit and leafy crops tend to have higher respiration rates and are more tender than when they reach maturity. Proper post-harvest handling procedures are critical to maximize shelf-life. Gentle handling and special packaging from harvest on are required to reduce bruising and dehydration. Rapid post-harvest cooling removes field heat and extends shelf-life. This may be combined with triple washing to remove soil and field debris followed by spin-drying as a method of adding value for leafy greens. Follow food safety recommendations to reduce the chance for food borne pathogens (*e.g.*, *E. coli* and *Salmonella*).

Plastic-lined cardboard boxes, clear plastic food-service containers (clamshells) and inflated, resealable, plastic bags are common packaging options. The industry has settled on 3-pound plastic-lined, or wax treated, cardboard boxes for the wholesale trade. Larger bulk boxes/lugs may be suitable to send these products to fresh-cut processors who eventually repackage their finished products in consumer-oriented plastic bags or clamshell boxes. This packing system allows modified atmosphere treatment to reduce decay while providing support throughout the bulk package to reduce bruising/injury caused by the weight of the product itself. Micro-greens that are harvested at the farm are offered the most protection by use of clamshell boxes. Determine the appropriate package for the intended market.

### **Mesclun (French)/Misticanza (Italian)**

Mesclun usually refers to mixed young/baby salad greens and herbs. Ingredients in mesclun blends vary, consisting of many varieties of the crops listed in Table A-3. Seed companies sometimes sell pre-mixed selections for mesclun production, but since different species emerge and grow at different rates, it is recommended to grow each separately and mix after harvest. This allows the grower to create unique blends and time production to allow harvest of similar stages of growth of each species.

**Table A-3. Potherbs and Salad Greens**

Leafy greens can be described simply as any plant grown for consumption of its fleshy leaves, petioles, and/or stems, either raw (salad greens) or cooked (potherbs) (see also Greens section in Chapter F).

<b>Arugula</b>	Arugula, Astro, Darkita, Esmee
<b>Asian Mustards</b>	Mibuna, Misuna, Mizuna, Pak Choy; Flowering Broccoli
<b>Cabbages</b>	Red, Green and Savoy, Chinese Napa
<b>Edible Flowers</b>	Nasturtium, Viola, Violets, Pansy
<b>Herbs</b>	Parsley, Basils, Borage, Chervil, Chives, Fennel, Salad Burnet
<b>Lettuces</b>	Iceberg, Romaine, Crisphead/Batavia, Leaf, Bibb, Boston
<b>Other Composites</b>	Endive and Frisee, Escarole, Radicchio, Dandelion
<b>Miscellaneous</b>	Beet tops and Chard, Belgian Endive, Mache/Corn Salad, Orach, Claytonia/Miner's Lettuce, Sorrel, Purslane, Pea tips, Nasturtium leaves
<b>Mustards</b>	Cress, Mustard, Turnip tops, Watercress
<b>Other Oriental Greens</b>	Tricolor Amaranth, Shungiku Chrysanthemum
<b>Spinach</b>	Usually Flat leaf varieties

### **Pest Control**

**Under Protected Culture** Specialty vegetable production can be extended in the field using floating row covers or grown nearly year-round using high-tunnels in most of the Mid-Atlantic states. Pests likely to be encountered in high density plantings growing in high humidity are slugs, white flies, and botrytis. Slugs can be trapped and there are parasites for controlling white flies. Maintaining constant air circulation and adequate ventilation to reduce humidity within the plant canopy will reduce the incidence of botrytis. If making multiple harvests, carefully remove all dropped cut leaves as botrytis and bacterial soft rot get started on injured tissue.

### **Weed Control**

Weed control may be the most difficult aspect of baby leafy green and herb production. Selecting fields with low levels of weed seedbanks and free of perennial species is important. Preventing weeds from producing seeds will help with control in subsequent seasons. Herbicides must be labeled for the specific greens and herbs grown; consult the weed control sections in this publication for herbicide recommendations for specific crops. Consult the herbicide

label to determine if the time between herbicide use and harvest is equal to or exceeds the required pre-harvest interval (PHI).

Use cultural weed control methods such as stale seedbeds or plastic mulch when applicable. Mechanical weed control must be done in a planned, timely fashion. Most crops relying on mechanical weed control will require multiple cultivations, which will be more difficult in high density plantings. Resort to hoeing and hand weeding when necessary.

### **Insect Control**

Careful crop monitoring is required to produce insect-free greens. Timing production and using physical insect barriers such as floating row covers or insect netting can effectively control insects on many of the shortest season crops. Longer season crops usually require insecticides of some type to protect them from an array of root maggots, lepidopteran larvae, aphids, thrips, flea beetles, and more. Beneficial insects are available to control some of these pests. It is important to apply them when the pest is first observed. Additionally, crop rotation and prompt destruction of crop residue help prevent buildup of flea beetles and other localized insect pests. Effective IPM scouting can identify pest population changes and alert the grower when a pest control application may be required. Given the diversity of crops within this group, there may be unexpected pests occurring on small plots of crop plants, making control even more difficult. Read pesticide labels carefully to ensure that a product is registered for use on a specific specialty crop. Many specialty vegetables fall under Crop Grouping labels. Consult the crop specific guidelines in this book for pest control recommendations.

### **Disease Control**

Scout plantings on a regular basis and adopt IPM practices that will help produce a disease-free crop. Use genetic resistance to help limit potential losses due to disease. Many specialty vegetables fall under Crop Grouping labels, therefore consult the fungicide label and crop guidelines in this book for disease control recommendations.

Many specialty vegetables, especially heirloom/ethnic types of eggplants and tomatoes, are not resistant to common diseases that most modern hybrids have been bred to resist. Therefore, sanitation is critical to avoid building up populations of these pests. Solanaceous crops are especially susceptible to soil borne root diseases such as *Verticillium* and *Fusarium*, rarely seen in today's hybrid crops, but they are re-emerging as serious pests in heirloom/ethnic crops. In addition to sanitation, grafting a susceptible plant to a resistant rootstock may be an alternative that allows the desirable, susceptible plant to be grown in infested soils. More information on grafting can be found in "Grafting Vegetables" in section A 5. Transplant Production.

## **4. Organic Production**

Organic sales in the U.S. continue to rise, creating an opportunity for certified organic farmers. Upfront costs can be high, mainly due to certification costs and fees, but returns can be higher than for vegetables grown using synthetic fertilizers and pesticides. The United States Department of Agriculture (USDA) regulates the term "organic" to ensure uniform standards are used and protect the industry from dishonesty. To become certified organic, you must follow production and handling practices contained in the **National Organic Standards** (NOS; see <https://www.ams.usda.gov/rules-regulations/organic>) and be certified by a USDA-accredited agency such as the Northeast Organic Farming Association of New Jersey (NOFA-NJ; <https://nofanj.org>), Maryland Department of Agriculture ([https://mda.maryland.gov/foodfeedquality/pages/certified\\_md\\_organic\\_farms.aspx](https://mda.maryland.gov/foodfeedquality/pages/certified_md_organic_farms.aspx)), or Pennsylvania Certified Organic (PCO; <https://paorganic.org/>). The USDA maintains a list of accredited certifying agencies at <https://www.ams.usda.gov/resources/organic-certifying-agents>. If annual gross income from organic products is \$5,000 or less, a farm can be exempted from certification, but production and handling practices must follow the NOS and some restrictions regarding labeling and combination with other organic products apply. Certified organic production typically begins with a 3-year transition phase during which soil and farming practices must adhere to the NOS. After the transition phase, you can apply for organic certification. The transition phase may be waived or shortened if you can document that no prohibited substances have been used on the land for at least 3 years prior to applying for organic certification.

Successful organic production is a long-term proposition. It usually takes a couple of years and may take as many as four years for a site managed organically to reach its full potential for profitability. Organic production is knowledge- and management-intensive. Organic certification can increase market access. However, becoming a certified organic farmer may require learning new production methods. Additionally, documenting practices and

## A. General Production Recommendations

materials used through careful record keeping is mandatory. Forms for record keeping may be obtained from your certifying agency or online. Evaluate new products and methods on a small scale prior to large-scale adoption to test their effectiveness under the specific conditions of your farm. When implemented well, organic methods can improve soil quality and tilth through increased numbers of soil microorganisms and improved organic matter recycling.

Becoming familiar with the production and handling practices contained in the NOS is a good place to start if you are considering organic production. Keep in mind that you do not need to transition your entire farm at once. You can begin with just a part of your operation. Below are a few questions to consider if you are interested in organic production.

- Does a market for organic vegetables exist in your area?
- Are adequate resources available, including approved inputs, labor, and technical support?
- Would you be able to ride out possible reduced yields without premium prices during the 3-year transition phase?
- Are you willing to devote more time to monitoring pests, managing soil fertility, and record keeping?

If you answered “yes” to all the above questions, organic production may be for you. If you are beginning the transition phase from non-organic to organic production, consider a pre-transition phase if pest pressures are high in the planting area. A pre-transition phase is a hybrid between organic and non-organic production. During the pre-transition phase, synthetic pesticides are used along with organic-approved tactics to reduce pest pressures. Once pest pressures are reduced, organic pest management tactics are used exclusively.

The steps for becoming certified organic can be found in the publication “Organic Vegetable Production” at <https://extension.psu.edu/organic-vegetable-production>.

## 5. Transplant Production

**These recommendations apply only to plants grown under controlled conditions in greenhouses or hotbeds. Field-grown plants are covered under the specific crop in Chapter F.**

Producing quality transplants starts with disease free seed, a clean greenhouse, and clean planting trays. Many vegetable disease problems including Bacterial Spot, Bacterial Speck, Bacterial Canker, Gummy Stem Blight, Bacterial Fruit Blotch, Tomato Spotted Wilt Virus, Impatiens Necrotic Spot Virus, and Alternaria Blight can start in the greenhouse and be carried to the field. A number of viral diseases are transmitted by greenhouse insects.

Buy disease-indexed seeds if available. To reduce bacterial seed-borne diseases in some crops (*e.g.*, tomatoes, peppers, cabbages), seeds can be hot-water treated. For some crops, chlorine treatment can also be useful, but this will not kill pathogens inside the seed. For more detailed seed treatment recommendations, see section E 4.3. Disease Control in Seeds, Plant Growing Mix, and Plant Beds.

Prior to seeding in greenhouse areas, remove any weeds and dead plant materials, and clean floors and benches thoroughly of any organic residue. Shop vacuums are useful in removing debris from concrete and covered floors. Irrigation systems should also be cleaned to remove dirt and microorganism buildup (biofilms). Growing benches should be washed with a detergent to remove soil and residues. Growing areas should be sanitized with an antimicrobial product.

### **Sanitizing Greenhouse Surfaces and Treatment of Flats and Trays**

There are several different groups of sanitizers that are recommended for plant pathogen and algae control in transplant greenhouses. When possible/practical, rotation between these disinfectants is recommended. Make sure to read and follow the label since these products have different properties:

- **Alcohol (70% isopropyl)** is an effective disinfectant that kills microbes on contact. Since it is volatile, its effect is not long-lived. It is best suited for disinfecting propagation and grafting equipment, such as knives or shears by dipping or wiping.
- **Quaternary ammonium chloride salts** (Q-salts or “quats” such as Green-Shield®, Physan 20®, KleenGrow™) are labeled for control of fungal, bacterial, and viral plant pathogens, and algae. They can be applied to floors, walls, benches, tools, pots, and flats as sanitizers.
- **Hydrogen Dioxide, Hydrogen Peroxide, and Peroxyacetic Acid containing products** (*e.g.*, ZeroTol® 2.0, OxiDate® 2.0, SaniDate®12.0 (all 3 are OMRI)) kill bacteria, fungi, algae, and their spores on contact. They are labeled as disinfectants for use on greenhouse surfaces, equipment, benches, pots, trays, and tools.

## A. General Production Recommendations

- **Chlorine bleach (sodium hypochlorite)** may be used for pots or flats, but is not recommended for application to walls, benches, or flooring. When used properly, chlorine is an effective disinfectant. A solution of chlorine bleach and water is short-lived and the half-life (time required for a 50 percent reduction in strength) of a chlorine solution may be as little as a few hours. Bleach should be used in a well-ventilated area.

**New flats and plug trays are recommended for the production of transplants to avoid pathogens that cause damping-off and other diseases. If flats and trays are reused, they should be thoroughly cleaned and disinfected as described below. Permit flats to dry entirely prior to use. Styrofoam planting trays can become porous over time and should be discarded when they can no longer be effectively sanitized.**

**Sanitizing Trays with Chlorine:** First, wash out all excess growing medium since organic matter will deactivate the active chlorine ingredients quickly. Dip flats or trays several times in a labeled chlorine sanitizer at recommended rates (3.5 fl oz of a 5.25% sodium hypochlorite equivalent product per gal of water). Cover treated flats and trays with a tarp to keep them moist for a minimum of 20 minutes. Wash flats and trays with clean water or a Q-salts solution to eliminate the chlorine. It is important that the bleach solution remains in the pH 6.5-7.5 range and that a new solution is made up every 2 hours or whenever it becomes contaminated (the solution should be checked for free chlorine levels at least every hour using test strips).

### Transplant Production

Transplant growth is affected by many factors including temperature, growing media, fertilization, water, and spacing. A good transplant is grown under the best possible conditions. A poor transplant usually results in poor crop performance. Transplant production includes germination, growth, and hardening off phases. Table A-4 presents optimum and minimum temperatures for seed germination and plant growth, the time and spacing (area) required to produce a desirable transplant, and the number of plants per square foot. Seedless watermelon transplant production has specific requirements (see Watermelons in Chapter F). Details on sweet potato plant production can be found in Chapter F Sweet Potatoes.

### Making a Plant-Growing Mix

Pre-mixed growing media are available commercially (see below), but a good, lightweight, disease-free, plant-growing material can also be made from peat and vermiculite/perlite. The main challenge of making one's own mix is having a uniform and consistent composition, but it can also be less costly. Formulas for simple mixes can be found in Chapter G Resources (Tables G-4 and G-5).

**Table A-4. Temperature and Planting Recommendations for Transplant Production**

Crop	Optimum Day Temperature (F)	Minimum Night Temperature (F)	Weeks to Grow	Square Inch per Plant	Number of Plants per Square Foot	1020 tray size cells
Broccoli	65-70	60	4-7	2-3	48	72
Cabbage	65	60	6-7	2-3	48	72
Cauliflower	65-70	60	6-8	2-3	48	72
Celery	65-70	60	9-12	2-3	48	72
Cucumber <sup>1</sup>	70-75	65	2-4	4	36	50 or 72
Eggplant	70-85	65	6-9	4	36	50 or 72
Endive, Escarole	70-75	70	5-7	2	72	72 or 96
Lettuce	60-65	40	4-6	1	144	96 or 128
Melon <sup>1</sup>	70-75	65	2-4	4	36	50 or 72
Onion	65-70	60	9-12	0.5-0.65	220-288	288 or 312
Pepper	70-75	60	7-9	2-3	48	72
Summer squash <sup>1</sup>	70-75	65	2-4	4	36	50 or 72
Tomato	65-75	60	5-6	2-3	48	50 or 72
Watermelon (seeded) <sup>1</sup>	70-75	65	3-4	4	36	50 or 72

<sup>1</sup>Seed directly in container; do not transplant prior to setting in the field.

### Commercial Plant Growing Mixes

Commercial media are available for growing transplants and are generally recommended for growing vegetable transplants. Most of these mixes will produce high quality transplants when used with good management practices. However, these mixes can vary greatly in composition, particle size, pH, aeration, nutrient content, and water-

## A. General Production Recommendations

holding capacity. Commercial growing media will have added lime and may or may not have a starter nutrient charge (added fertilizer). Plants grown in mixes without fertilizer will require supplemental liquid feedings after seedling emergence. Plants grown in mixes with added fertilizers will require liquid feeding starting 3-4 weeks after emergence. If you experience problems with transplant performance, the growing medium should be sent to a testing laboratory. It is recommended to mix 3 to 4 bags of commercial product together before filling trays. Baled commercial mixes must be loosened before mixing. Before filling trays, the medium should be moistened so that it feels slightly damp but not wet. Medium should be used in the growing season in which it is purchased.

### **Transplant Trays and Containers**

Most transplants are grown in plastic trays with individual cells for each plant. Standard 10 x 20-inch plug trays (or more commonly 11 x 22-inch) can have 32 to over 500 cells. Larger cell sizes (32, 50, or 72) are best used for vine crops and for rooting strawberry tips. 72-cell and 128-cell trays are suitable for tomatoes, peppers, eggplant, and cole crops. Smaller cell sizes (128, 200, 288) may be appropriate for lettuce and onions. Larger Styrofoam and heavy plastic transplant trays are also available in similar cell sizes. Larger cells hold more growing mix/soil and result in better transplant survival in the field, but use more greenhouse space and it takes longer to produce the root ball. Individual plant-growing containers may also be used for vine crops and early market crops of tomatoes, peppers, and eggplant. Various types of fiber or plastic pots or cubes are available for this purpose. If plastic pots or trays are reused, disinfect as described previously.

### **Seed Planting and Germination**

Seeds that are over-sown in flats without cells to be “pricked out” (thinned or transferred bare rooted at a later date) should be germinated in 100% vermiculite (horticultural grade, coarse sand size) or a plant growing mix. It is recommended that no fertilizer be included in the mix or the vermiculite until the seed leaves (cotyledons) are fully expanded, and the true leaves are beginning to unfold. Fertilization should be in liquid form and at one-half the rate for any of the ratios listed in the “Liquid Feeding of Transplants” paragraph below. Seedlings can be held for 3 to 4 weeks if fertilization is withheld until 3 to 4 days before “pricking out.” These then can be transplanted into individual cell trays or grown on to use as bare-root transplants. This system can be used for tomatoes, peppers, eggplant, cole crops, and lettuce. Do not use this system for cucurbit crops.

More commonly, one seed is planted per cell directly in growing trays. Seeds that are sown in tray cells, pots or other containers can be germinated in a mix that contains fertilizer.

For earlier, more uniform emergence, germinate and grow seedlings on benches with bottom heat or in a floor-heated greenhouse. Minimum growing temperatures are listed in Table A-4. Germination rooms or chambers also ensure even germination, where higher temperatures can be maintained for the first 48 hours. Trays may be stacked in germination rooms during this period, but must be moved to the greenhouse prior to seedling emergence.

### **Plant Growing Facilities**

Good plant-growing facilities (greenhouses) provide maximum light to the seedling crop. The greenhouse cover material (glass, rigid polycarbonate, and polyethylene film) should be clean, clear, and in good repair. The ideal greenhouse will provide good heating and ventilation systems for effective environmental control. For hot air heating systems, place thermostats at plant level to maintain proper growing medium temperature. Combustion heating units located inside the greenhouse must be vented and have outside fresh-air intake and exhaust systems to provide air to and from the heater. Ventilation units must be adequate in size, providing 1.2 to 1.4 sq ft of opening for each 1,000 cubic feet per minute (cfm) fan capacity. Bottom-heating systems using circulating hot water, either on benches or on the floor, are better than hot air systems for germinating seeds and growing uniform transplants. This system heats the area around the plants, not the whole greenhouse, and can also result in reduced heating costs.

### **Liquid Feeding of Transplants**

In most instances, additional nutrients will be needed by growing transplants. Commercially available 100% water soluble greenhouse fertilizer formulations are recommended (see also Chapter C Irrigation Management, section C 3. Fertiligation). For most crops use a formulation with lower P than N and K levels (*e.g.*, 21-5-20, 13-2-13, 20-10-20, 17-5-17, 18-9-18). If you plan to fertilize with every watering, begin with N concentrations in the 30 to 50 ppm range and modify the concentration as needed. Use higher rates for tomato, pepper, and cole crops and lower rates for cucurbits (*e.g.*, watermelon, squash). Use higher rates when temperatures are high (late spring and summer) and lower rates when temperatures are cooler. Fertilizer requirements may vary substantially with crop and growing

## A. General Production Recommendations

conditions. For example, if fertigation is scheduled only once a week, N concentrations of 200 to 250 ppm may be required. Some growers may use a growing medium with no starter fertilizer. If that is the case, use 50 ppm N from emergence to first true leaf every 3 days, and 200 ppm N every other day from first true leaf to second true leaf.

For a less sophisticated way of applying nutrients, the following materials can be used for general use on transplants. Over an area of 20 sq ft, use 1 to 2 oz of 20-20-20 dissolved in 5 gal of water, or 2 oz of 20-10-15 dissolved in 5 gal of water. Rinse leaves after liquid feeding. Applications should be made weekly using these rates.

When using starter solutions for field transplanting, follow the manufacturer's recommendation. If concentrations are above recommended levels, they can cause excessive growth and reduce transplant quality. Highly concentrated nutrient solutions often can cause plant salt injury and leaf burning. Over-fertilized transplants will often "stretch" and have impaired field survival. **Caution: Fertilizers supply nutrients in the form of salts. High rates of starter solution can become concentrated and burn transplant roots when the soil becomes dry.**

### Watering

Keep the mix moist but not continually wet. Water less when the weather is cloudy. Watering in the morning allows plant surfaces to dry before nightfall and reduces the possibility of disease.

### Transplant Height Control

One of the most important considerations is managing "stretch" or height of transplants. The goal is to produce a transplant with a size that can be handled by mechanical transplanters or hand without damage, and that is tolerant to wind.

Most growth regulators that are used for bedding plants are not registered for vegetable transplants. One exception is Sumagic®, which is currently registered for use as a foliar spray on tomato, pepper, eggplant, ground cherry, pepino, and tomatillo transplants. The recommended label rate is 0.52 to 2.60 fl oz/gal (2 to 10 ppm), and 1 gal should be sprayed so it covers 200 sq ft of transplant trays (use 2 qt per 100 sq ft). The first application can be made when transplants have 2 to 4 true leaves. One additional application may be made at the low rate, 0.52 fl oz/gal (2 ppm), 7-14 days later, but do not exceed 2.60 fl oz per 100 sq ft for a season. Growers are advised to perform small-scale trials on a portion of their transplants under their growing conditions before large scale use.

For other crops, alternative methods for height control must be used, *e.g.*, the use of temperature differential or DIF, the difference between day and night temperatures in the greenhouse. In most heating programs, a greenhouse will be much warmer during the day than at night. The critical period during a day for height control is the first 2 to 3 hours after sunrise. By lowering the temperature during this 3 hour-period, plant height in many vegetables can be modulated. Drop the air temperature to 50-55°F for 2 to 3 hours starting just before dawn, and then return to 60-70°F. Crops vary in their response to DIF, *e.g.*, tomatoes are very responsive, while cucurbits are much less responsive.

Mechanical movement can also reduce transplant height. This may be accomplished by brushing over the tops of transplants twice daily with a tube, pipe or wand made of soft or smooth material. Brush plants when the foliage is dry and if plant damage is seen then reduce the number of times you are brushing the plants. Crops responding to mechanical height control include tomatoes, eggplant, and cucumbers. Tender plants such as peppers may be damaged by this method.

For some vegetables, managing water can be a tool for controlling stretch. After plants have reached sufficient size, expose them to stress cycles, allowing plants to approach the wilting point before watering again. Be careful not to stress plants so much that they are damaged.

Maximizing the amount of light plants receive helps reduce plant stretch. Giving plants adequate space and replacing plastic coverings as needed helps prevent reduction in light levels. Managing greenhouse fertilizer programs is another method for controlling transplant height. Most greenhouse growing media come with a starter nutrient charge, good for about 2 to 3 weeks after emergence. After that, apply fertilizers, usually with a liquid feed program. Fertilizers that are high in phosphorus will promote transplant stretch.

### Hardening

It is recommended that transplants be subjected to a period of "hardening" prior to transplanting to the production field. Reducing the amount of water, lowering temperatures, and limiting fertilizers causes a check in growth (hardening), which prepares plants for field settings. When hardening vine crops, tomatoes, peppers, or eggplants, do not lower the temperature more than 5°F (3°C) below the recommended minimum growing temperatures listed in Table A-4. Too low a temperature may injure plants and delay regrowth after transplanting. Exposing plants to

## A. General Production Recommendations

outside conditions is used for the hardening off process prior to transplanting. You can also use this for transplant height control during the production period. Roll-out benches or wagons that can be moved outside of the greenhouse for a portion of the day can be used for this purpose (see below).

Another tool is available for reducing transplant shock. The chemical 1-methylcyclopropene (1-MCP), which is marketed as the product LandSpring reduces ethylene production and stress on young plants. Ethylene is the plant hormone released when plants are injured or under stress, as is common during transplanting. Excess ethylene can cause leaf drop and wilting and can increase transplant losses. 1-MCP blocks ethylene from causing damage. LandSpring is labelled for broccoli, Brussels sprouts, cabbage, cantaloupe, cauliflower, cucumber, eggplant, muskmelon, bell pepper, non-bell pepper, summer squash, tomato, and watermelon. Apply to seedlings 1-5 days before transplanting,

### **Common Problems**

Poor growth and yellow or stunted plants may be attributable to the greenhouse growing medium. Greenhouse media manufacturers use good quality control measures, but things can go wrong, *e.g.*, through inadequate mixing, missing or the wrong proportions of critical components (wetting agents, fertilizers, lime), or defective components (poor quality). Media can also be affected by poor handling and storage, most commonly when media are stored outside or when bales or bags get wet, or if stored past the shelf life. Old media often are dried out and hard to rehydrate. If the medium is over a year old or possibly compromised, it should not be used (contact your supplier for inspections and tests on any suspect media). Avoid using overly dry or caked media, media that are difficult to loosen, media with a foul odor, waterlogged media or media that are resistant to wetting.

Most (but not all) media include a starter lime and fertilizer charge. The fertilizer is designed to provide 3-4 weeks of nutrients. If the fertilizer is missing, improperly mixed, or in the wrong proportions, seeds will germinate but seedlings will remain stunted. In this case, liquid fertilizer applications should start early.

Peat-based media are acidic in nature. Plants will perform well from pH 5.4 to 6.4. Lime is added to peat-based media and reacts over time with water to increase pH. Above pH 6.4, iron deficiencies in transplants are common. This also occurs if irrigation water is alkaline (has high carbonate concentrations).

In high pH situations (over 7.5), use an acidifying fertilizer (high ammonium content) for liquid feeds. Use of iron products such as chelated iron as a foliar application on transplants can accelerate plant recovery prior to the pH drop with the acid fertilizer. In cases with very high media pH, use of iron sulfate solutions may be needed to drop the pH more rapidly. Addition of dilute acid solutions to greenhouse irrigation water may also be considered in cases of excess alkalinity.

If lime is missing or inadequate from the growing medium, and pH is below 5.2, plants may exhibit magnesium deficiencies or iron or manganese toxicities. This also occurs in media that have been saturated for long periods of time. To correct this situation, apply a liquid lime solution to the medium and irrigate liberally.

Media that are difficult to hydrate may not have sufficient wetting agent or the wetting agent may have deteriorated; additional greenhouse grade wetting agent may be needed.

If the initial medium fertilizer charge is too high, or if excessive liquid or slow-release fertilizer feed is used, high salt concentrations can build up and stunt or damage plants (possible symptoms: leaf edge burn, "plant burn", plant desiccation). Test the media for electrical conductivity (EC) to see if salt levels are too high. The acceptable EC will depend on the type of test used (saturated paste, pour through, 1:1, 1:2) so the interpretation from the lab will be important. If salt concentrations are too high, leaching the growing media with water will be required.

Poor transplant growth or injury can also result from the following:

- Heater exhaust in the house caused by cracked heat exchanger, inadequate venting, use of non-vented heaters
- Phytotoxicity from applied pesticides or plant growth regulators
- Use of paints, solvents, wood treatments, or other volatiles inside the greenhouse
- Use of herbicides in the greenhouse or near greenhouse vents
- Low temperatures due to inadequate heater capacity or heater malfunction or excessively high temperatures due to inadequate exhaust fan capacity or fan malfunction

### **Grafting Vegetables**

Grafting is the act of joining two plants together. The upper part of the graft (the scion) becomes the top of the plant, and the lower portion (the rootstock) becomes the root system or part of the trunk. Grafting has resulted in increased yields, fruit quality, and tolerance to abiotic and biotic stresses. Research on annual vegetable crops was limited until the last decade when the grafting movement started in Asia and Europe. Grafting is used extensively in the

production of watermelon, cucumber, melon, tomato, and eggplant. Grafting can overcome tissue damage and/or plant mortality caused by the soil-borne diseases *Fusarium* and *Verticillium* Wilt, Bacterial Wilt, and nematodes. Grafting may reduce or eliminate the use of certain pesticides (especially soil fumigants) because the appropriate rootstocks will provide tolerance to many soil insects and disease pests. Grafting is also used to impart additional vigor to plants and to increase yields. Specific rootstocks have been developed for grafting the vegetables listed above. The selection of rootstocks will depend on the specific goals for grafting. There are often many rootstocks available. Consult your seed suppliers for more information.

Some commercial nurseries are starting to feature grafted transplants. As a rule, they are substantially more expensive than conventional transplants, so there should be reasonable assurance of the economic benefit. Any grower seeking to produce grafted plants should first consult technical resources, such as the websites in this section. Upgraded facilities and employee training will likely be necessary.

Two successful and easily performed grafts are the tube and cleft graft. The tube graft uses a 45° cut in the rootstock and the scion. The two pieces are subsequently joined together with the angles complementing each other and held together with a clip. The cleft graft utilizes a 90° cut in the rootstock perpendicular to the soil surface. The rootstock stem is then cut in half down the center; this cut should be around ½ inch depending on the size of the rootstock stem and scion. The base of the scion is then cut to form a “V” that will fit the notch that was cut into the rootstock. A grafting clip is secured around the graft junction. This type of graft often requires a larger grafting clip than the tube graft. It is important that both the scion and rootstock stem diameters are similar. Several trial seedlings should also be grown prior to any large grafting operation to ensure that the rootstock and scion seedlings grow at the same rate; if not, the stem diameters may not coincide, which can lead to a poor graft union.

Cucurbits such as watermelons, cucumbers, and muskmelons are often grafted using the one-cotyledon splice graft method. In this method, rootstock seedlings should have at least one true leaf and scion seedlings should have one or two true leaves. With a single angled cut, remove one cotyledon with the growing point attached. It is important to remove the growing point and the cotyledon together so that the rootstock seedling is not able to grow a new shoot of its own after being grafted. Cut the scion and match the rootstock and scion cut surfaces and hold in place with a grafting clip.

One of the most crucial aspects of producing grafted seedlings is healing the graft junctions. After the grafts are clipped back together, they need to be placed in a high humidity environment known as a healing chamber. A healing chamber can be constructed in various ways using wooden or metal frames and a plastic covering. The goal is to create a closed environment in which the humidity can be increased, and the temperature controlled. Open warm water pans or commercial humidifiers can be used to increase humidity. Propagation heat mats can be placed on the floor to control temperature. For the first several days in the healing chamber, light should be excluded as much as possible. The increase in humidity and decrease in light slow transpiration and keep scions from desiccating while vascular tissue reconnects the scion and rootstock. After 5 to 7 days in the healing chamber, seedlings can harden off in a greenhouse for several weeks before moving to the field. Grafting generally adds 2 weeks to seedling production. Grafting can be performed at various plant growth stages ranging from the 2 true leaf stage on.

## 6. Conservation Tillage Crop Production (No-Till, Strip-Till)

Conservation tillage crop production systems are beneficial for a variety of reasons, but they require different management than conventionally tilled soils. Some benefits from no tillage can be observed quickly, such as reduced soil erosion, conservation of soil moisture, and reduction in fuel and labor costs. Others benefits occur over time, such as reduction in soil compaction, improved soil structure, and increased soil organic matter. Eliminating tillage can also influence weed and disease severity and produce cleaner harvested products in vegetables growing on the ground.

Conservation tillage crop production systems can also pose several crop management challenges. Soil temperatures do not warm up as quickly in the spring and this can affect seed germination, nutrient cycling from crop residues, and transplant vigor. Type of crop residue, amount of residue, and desiccation timing all impact soil temperature and should be taken into consideration. Modifications to planters and heavier equipment may be needed to accommodate no-till production. Small-seeded crop species may be more difficult to plant in no-till systems.

Conservation tillage systems eliminate mechanical weed control. Since tillage used for seedbed preparation is eliminated, fields receive additional herbicide treatments to control emerged weeds and vegetation prior to or at planting. Thermal weed control (such as flammers) may be an option, but most other tactics (*e.g.*, mowing) are not

## A. General Production Recommendations

effective. Interrow cultivation with no-till cultivators has been used with some success in conservation tillage programs for weed management, but these implements are not readily available.

Nitrogen fertilizer must be managed properly when utilizing a conservation tillage production system. Crop residues typically contain an enzyme (urease) which can increase nitrogen volatilization from urea-containing fertilizer sources such as urea, liquid urea ammonium nitrate, or a variety of blends currently available. Management practices such as banding or incorporating nitrogen fertilizer with irrigation or rainfall should be considered to reduce urea-containing fertilizer contact with urease.

Nitrogen management in conservation tillage systems must account for microbial “tie-up”. High levels of crop residue, cover crops, or weed vegetation on the soil surface will result in microbes assimilating nitrogen and immobilizing it (*i.e.*, it is unavailable to the crop). Research has shown that 25% or more nitrogen fertilizer may be necessary in the initial conversion years from conventional to conservation tillage until the soil reaches an equilibrium. Previous crop residue (type and amount), current soil nitrogen concentrations, fertilizer sources, application timing, and application methods all need to be considered when making necessary nitrogen rate calculations.

Maintaining proper soil pH is one of the most important crop production considerations in conservation tillage and has a significant impact on nutrient availability and toxicity. Mixing lime into the soil with tillage is not an option with no-till systems, so consider adjusting pH to the optimal level prior to initiating a continuous conservation tillage system. Lime has relatively low water solubility and leaches slowly through the soil profile. Eventually, fertilizer, organic matter decomposition, and rain will lower soil surface pH, but changes to subsoil pH will take a longer time. Continued liming based on soil test recommendations will maintain the proper pH.

**Strip-till is a blend of tillage and no tillage within the same field.** A narrow strip of soil is mechanically tilled with specialized tools to incorporate fertilizers and plant residues, warm soils, and improve soil to seed contact. The area between the crop rows is managed as no-till.

No-till and strip-till production systems often use cover crops to provide a mulch that the vegetable crops are grown on. Rye, hairy vetch, crimson clover, and mixtures of these crops provide biomass that forms this mulch. Hairy vetch and crimson clover also provide nitrogen in the system. These mulches are often rolled with a roller-crimper prior to planting to provide the mulch base.

Under conventional tillage the plant residue is incorporated into the soil. However, in no-till systems, the cover crops may add additional plant residue that needs to be considered with management decisions such as preplant vegetation control, slower soil warming, plant residue management at planting, and fertility management. The amount of cover crop biomass (determined by when the cover crop is terminated) will dictate whether additional management is necessary.

## 7. Mulches and Row Covers

A favorable environment for plant root systems can be achieved by using plastic mulches and drip irrigation. Additional advantages of using row covers early in the season include increasing day time air temperatures and holding ground heat over night. This improvement in temperature can speed plant growth resulting in an earlier harvest. Mulches may discourage weeds and insect pests depending on the type.

### Plastic Mulches

Black and white-on-black polyethylene film (0.75-1.25 mil) are the most popular mulches. Other mulches include blue, red, green IRT and metalized. Black mulches are used to warm the soil and white-on-black mulches are used to cool the soil. Different mulch colors and compositions impart new functional properties to the mulch. Green ‘IRT’ types increase soil temperatures more than black plastic and suppresses most weeds including nutsedge. Results for other colored mulches such as red and blue have been inconsistent. Metalized or aluminized mulches repel certain insect pests (aphids, thrips, whiteflies) early in the crop growing cycle due to the reflectance of UV rays, but this benefit is lost once the crop canopy covers the mulch. This can be useful in cucurbit and tomato crops to delay the onset of certain virus diseases vectored by thrips, aphids, and whiteflies. Yellow mulches attract cucumber beetles and may also attract other insect pests. Note that planting date and environmental conditions influence crop responses to mulch color.

Soil fumigation may be used in conjunction with any type of plastic for weed, disease, and insect management, depending on the fumigant label. As the cost of soil fumigation increases, growers will likely need to reduce

application rates to maintain profitability. New mulches have been developed that have decreased permeability to fumigants. These “virtually impermeable film” (VIF) mulches keep the fumigant in the ground longer which allows for reduced application rates while maintaining efficacy. VIF mulches come in various colors for fall and spring plantings. Consult the fumigant label for the allowable reduction in use rate under VIF mulch and plant back restrictions. The cost of VIF mulches is higher than that of low-density mulches but this increase is usually offset by the savings gained from reduced fumigant rates. Another mulch type has been developed that is more retentive than VIF mulch, *i.e.*, “totally impermeable film” or TIF. Soil fumigant use rates may be further decreased if used in combination with TIF, consult the fumigant label (see also section E 1.5. Soil Fumigation).

### **Fertilization**

Measure soil pH before considering a fertilization program for mulched crops. If a liming material is needed to increase the soil pH, the material must be applied and incorporated into the soil as far ahead of mulching as practical. For most vegetables, the soil pH should be at or near 6.5. If the pH is below 5.5 or above 7.5 nutrients may be present in the soil, but not available to the plants.

Ideally a drip irrigation system is used with plastic mulch. When using plastic mulch **without** drip irrigation, all plant nutrients recommended for standard cultural practices should be incorporated in the top 5 to 6 inches of soil before laying the mulch. If equipment is available, apply all the fertilizer required to grow the crop to the soil area that will be covered with mulch. This is more efficient and effective than a broadcast application over the entire field. Non-localized nutrients may promote weed growth.

All essential plant nutrients, including major nutrients (N, P, K) as well as secondary and micronutrients, should be applied according to needs from soil test results and incorporated in the manner described above. Placing some of the required N under the mulch and then side dressing the remainder of the needed N along the edge of the mulch or in the row alleys after the crop becomes established has been found to be ineffective.

Applying some of the required N under the mulch and the remainder through the drip irrigation system is an effective way to fertilize. If using drip irrigation, see “Drip/Trickle Fertilization” in the crop sections in Chapter F (*i.e.*, eggplants, muskmelons, peppers, and tomatoes) for specific application rates.

### **Soil Conditions for Laying Mulch**

Soil texture should be similar, and plastic should be laid so that it is tight against the soil in a firm bed for effective heat transfer. Prepare the soil by incorporating crop residues, minimizing large soil clods, and removing rocks and other debris that could interfere with good contact between the soil and plastic. Plastic can be laid flat against the ground or on raised beds. Raised beds offer additional soil drainage and early warming. Use of a bed shaper prior to laying plastic allows for fertilizer and herbicide incorporation and can assist in forming a firm bed. Combination bedder-plastic layers are also widely used.

Before any mulch is applied, check the soil moisture level. Optimally the soil moisture level is at or near field capacity (field capacity is the amount of moisture left after a rain or irrigation event after surplus water has moved out of the root zone by gravity). Being at field capacity is extremely important when drip irrigation is not used because this moisture is critical for early growth of the crop plants as soil moisture cannot be effectively supplied by rain or overhead irrigation to small plants growing on plastic mulch.

### **Biodegradable Mulches**

Biodegradable plastic mulches have many of the same properties and provide comparable benefits as conventional plastic mulches. They are made from plant starches such as corn or wheat. These mulches are weakened by exposure to sunlight but are designed to degrade into carbon dioxide and water by soil microorganisms when soil moisture and temperatures are favorable for biological activity. Soil type, organic matter content, and weed pressure are other factors affecting breakdown. Unlike petroleum-based mulches, biodegradable mulches will usually be retained on the surface of the soil rather than be blown away from the application site. Most of the biodegradable mulch will eventually degrade or fragmentize, including the buried tucked edges. However, biodegradation is often unpredictable and incomplete. It is recommended that biodegradable mulch be incorporated into the soil at the end of the harvest or growing season. Cover crops can be planted the day after biodegradable mulch has been disked into the soil. In 2012, the National Organic Standards Board passed a motion allowing the use of ‘biodegradable bio-based mulch film’ provided that the mulch is ‘produced without organisms or feedstocks derived from excluded methods’ and meet certain degradation standards (at least 90% degraded in 2 years or less). However, currently only certain paper mulch products meet the organic requirements.

## A. General Production Recommendations

Field research has demonstrated that crop yields are comparable between biodegradable and non-degradable plastic mulches. Growers may be apprehensive about the cost of biodegradable mulch and the unpredictability of degradation rate. However, the initial cost is offset because disposal costs are eliminated. Below are some tips on using biodegradable mulch (excerpted from A. Rangarajan, Cornell University):

- **Storage:** Buy what you need each year. Product performance will be best with new products; more rapid degradation may be seen with older products. Store mulch rolls upright, on ends; pressure created from stacking may lead to mulch binding together or to degradation. Store mulch rolls in a cool, dark, and dry location; these products will start to degrade if stored warm, in sunlight and if rolls get wet.
- **Application:** Do not stretch biodegradable mulch as tightly over the bed as standard plastic mulches (contrary to recommendations for standard plastic that performs best when laid tightly over the bed). Stretching starts the breakdown of the biodegradable mulch, and will increase the rate of breakdown. The product will mold to the bed like commercial food wrap soon after application. Apply immediately prior to planting. If applied too far in advance of planting, the mulch may not last as long as needed. Sunlight and moisture will start the breakdown.
- **Incorporation into Soil:** Chisel or till the mulch into the soil as soon as possible after harvest to maximize breakdown; breakdown requires warm soil temperature and moisture. If mulch is incorporated after soil temperatures have dropped it may still be visible in the spring. However, as the soil warms, the product will further degrade and fragment. Rototilling will result in smaller mulch pieces that breakdown faster. Mulches will break down more quickly in soils with higher organic matter content.

### Floating Row Covers and Low Tunnels

These systems are being used for frost, hail, and wind protection, to hasten the maturity of the crop and to effectively exclude certain insect pests. Vented clear and translucent plastic covers are being used in low tunnels, supported by wire hoops placed at 3 to 6 feet intervals in the row. Porous floating row covers are made of lightweight spun fibers (polyester or polypropylene). They may be supported with wire hoops, PVC pipes or metal conduit hoops for plants that require a higher volume to grow, or they can be placed loosely over the plants without wire hoops for low growing plants such as vine crops and strawberries. Upright plants have been injured by abrasion when the floating row covers rub against the plants.

Clear plastic can increase air temperatures under the row cover on warm sunny days, resulting in a danger of heat injury to plants. Therefore, vented materials are recommended. Even with vents, clear plastic has produced heat injury, especially when plants have filled a substantial portion of the air space in the tunnel. Heat injury has not been observed with translucent materials.

Row covers are usually installed over plastic mulch using a combination of mechanical application and hand labor. Equipment that will cover the rows in one operation is available. However, farmer-made equipment in conjunction with hand labor is currently the most prevalent method used.

When considering mulches, drip irrigation, and/or row covers weigh the economics involved. Does the potential increase in return justify the additional costs? Are the odds of getting the most benefit in terms of earliness and yield from the mulch, drip irrigation, or row covers favorable? Does the market usually offer price incentives for the targeted earlier time window? Are you competing against produce from other regions? Determine the costs for your situation, calculate the potential return, and come to a decision as to whether these strategies are beneficial.

### Plastic Mulch Removal

Several methods of plastic mulch removal have been tried, but on small acreages it is removed by hand by running a coulter down the center of the row and picking it up from each side. Commercial tractor mounted mulch removal equipment is also available. High-quality, plastic mulch can be used for two successive crops during the same season when care is taken to avoid damage to the film. Thin wall (4 to 8 mil) drip irrigation tape cannot be removed and reused. However, high-quality, 16-mil drip tubing can be used for a second season if damage is minimal, and emitters do not become clogged. Crop foliage and weeds may hamper mulch removal. Prior to replanting or removing mulch, vegetation may be eliminated by using herbicides (see specific crop sections in chapter F), or delay removal until after frost.

### Plastic Mulch Disposal

Dispose of plastic in an environmentally responsible manner. Disposal regulations vary between states and municipalities. Contact your local solid waste authority for recommended methods of disposal in your area. Some states have developed recycling programs for agricultural plastics; consult state authorities.

## 8. Staking and Trellising

Many vegetable crops benefit from the addition of structural supports in the field. The benefits include: 1) better use of the available space and light; 2) improved air flow and more rapid drying of foliage; 3) reduction in certain disease pathogens; 4) protection against plant breakage; 5) protection of developing fruits and other plant parts against rain, dew, and sun; 6) ease of harvest, and 7) possible higher net yields. The disadvantages include the cost of materials, installation, removal, and disposal. Assess on a case-by-case basis if a structural support system is desirable.

Structural support systems have been used successfully for fresh market slicing, cluster, grape and cherry tomatoes, peppers, eggplants, cucumbers, climbing beans, and peas. The types of materials and how they are assembled differ for each crop. Specifics of the design and installation are included in Chapter F. If materials fail during the growing phase, the resulting damage can be catastrophic. Use high quality construction materials and adhere to minimum size and spacing recommendations. For wooden stakes, it is recommended that a clear hard wood source be used.

It is a common practice to re-use wooden stakes over many seasons. Because stakes are in contact with the environment and plant material, there is a significant probability that surfaces will become infested with pathogens, especially bacteria. If left untreated, infested stakes may re-introduce diseases into the field, although the extent of this problem has not been determined. It is recommended that stakes are thoroughly disinfested before re-use.

The preferred (and most expensive) method of stake disinfestation is a heat treatment. Pathogens are eliminated from wooden stakes with exposure to  $\geq 220^{\circ}\text{F}$  for  $\geq 15$  minutes. This can be accomplished in a large capacity autoclave, or seed dryer. It is unlikely that most growers will have access to such equipment. Alternatively, therefore, stakes may be exposed to disinfectants such as commercial chlorine solutions (sodium hypochlorite) or Oxidate® (hydrogen dioxide; see below). Research has shown that a 20-minute soak in a solution made of 5 to 20 parts by volume sodium hypochlorite (commercial bleach) to 80 to 95 parts by volume water is effective in eliminating pathogens **only from the surface** of wooden stakes. It is crucial to maintain the pH of the bleach solution within the 6.0 to 6.5 range, as effectiveness decreases at lower and higher pH levels.

Studies on stakes treated with bleach solutions show that pathogens may still be present beneath the surface at depths  $\geq 1/16^{\text{th}}$  inch. Pathogens embedded within the stake may be able to migrate back to the surface and re-infest plants, although this has not yet been demonstrated. To improve the effectiveness of procedures for removing microbial pathogens from stakes, consider the following: Add a non-ionic surfactant to the disinfecting solution; increase the soaking time to  $\geq 1$  h; apply a vacuum during the stake soak; use a higher concentration or more potent source of hypochlorite (such as “heavy duty” or swimming pool grade chlorine); or use stakes comprised of non-absorbent stake materials (such as plastic, fiberglass, or metal). Many growers have successfully used the commercial product Oxidate® or chlorine dioxide to disinfest stakes. Oxidate® is OMRI listed and has been demonstrated to be an effective control agent for several important plant pathogens. However, data on the efficacy of this treatment as compared to using heat or commercial chlorine solutions are not available.

## 9. High Tunnels

High tunnels are relatively low-cost, plastic-covered structures which extend the traditional growing season and protect the crop from stress. High tunnels are either freestanding or connected at the gutters to cover larger areas. Freestanding tunnels are generally between 14-36 ft wide and 120 to 200 ft long. High tunnels are typically tall enough for a person to stand straight up in at least part of the structure. While high tunnels are not greenhouses (generally no heat or automatic ventilation), the basis for their function and design are similar to those of a greenhouse. In the Mid-Atlantic region, year-round production of specialty crops is possible using freestanding high tunnels.

Taking the time to level the tunnel site prior to construction will make subsequent steps much easier. Spacing between high tunnels should be approximately 1.5 times the height of the nearest structure to facilitate snow removal, to provide for cross ventilation, and to reduce mutual shading. Runoff of rainwater from the structure should be channeled into storage tanks or through drainage pipes to avoid backwashing of water into the high tunnel. For freestanding high tunnels, metal bows approximately 1.75-2 inches in diameter are used as the support frame for a single or double layer of polyethylene covering (typically 6 mil greenhouse plastic that lasts 3-4 years; a double layer of plastic with an air gap between layers is especially recommended in colder areas and for growers wishing

## A. General Production Recommendations

to grow produce in winter). Polyethylene films characterized by different levels of light diffusivity and photo-selective properties are available on the market and can be used to create different microenvironmental conditions, with direct effects on crop performance, pathogens and insect incidence, and produce quality. The metal bows are spaced 4 feet apart and are connected to metal posts, which are driven at least 2 feet deep into the ground. End walls can have removable framing to allow the use of power tillage and bed maker/mulch layer equipment within the tunnel. High tunnels can also be designed as movable structures, which sit on wheels, rails, or skids, allowing growers to relocate the structure to different areas of their field, thus enabling better crop rotation and management of soilborne pests and pathogens, soil fertility, and overall soil health. Anchoring movable structures to the ground is critical to avoid losing or damaging the structure during windstorms.

A soil nutrient test should be conducted prior to constructing the high tunnel. Once the high tunnel is covered with plastic film, prepare the soil, apply, and incorporate lime and preplant fertilizer as recommended for the intended crop or crops (see Chapter F). Avoid using excessive levels of organic amendments at one time such as animal manure composts to avoid high concentrations of soluble salts and other nutrients such as phosphorus.

High tunnels can considerably increase yield potential, thereby increasing nutrient requirements. Plant tissue testing should be conducted at important growth stages during the season to ensure adequate fertility requirements are maintained. See Chapter B Soil and Nutrient Management for more details. Make beds, if needed, and install drip irrigation to supply moisture. Using a small bed maker/mulch layer, cover soil or beds with black or clear polyethylene to warm soil for spring crops. When transplanting crops into tunnels during July and August, use white or silver polyethylene mulch on the soil or beds rather than black polyethylene to reduce soil temperature and excessive heat buildup in tunnels. Shade fabric may be installed over the tunnels or white shade paint may be sprayed on the plastic cover in areas with high summer temperatures.

For freestanding high tunnels, snow removal from the top of the tunnels may be necessary after heavy snowfalls. Snow may need to be removed from the sides of the tunnels as well to reduce/eliminate outside water intrusion into tunnels and collapse of tunnel sidewalls. Gutter-connected or multi-bay high tunnels are constructed with much lighter posts and bows and cannot be used for crop production during the winter. During the winter season, the plastic on gutter-connected high tunnels must be bundled and moved to the gutters for storage. Low-cost caterpillar tunnels can be used for spring, summer and fall production. Hence, freestanding high tunnels allow for year-round production while gutter-connected and caterpillar tunnels do not.

The keys to successful production of vegetables and other horticultural crops in high tunnels are soil health management, crop scheduling, fertilization, ventilation, and irrigation. Table A-5 provides a relative planting and harvesting schedule for some vegetable crops produced using freestanding high tunnels in the Mid-Atlantic region. When planting high tunnel crops in the spring, it is generally recommended to transplant vegetable crops 2-4 weeks earlier than the earliest planting date in the field on bare ground. If unusually cold night temperatures are experienced several days to weeks after planting vegetable crops in high tunnels, floating row covers, low tunnels, thermal blankets and/or portable clean burning propane heaters (11,000 to 44,000 Btu per hour) can be placed in high tunnels until more seasonal temperatures return.

The most critical component of the system is ventilation. In freestanding high tunnels, ventilation is accomplished by rolling up the side walls to the batten boards, approximately 5-6 ft above the ground on each side of the tunnel. In gutter-connected high tunnels, ventilation is accomplished by sliding the plastic covering aside creating ventilation openings in the roof bows, as well as by opening the end walls. The use of a ridge vent may significantly reduce relative humidity and temperature fluctuations. Maintaining optimum growing conditions inside high tunnels without having extreme fluctuations in temperature and high humidity conditions can lead to early, high yielding and high quality crops. Checking and adjusting high tunnel internal temperature and humidity conditions several times a day will help ensure increased crop yields and profitability. Automation of the openings based on minimum and maximum temperature thresholds, wind and precipitations may facilitate the maintenance of optimal temperature and humidity conditions within the high tunnel.

Depending on the crop grown, there are several production systems that can be used in high tunnels. Conventional tillage and establishment of crops may be efficient for cool season crops that can be direct-seeded or transplanted such as, lettuce, onions, Swiss chard, spinach, collards, or kale. For warm season crops, especially cucurbits (cucumbers, squash, cantaloupe, and watermelon) and solanaceous crops, (potato, tomato, pepper, and eggplant) use of raised beds with plastic mulch and drip irrigation is required for optimum yield, maturity, and quality. Warm season vegetable crops dramatically benefit from higher soil temperatures in early spring in high tunnels. In addition, multiple cropping is possible from the initial raised bed/plastic mulch – drip irrigation system established in the spring. Permanent raised beds with a width of 24-48 inches may also be constructed in high

tunnels using wooden boards measuring 2 by 12 inches. Use of permanent raised beds may limit crops grown on them depending on the distance between raised beds (center-to-center) within the high tunnel. Some growers successfully use 30-36 quart potting soil bags that are drip irrigated to grow high tunnel crops. These bags are placed end-to-end in rows and on landscape fabric. Either one or two drip irrigation lines are inserted in planting holes in each bag. Additionally, small holes are cut on the bottom of the bags for drainage. Warm season vegetables can be grown using conservation or no till production practices within high tunnels. A winter cover crop is established the previous fall and terminated with silage tarps or mowing. The cash crop can be planted directly in the mulch or planted using woven ground cover as the mulch. High tunnel culture minimizes some diseases by reducing splash dispersal. In addition, appropriate adjustment of the plastic sides will also minimize leaf wetness duration.

Crops within high tunnels usually have more vegetative growth relative to open-field crops. Therefore, crops such as tomatoes and cucumbers should be pruned to manage vegetative growth and fruit load. Vining or indeterminate crops such as beans, tomatoes, and cucurbits should be trellised which increases marketable yield.

Preserving and improving soil health is critical for the long-term sustainability of high tunnel production systems. Implementing adequate crop rotations, the use of cover crops, and organic amendments can prevent the emergence of soilborne pest and pathogen issues. Monoculture and intensive cultivation may lead to a rapid decline of soil health and the emergence of several soilborne pests and pathogens including fungal pathogens (*i.e.*, Fusarium wilt, Fusarium crown and root rot, Timber rot, Verticillium wilt, Corky root rot, Southern blight), bacterial pathogens (*i.e.*, Bacterial wilt), and plant-parasitic nematodes (*i.e.*, root-knot nematodes, cyst nematodes, lesion nematodes, and dagger nematodes). The emergence of one or multiple soilborne pathogens may compromise and even prevent the successful cultivation of vegetables and other horticultural crops and requires the implementation of adequate soil disinfestation strategies. Historically, soilborne pests and pathogens have been managed using chemical fumigants or soil steaming; however, the application of these solutions in high tunnels poses several practical challenges, and is considered not sustainable, for their environmental impact and the negative effects on the overall soil health. Alternative soil disinfestation approaches applicable within the Mid-Atlantic region and considered more sustainable include biofumigation, biosolarization, and anaerobic soil disinfestation (ASD). Among those, ASD, also known as biological soil disinfestation, is emerging as a sustainable broad-spectrum pre-planting soil disinfestation strategy, capable of suppressing fungal and bacterial soilborne pathogens, plant-parasitic nematodes and weeds. As a microbial driven process, ASD requires minimum soil temperature of 60 F and is implemented amending the soil with readily decomposable organic amendments (wheat middlings, sugarcane molasses, crop residues), saturating the soil with water, and tarping the soil with an impermeable film to limit gas exchanges for a period of 3 to 4 weeks. The temporary development of anaerobic conditions, and the fermentation of the organic amendments, leading to the generation of organic acids and volatile organic compounds, allows the suppression of soilborne pests and pathogens, while preserving soil health, and upon re-aerating the soil, it is possible to establish a new crop. Vegetable grafting with the use of resistant or tolerant rootstocks is another sustainable approach proposed for addressing specific soilborne pest and pathogen issues affecting fruiting vegetables belonging to the Solanaceae (tomato, pepper, eggplant) and Cucurbitaceae (cucumber, watermelon, melon) family. Vegetable grafting may be also used to increase plant vigor and enhance tolerance to abiotic stress conditions such as sub-optimal temperatures, water deficit, salinity, and nutrient excess or shortage. Some airborne diseases and pest issues are prevalent in high tunnel environments. Leaf mold, gray mold, and powdery mildew, as well as thrips, whiteflies, aphids, and spider mites can become problematic in high tunnels. Synthetic fungicides and several fungicides approved for organic production are available for in-season management of fungal pathogens. When high tunnel sides are raised, fungicides and bactericides labeled for field use are allowed. When sides are lowered, fungicides and bactericides labeled for greenhouse use should be used (see Table E-14 “Selected Fungicides Labeled for Greenhouse Use” for specific disease and crop recommendations). See also Rutgers Cooperative Extension Fact Sheet No. 358 titled: “Important Diseases of Tomatoes Grown in High Tunnels and Greenhouses in New Jersey” (<https://njaes.rutgers.edu/fs358/>). This information is applicable to all states in the Mid-Atlantic U.S. region.

Integrated Pest Management for key pests affecting high tunnel crops is based on the principle of economic thresholds. Random plant inspection and use of sticky cards is encouraged in the semi-enclosed plant production spaces. The hot and dry environment can elevate certain pest problems as compared to the field situation. Many studies show that the use of beneficial insect or mite biocontrol agents is an effective pest management technique in high tunnels. It is important to know the right biocontrol agent, recommended release times and adequate temperatures and humidity at which they function best. Using habitat plants, such as alyssum and marigolds, and banker plants in a biocontrol program is beneficial and contributes to plant diversity in the high tunnels.

## A. General Production Recommendations

**Table A-5. Planting and Harvesting Schedule for Freestanding High Tunnel Vegetable Crop Production**

Crop	Method <sup>1</sup>	Average High Tunnel Planting Dates	Average High Tunnel Harvest Dates
Beet	TRP or DS	February-April; August-October 15	October-May
Bean (Snap)	TRP or DS	April-September 1	June-October
Bok Choi	TRP or DS	February-November	Year-round
Broccoli (crown)	TRP or DS	March-April; August	May-June; October- November
Broccoli (sprouting)	TRP or DS	March-April; August-October	May-June; October-April
Cabbage (Chinese)	TRP or DS	February 15-April 15; August 1-September 30	April-June; October-December 10
Cabbage (Green)	TRP or DS	March 15-May 15; August-September	May-December
Cantaloupe	TRP or DS	March 21-July	June-October
Carrot	DS	February 1-April 15; August-October	March-June; November-April
Cauliflower	TRP or DS	March 15-April 15; August	May-June; October-December 10
Chard	TRP or DS	Year-round	Year-round
Collards	TRP or DS	March-April; August-September	April-May; October-February
Cucumber	TRP or DS	April-September 1	May-October
Eggplant	TRP	April 15-August 15	July-October
Garlic	DS	October-November	June-August
Kale	TRP or DS	January-April 15; August-November 1	February-June; September-January
Kohlrabi	TRP or DS	March-April; August-September	May-June; October-December
Leek	TRP or DS	February 15-November 15	April-May; November-April
Lettuce	TRP or DS	Year-round	Year-round
Onion (Bulb)	TRP	February-March; October-November	April-July
Onion (Bunching Green)	TRP or DS	September-December; February-June	March-December
Pea	TRP or DS	February-April	May-June
Pepper (Bell)	TRP	April-July 20	June-November
Potato (Irish)	DS	February 14-March 15; August	May-June; October-December
Radish	DS	February-April; September-December	February-May; November-January
Spinach	DS	January 1-May 1; August-December	January-May; October-December
Summer Squash	TRP or DS	April-May	May-June
Tomato	TRP	March 15-July 15	June 1-December 5
Turnip	DS	February-April; September-December	February-May; November-January

<sup>1</sup>TRP=Transplanting, DS=Direct Seeding.

## 10. Greenhouse Production

Many growers have an interest in increasing productivity and quality as well as having seasonal products such as tomato, sweet pepper, cucumber, lettuce, arugula, and herbs in the off seasons or year-round. To do this in the Mid-Atlantic U.S., a temperature-controlled greenhouse structure might be needed. Greenhouse production requires a much greater level and often entirely different management strategies than field production. There are multiple greenhouse designs in the market, each with its own advantages and disadvantages, depending on the farmer's needs, environmental conditions, and level of economic investment. Greenhouse production generally requires different varieties (often of indeterminate growth habit), nutrient sources and regimens, plant training, and pest management than field production. Hydroponic systems are commonly used, although production in soil is also possible. Other fully lighted “warehouse” or vertical production systems under artificial light have been developed.

The extensive differences between greenhouse and field production preclude the inclusion of these techniques in this guide. There are many complete guides for the production of vegetables in greenhouses that have been developed and distributed through the cooperative extension service in various states. Links to some guides are provided below. This list is not all-inclusive and does not endorse these guides exclusively.

[https://edis.ifas.ufl.edu/collections/book\\_florida\\_greenhouse\\_vegetable\\_production\\_handbook](https://edis.ifas.ufl.edu/collections/book_florida_greenhouse_vegetable_production_handbook)

<https://ceac.arizona.edu/resources/intro-hydroponics-cea>

## 11. Wildlife Damage Prevention

### Philosophy of Damage Management

Because agricultural lands offer abundant opportunities to successfully find food and shelter, many species of wildlife view farms as viable and attractive habitats to visit or occupy. Not all wildlife species damage agricultural crops, yet some can inflict serious economic losses on growers. Additionally, resolution of wildlife conflicts sometimes can be difficult for growers because they often have little direct control over what happens on the lands surrounding their farms, areas that potentially can provide refuge for the wildlife causing damage on their farms.

One of the realities of running a farm is that interactions with wildlife on agricultural lands are inevitable. However, growers must recognize that trying to eliminate all wildlife damage is not realistic and that some level of damage attributed to wildlife represents another of the costs of doing business, much like the effects of weather or disease outbreaks. The priority should be a focus on reducing damaging or repetitive episodes of wildlife inflicted loss to a tolerable level. To achieve this goal, developing and implementing a well-designed **wildlife damage management plan** is strongly recommended, one that proactively (not reactively) will help prevent or reduce damage. An effective management plan will identify the full suite of legal damage management techniques and strategies applicable to the different types of conflicts being experienced. It should clearly establish when and under what conditions those options could be employed (*e.g.*, are special permits or authorizations necessary and, if so, from what entity?). Such a plan should also delineate areas of the property where the need for damage prevention is greatest, as well as areas where some level of damage can be tolerated; this allows one to prioritize allocation of resources to areas of highest value or concern.

### Considerations in Developing Damage Response

Wildlife damage management practices fall into 3 categories: husbandry methods, non-lethal techniques, and lethal options. This categorization also represents the hierarchical order in which these methods should be implemented: start with husbandry methods that typically are simple, less costly, and easy-to-implement behavioral or operational modification approaches; if those don't provide the desired level of relief, consider non-lethal techniques, which inherently are more costly, time-consuming to install, and likely require a greater level of maintenance to remain functional. Lethal options must be viewed as methods of last resort, not options of first choice.

Different damage abatement strategies obviously will have varying levels of effectiveness and inherent risks. Generally, a damage management plan that proactively integrates several different damage abatement techniques, whether used concurrently or in rotation, will be more effective than a reactive strategy that repeatedly relies on only a single technique. Although a variety of damage management options exists, not all are suitable or legal for use in all conflict situations. Many options legally are restricted from use depending upon the species of wildlife involved and the site or location where the conflict is occurring. By design, some damage abatement options are most effective for temporary, short-term, or localized use, whereas others are better suited to permanent, long-term needs. Each situation of human-wildlife conflict can be unique, so management options usually need to be tailored to the needs of a specific site.

Acquisition and implementation costs associated with each management option can vary substantially. Before deciding on a particular management technique, you should first estimate both the direct and indirect costs you actually are experiencing from wildlife damage. A direct cost would be the estimated annual loss of value from reduced yield or production through consumption or ruin of the crop by wildlife. An indirect cost would be the annual monetary investment (in time and/or labor) expended in monitoring for or detecting new damage. Calculating this estimated total annual cost, in terms of the actual economic loss *due only to wildlife*, will help you decide which, if any, strategies are cost-effective. Remember, if you are spending more in your attempts to acquire and implement preventive techniques than you are experiencing in actual annual damage, you are likely to lose money; but don't forget that some acquisition/operational costs can be pro-rated over the anticipated life expectancy of certain physical techniques, like fencing, thereby reducing the actual annual estimate of incurred acquisition cost. Though hard to swallow, the reality is that, for some producers, it may be more practical and economically feasible to simply tolerate some level of damage than to attempt to eliminate it. To determine if control is feasible and to inform your decision-making on selecting appropriate control techniques, you must have good estimates of the level of economic damage you are incurring and reliable estimates of what each abatement technique costs to implement. More importantly, to assess whether the techniques you chose to utilize actually provided cost-effective management of your damage, you must have good pre- and post-treatment economic estimates of the amount of

## A. General Production Recommendations

damage for comparison. If you find that you are spending more on your effort to manage damage than you are realizing in savings, you're obviously losing money. In such cases, you should reexamine the suite of available methods to find a more cost-effective approach or accept that trying to reduce this damage simply is not economically feasible.

Prior to employing any damage abatement practice, you must be sure that you have correctly identified the species causing the damage. Do not assume that any animal you see in your fields is the culprit responsible for any damage you detect - presence alone does not equate to guilt. Other forms of evidence must be present to confirm culpability (such as direct observation of foraging, signs of unique feeding behavior, etc.). Wildlife populations are regarded as a public resource and, as such, the animals visiting and perhaps causing damage on your farm are managed and protected by state and federal laws, regulations, and elements of Code. Most of the restrictions imposed on the use of damage management practices (e.g., trapping, shooting, pesticide applications) are specific to individual species and can become quite technical. If you mistakenly assign blame for damage to the wrong wildlife species and employ a technique that is not authorized for use on that species, you can be cited for violation. Therefore, before employing any management practices, check with your county extension agent, local conservation police officer/game warden, or your district wildlife biologist to review depredation permit requirements and/or legal issues related to "take" or use of wildlife.

## Damage Management Strategies for Common Wildlife Conflict Situations

Below you will find information on managing conflict situations involving some of the commonly recognized wildlife species known to be responsible for agricultural damage. This presentation is not a comprehensive summary of all possible options - the objective is to provide sufficient understanding that can assist decision-making of affected growers. This presentation purposefully takes a generic overview approach, given the breadth of the geographic region covered and the diversity of state-specific legal requirements and statutes. Producers are encouraged to work closely with the authorities in your state regarding the best and most legally accurate advice.

### Bears

Damage caused by black bears (*Ursus americanus*) to field crops often is characterized by trampled, pulled down, or broken plants. In corn fields (especially sweet corn), bears create circular patches within the field where the stalks are pulled inward toward the center of the circle; this behavior typically occurs just prior to the corn's peak maturity or what some call the "milk stage." Within these circles, a bear will remove an ear of corn from the stalk and consume the majority of kernels on the ear before moving on to the next ear. It can be difficult to detect bear activity from the ground at the edge of a corn field because bears often wade deep into the field before beginning to start this feeding behavior - the presence of damage and its extent often is first detected once the harvest is underway. Bears also will climb fruit trees to access the maturing fruits and, in the process, may cause substantial damage by breaking or cracking the branches of affected trees. Small fruit growers (e.g., blueberries, brambles) similarly can witness bear-caused damage in the form of broken branches or splintered bushes from the bear pulling down branches to gain better access to the berries.

In livestock depredation settings, the pattern of the kill often gives clues to the culprit. Bears typically attack from the rear and try to bring down the targeted animal forcibly using its weight and strength and via bites to the top of the neck; strafing claw marks often are present on the carcass. Scat (feces) and footprints are usually present in the area of feeding activity.

There are no guaranteed bear management strategies that offer complete protection against all types of damage, but several strategies used in combination may offer some relief:

- **Cultural practices and habitat modification** can help to deter bears. First and foremost: Do Not Feed Bears! Restricting access to potential food resources, such as storing feed in bear-resistant containers, timely collection and disposal of animal carcasses (via deep burial), and removing fruit drops and other organic wastes often can reduce the attractiveness of the site to bears. Containing livestock in pens away from wooded areas may reduce negative interactions, particularly during calving/lambing season. Because bears generally avoid open areas away from protective cover, maintaining a mowed buffer approximately 50 yards wide around crop fields, particularly where fields are adjacent to the woods, may reduce bear activity. Alternating or strip planting row crops, especially those deemed more attractive to bears, may help hide vulnerable crops from the bear's direct line of sight.
- **Fencing** is effective in reducing bear damage; however, fencing can be expensive and may not be cost-effective for all farmers. Electric fencing is the most effective design to deter bears and would be recommended in most

instances, where allowed by law. Electric fences should utilize high voltage (~6,000 volts), low-impedance (short pulsed) charger systems. When first installed, all electric fences should be baited with an attractant (e.g., peanut butter, sardines) to lure bears close so they learn to associate the fence with a negative consequence. These attractants should be placed at the bear's head height (approximately 3 ft) along the entire perimeter to better deliver the shock to the bear's muzzle.

- **Visual/auditory deterrents or harassment techniques** have been used to deter black bears from crop fields. Pyrotechnics, horns, bright lights, propane cannons, and other devices provide both visual and auditory stimulation. When damage coincides with the "bear chase season," allowing a local bear hunt club to run dogs through a field often provides temporary relief against bears that are using those fields. The success of these techniques is highly variable. Bears can become habituated to constant or repetitive disturbance, so a rotation of several types of sensory deterrents should be used and their position in a field should be relocated frequently. Where bears have become tolerant of human activity, sensory deterrents often will not be effective. Human-conditioned bears can be dangerous, and caution is advised.
- **Aversive conditioning techniques** are intended to more forcibly change or eliminate undesirable bear behavior. Options that potentially may be available to harass a troublesome bear include using paint balls, projected bean bags, or rubber bullets, all of which are intended to send a strong message without injuring or maiming the animal. These approaches may not be allowed in all states, so check with your state wildlife department before use.
- **Shooting** problematic black bears is viewed as a last resort management practice, but one that may be necessary to reduce persistent crop damage caused by a single returning individual or family group. Special "kill permits" are required to shoot a bear outside the regulated open season, so farmers need to work closely with their state wildlife agency. Farmers who have persistent damage should develop a relationship with local bear hunters or chase clubs to increase the level of pursuit activities, when legally allowed, on or adjacent to the farm as a means of reducing future losses. **This practice is not permitted in some jurisdictions. Consult your local Wildlife Management Authority.**

## Birds

Bird damage often takes the form of the total consumption of grains or small fruits or the creation of holes or other surface blemishes inflicted from the birds' pecking on ripening fruits, vegetables, or woody stems. Proper identification of the offending bird species is necessary before implementing any management strategy because most bird species are protected under federal statutes (e.g., the Migratory Bird Treaty Act). That said, growers who face serious damage events may be allowed certain latitude under special permits to use techniques that might otherwise be prohibited. For example, producers may be allowed to "take" flocking blackbirds causing damage under provisions of the Migratory Bird Depredation Order (50 Code of Federal Register 21.100). "Blackbirds" generically refers to a group of about 10 species, often including Common Grackles (*Quiscalus quiscula*), Brown-headed Cowbirds (*Molothrus ater*), and European Starlings (*Sturnus vulgaris*). However, growers still need to apply for and receive authorization from the U.S. Fish and Wildlife Service before any lethal action is taken. Starlings, House Sparrows (*Passer domesticus*), and Common Pigeons (*Columba livia*) also are common to farms, where they inhabit the rafters of barns, warehouses, and other structures or contaminate feed bunks with their droppings while foraging on livestock feed; these species are considered exotic or introduced species and thus have no legal protection under the Migratory Bird Treaty Act - lethal "take" can occur at any time, so long as the state's lawful methods of take are followed.

Birds inside packing houses represent a serious source of fecal contamination, which may violate USDA food standard guidelines. Fecal contamination of fruits and vegetables in the field also can occur if fields are located near a bird roost where large numbers of birds congregate. Such situations can complicate a producer's ability to fulfill Good Agricultural Practices (GAP) standards and typically create health hazard conditions for which relief also may be allowed under the 50 CFR 21.100 statutes. It is imperative to check with the state Fish and Wildlife authority before implementing any management to ensure compliance with state and federal wildlife laws.

As was true with bears, there are no guaranteed management strategies that offer complete protection against all types of bird-inflicted damage; strategies that may offer some relief include the following:

- **Cultural practices and habitat modification** may reduce some forms of crop damage. Because the most severe instances of blackbird damage commonly occur within 5 miles of a roost, locating plantings of highly attractive crops outside this radius is recommended wherever established roosts are known to exist. Blackbirds generally do not prefer soybeans, hay, wheat, or potatoes. Obviously, large fields of a single crop (e.g., sunflowers)

## A. General Production Recommendations

are highly attractive to large flocks of birds; damage in such crops is likely to be spread sporadically throughout the entire field, but the heaviest impact typically is concentrated toward field edges closest to cover or the direction from which birds have come from the roost site. Modifying the physical structure of roosts or encouraging birds to abandon existing roosts may reduce the number of birds in the area. For example, eliminating stands of introduced bamboo or thinning dense stands of conifer trees have been shown to reduce crop damage by removing favored roosting spots. Removal of about 1/3 of a tree's crown or 1/3 of a stand of trees has been successful in reducing or dispersing birds from a roost. Keep in mind, though, that you also will modify habitat used by other non-destructive and potentially desirable bird species. Installing artificial hunting perches in or adjacent to fields to enhance the success of raptors that feed on other birds may reduce blackbird numbers as a result of the threat of predation.

- **Exclusion** (*e.g.*, netting) typically is practical only on small acreages or for high-value crops. Lightweight netting has been used successfully to prevent bird damage either by draping it over individual plants or constructing a frame and stretching netting over an entire block of plants. To prevent birds from entering packing houses, netting or some other type of barrier should be placed over openings larger than 1/2 inch. In doorways where frequent pedestrian, vehicle, or machinery traffic occurs, hang heavy translucent plastic or rubber strips, or install self-closing doors to prevent birds from accessing the building.

- **Repellents** can be used to mitigate bird damage. Methyl anthranilate, the primary ingredient of artificial grape flavoring, is registered by EPA for use as a bird repellent for certain species. However, methyl anthranilate remains viable for only short periods of time as it loses efficacy when exposed to UV radiation and weathering. Sucrose solutions may be applied to fruits to deter birds, but the efficacy of this method is not well documented and actually may attract other pests, such as Japanese beetles.

- **Audio/visual scare tactics** usually are effective only when used to provide short-term protection for crops, such as the period immediately before peak condition is attained and just before harvest normally would begin. Blackbirds are intelligent animals and demonstrate rapid habituation to repetitive or predictable patterns and disturbances. Frightening methods must be changed and/or relocated often to maintain the desired effect. Frightening devices include both visual and auditory deterrents. Pyrotechnics (*e.g.*, propane cannons and shotguns), Mylar balloons and reflective metallic tape, raptor-shaped kites, scarecrows, flashing lights, water sprayers, and tape-recorded bird-distress calls or predator attack calls all represent examples of harassing techniques; success among these devices varies substantially. Based on reported success in repelling fish-eating birds in aquaculture facilities, use of lasers to reduce bird damage has garnered attention of producers and may offer a viable option. In general, scare tactics should be activated early to mid-morning and mid- to late afternoon, when birds are most active. For maximum effectiveness, it is best to use two or more devices in combination, varying the times and places these devices are employed, and be persistent.

- **Lethal Methods**, other than those allowed under provisions of the Migratory Bird Treaty Act noted earlier in this section, typically are prohibited or tightly controlled by regulation or statute, so caution is advised here. Flocking birds that ingest registered treated bait material will fly in an erratic fashion, produce distress calls, and ultimately die, but this unusual behavior typically triggers an alarm response among the remaining birds in the flock, causing them to vacate the area. NOTE: use of such chemicals typically is restricted only to certified applicators (usually representatives of USDA APHIS-WS), so check with state and/or federal authorities about the legality and possibility of employing chemical frightening agents on your farm. Where shooting via permit is allowed, it helps to maintain good working relations with local authorities; farmers should alert local police and/or abutting neighbors about the intention to conduct a shooting program. This helps the police avoid responding to calls received at the local "911" Emergency Dispatch system when the shooting starts.

## Deer

Deer damage may take the form of direct foraging, antler rubbing, and/or trampling of crops. Foraging damage from deer is characterized by a torn, jagged appearance on vegetation or a ragged break on woody stems. Most browsing damage occurs from ground level and up to a height of about 6 ft. Residual damage may occur from the trampling or matting down of vegetation as deer travel through crop fields or bed down to rest. Antler rubbing damage occurs as males try to shed the velvet coating from their antlers each autumn and typically leaves behind scarred saplings, broken limbs, bruised bark, and/or exposed wood. Rubs usually are located about 2-3 ft. above the ground on the trunks of small trees or widely branched shrubs.

Effective management of deer damage often requires using several deterrent strategies and should embrace options from within the full suite of available husbandry, non-lethal, and, where warranted, lethal options. Each

method carries certain benefits and inherent drawbacks; therefore, an accurate assessment of management needs and desired outcomes is critical to selecting the most appropriate techniques.

- **Habitat modification** includes efforts that seek to change the landscape and make it less attractive to deer. White-tailed deer (*Odocoileus virginianus*) are creatures of edge habitats; they prefer locations where two or more vegetation types or age classes meet. Habitat modification usually involves eliminating preferred types of vegetation, planting non-palatable (“deer-resistant”) species, or creating alternative foraging areas to attract deer away from crops. Although these strategies have been used to reduce incidences of deer-vehicle collisions and browsing on residential vegetation and commercial landscaping, habitat manipulation alone is unlikely to significantly reduce damage in agricultural crop fields or nurseries.

- **Harassment and audio/visual scare tactics** are intended to persuade deer to leave an area where they are not desired. Examples of hazing and scare techniques include dogs, auditory deterrents (such as propane cannons, pyrotechnics, playback of recorded distress calls), and visual deterrents (such as bright flashing lights, lasers). An increase in the use of free-roaming dogs contained within invisible fencing has brought some success to farms but is dependent on the number and aggressiveness of dogs used relative to the size of the area needing protection. Dogs tethered by chains or ropes are not effective because deer quickly detect that the dog’s movement is restricted to a certain range, beyond which deer may move about freely. If implemented, hazing campaigns need to be relentless and persistent; they generally are better suited to small areas where damage from deer is minor or where other strategies (*e.g.*, hunting) may be prohibited.

- **Fencing** can be an effective management tool for eliminating or reducing deer damage and, in many cases, may be the most effective damage abatement option. When attempting to protect large areas, permanent high-tensile wire (HTW) electrified fences are recommended. These fences consist of a series of electrified smooth wires spaced about 8 inches apart that extend to an overall height of 10 feet. HTW fences are durable and long-lived but do require regular monitoring (to check that sufficient voltage is flowing) and periodic maintenance (*e.g.*, removal branches or other material that may short-out the flow of current). Temporary HTW electric fencing or fences that use polytape strands are available; these usually are better suited for use on smaller acreages or where protection is needed for only a short period of time. When using any form of electrified fencing, it should be charged at all times to prevent deer from becoming emboldened or gaining confidence from the lack of deterrence during a shutdown. Electric fences that have been baited with an attractant (*e.g.*, peanut butter) demonstrate noticeable enhanced success over non-baited fences, as deer are more likely to develop an immediate association between the fence and its negative consequence when drawn in by baiting. The addition of cloth strips, flagging, and reflectors certainly increase the visibility of a fence, but these additions display only marginal improvement in efficacy over fences lacking such visual cues. Although other fencing alternatives exist, such as slant fencing, double-barrier fencing (2 parallel rows of fence placed approximately 3-4 feet apart), or heavy-duty extruded plastic mesh fencing, these may not provide the level of protection or cost-effectiveness of a well-designed and properly installed and maintained electric HTW fence. It is important to note that no type of HTW fence will eliminate all penetration by deer. If absolute protection from deer is desired, the only fence design that can guarantee that outcome is a 10-ft tall woven wire fence. However, in most situations, producers typically cannot justify the costs of procurement and installation of such a fencing system.

- **Repellents** are products that have strong or repugnant tastes, odors, or a combination of both taste and odor. Those that animals find most offensive will encourage them to avoid the area being protected. There are 2 types of repellents: contact repellents and area repellents. Contact repellents are applied directly to vegetation or objects via spray, shakable powder, or brush. Area repellents are applied in the general vicinity of the object where protection is desired (often on the ground surface) and repellency is derived primarily by odor. Repellents can be expensive, based on the initial cost of materials, but more so by the need for frequent reapplication. Rain and weathering eventually will wash a repellent off vegetation, even if a “sticker” is used. Based on multiple research studies, products derived from or containing a high concentration of putrescent egg solids and/or animal fat-based products provide greater repellency than those without these active ingredients. The level of attractiveness (*i.e.*, palatability) of a food resource to deer, the density of deer in the area, and the availability (or lack thereof) of other natural foods in the area all will influence the effectiveness of repellents. Many deer repellents are labeled for use only during the dormant season or on non-consumable vegetation, so growers must be sure to follow the manufacturer’s label instructions. Repellents used during the growing season must be applied frequently to keep pace with the emergence of new, but as yet unprotected, growth. Regardless of the type of repellent used, all repellents are intended to reduce,

## A. General Production Recommendations

rather than eliminate, deer damage; repellents should be used in conjunction with other damage abatement techniques to maximize overall success.

Results from recent research on the use of an auditory repellent offer some encouragement for those seeking to protect small acreage plots from deer (see Hildreth, A.M., S.E. Hygnstrom, and K.C. VerCauteren. 2013. *Deer-activated bioacoustic frightening device deters white-tailed deer*. Human-Wildlife Interactions Vol. 7(1):107-113, available at <https://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=1128&context=hwi>). Tests examining the efficacy of bioacoustics revealed that entry by deer to a protected area was reduced by 99% and acquisition by deer of bait materials placed within the plot was reduced by 100% following deployment of the device. Pre-recorded playback of distress calls made by deer, triggered by activation of motion detectors as deer approached the test area, caused deer to immediately flee the area. There were limitations on the size of the area over which protection successfully was extended by a single device, so the suitability of this approach to large acreages likely would be reduced.

- **Reproductive abatement:** Although there is great interest in and much research being conducted on the use of sterilization or contraceptives (*i.e.*, hormones administered to female deer to disrupt reproductive behaviors), only specially trained wildlife professionals are permitted to administer this treatment. To date, no effective reduction in population numbers, and thus a concurrent reduction in damage, has been achieved using contraceptives in free-roaming populations of deer. Success has been realized only in isolated contained populations where access to nearly all members of the population can be attained (*e.g.*, on islands, in confined city parks, etc.). This is a labor-intensive and costly strategy, and because individuals consistently move into and out of a population, it is extremely difficult to treat a sufficient number of individuals or to know which individuals already may have been treated. Research to improve fertility control methods is ongoing.
- **Trap and transfer** methods involve capturing deer in a specific area and physically moving them to another location for release. There are several techniques for capturing deer, including box traps, Clover traps, netted cage traps, drive nets, drop nets, rocket nets, corral traps, net guns, and immobilization drugs delivered through a dart. This strategy is labor-intensive, costly, impractical at large scales due to a lack of suitable relocation sites and carries the risk of spreading disease. Most states ban the translocation of deer due to concern about the spread of disease and the potential for injury to trapped animals. The approach does little to resolve damage situations and may simply transfer habituated deer, and the damage problems they cause, to new areas. **This practice is not permitted in some jurisdictions (*e.g.*, Virginia). Consult your local Wildlife Management Authority.**
- **Regulated hunting** involves the use of hunters to harvest deer in accordance with defined open seasons, bag limits, and population objectives. Hunting legally takes place during any of the various deer hunting seasons (archery, muzzleloaders, shotguns, and general firearms) established by the state Fish and Wildlife authority. Regulated hunting is the most cost-effective and efficient method to manage deer populations and is the only effective means to manipulate deer numbers statewide. See your state Fish and Wildlife authority for details on when the regulated season is open. Many states offer additional deer tags (“bonus tags”) to farmers who then distribute them to hunters who offer to harvest additional deer beyond what normally would be allowed on an individual’s hunting license. Check with your local wildlife authority to see if this option is available.
- **Permits to Shoot**, commonly referred to as a “Depredation Permit” or “Kill Permit,” are issued by the state Fish and Wildlife authority to owners or lessees of land who are experiencing crop damage. These special permits are highly variable among jurisdictions, but, where available, allow growers a mechanism to manage damage situations during times of the year when the regulated hunting season is closed and “take” normally would not be allowed. Depredation permits can help regulate local deer populations, particularly in areas that receive only limited hunting pressure (*i.e.*, farms surrounded by residential properties). For more information or to apply for a depredation permit, contact your state Fish and Wildlife authority.
- **Controlled hunts** combine conventional deer hunting methods with more stringent controls and restrictions on hunter activities. Participants in controlled hunts are chosen by various methods, ranging from selection by random lottery of interested licensed hunters to a more rigorous process involving assessment of hunting skill and weapon proficiency. Specific restrictions and controls applied to hunting activities depend upon the needs and concerns of landowners, elected officials, and other stakeholders, but usually will be similar to hunting regulations imposed during the regular open deer hunting season.

Because deer populations range over multiple parcels or farms, management of deer numbers often cannot be implemented effectively on a single property. Research clearly indicates that greater success in attaining population objectives can be achieved by developing and implementing a comprehensive **Community-Based Deer**

**Management Program**, especially in environments where traditional population management methods are not an option. Under this program, the state Fish and Wildlife authority cooperates with municipal, county, and, if applicable, other federal agencies to provide technical assistance in developing alternative deer management options. Some options include employing baiting and sharpshooting, use of noise-suppressed firearms, and other methods that otherwise would not be legal under traditional hunting approaches. State authorities can issue permits to conduct special deer management activities in areas where regulated hunting is not possible or deemed safe. **Consult your local Wildlife Management Authority to see if this approach may be applicable to your area.**

## Feral Swine

The appearance of feral swine on the landscape likely is an unfamiliar threat that many commodity producers in the Mid-Atlantic region have never experienced before. However, new populations of hogs can appear overnight and spread rapidly as a result of illegal importation and release of animals trapped elsewhere, creating a serious threat for producers. Whether from direct foraging or rooting behavior, the amount of damage inflicted by feral swine can be extensive and devastating in a short period of time, depending on the size of the sounder (*i.e.*, the family or social grouping) that takes up residence in the area.

Being an introduced, non-native species, the feral swine often has no protection under wildlife or game laws in many states, so hogs may be taken by legal methods at any time where such restrictions do not exist. However, in some states, feral swine are treated as “game” animals and are regulated via established open/closed seasons and bag limits on take. A substantial difficulty is the ability to accurately distinguish feral animals from true domestic swine that recently may have escaped, which constitute “property” to the swine owner. Thus, in some states, feral swine population management and damage mitigation have become complicated by questions of ownership and the legalities related to the take of property - thus, caution is urged before any hog removal action is implemented.

That said, growers confronted by the presence of feral swine should immediately contact the USDA-APHIS Wildlife Services Office that serves their state and request assistance in devising an effective hog eradication plan. Trap and bait techniques have proven to be the most reliable and successful approach to manage hog populations, but such operations take time and persistence to achieve the desired outcome. Shooting alone often proves ineffective in that only a small number of hogs in a social group can be taken at once and those that escape quickly learn to avoid hunters in the future. Although on-going research on potential toxicants has been encouraging, registration of an approved formulation remains years away. At this time, keen observation, and attentiveness to the first signs of hog presence are imperative to stay ahead of this threat.

## Groundhogs

The most obvious signs of groundhog (*Marmota monax*) presence, aside from actually seeing the animal or the symptoms of the damage they cause, are the entrances to a groundhog burrow system. Groundhog burrow systems are characterized by a large mound of excavated earth distributed about the main entrance and several discretely located secondary exits. The diameter of the main entrance may measure 10-12 inches. There are usually 2 or more additional entrances to a burrow system, which are well hidden and do not display an apron of excavated soils distributed around the opening. Although groundhogs prefer leafy green vegetation, they will use many different crops throughout the growing season. Seasonal or cyclic reproductive patterns typically influence groundhog abundance and the extent of damage from year to year. Damage management options include:

- **Habitat modification**, such as thinning or removing dense vegetation that provides protective cover, typically is not a feasible strategy for minimizing groundhog damage at the commercial scale of most farms and rarely proves to be cost-effective.
- **Exclusion** with fencing can be an effective short- or long-term strategy, depending on the type of fence and the size of the area to be protected. An electric wire placed 3-4” above the ground can deter groundhogs from entering a protected area. However, a determined groundhog eventually will dig under the wire and gain access to the area. Woven wire mesh or chicken wire fencing provides a more permanent solution. Mesh openings should be < 2 inches, and the fence should extend at least 3-4 feet above the ground. The top 15 inches of the fence should extend outward at a 45° angle to prevent individuals from climbing over the top. To prevent groundhogs from digging under the fence, the bottom edge of the fence should be buried at least 10 inches beneath the ground, with an additional 6-8” section bent outward at the bottom of the trench. Groundhogs are excellent climbers, so fence posts should be placed on the inside of the fence and greater deterrence has been achieved where the fence material is not drawn taut or rigid but instead left somewhat loose.

## A. General Production Recommendations

- **Fumigants** are effective in reducing groundhogs. Gas cartridges (sodium nitrate) currently are registered for this purpose. An ignited gas cartridge is placed in the main entrance of the burrow system after all other entrances have been located and sealed. As the cartridge burns, thick smoky fumes are emitted and fill the burrow system once the main entrance also has been sealed. Burrows can be treated with gas anytime of the year, but this method is most effective in the spring before the first litter of young emerges. Gas cartridges can be purchased at most farm supply stores. A note of caution when using gas cartridges - because the gas cartridge must be ignited for proper use, a fire hazard does exist. Therefore, gas cartridges should not be used in burrows located under wooden sheds, buildings, or near combustible materials. New resident animals may recolonize an empty or recently vacated burrow system, so continued vigilance is recommended.

Aluminum phosphide tablets, placed deep inside the main burrow entrance, are another type of fumigant that can provide effective groundhog control. The tablets react with moisture in the soil, creating hydrogen phosphide gas. Soil moisture and tightly sealed burrow entrances are important for the fumigant to be used effectively. The tablets are approved for outdoor use on non-cropland and orchards. Aluminum phosphide should not be used within 15 ft of any occupied building or in areas where gas could escape into areas occupied by animals or humans. Aluminum phosphide is a Restricted Use Pesticide (RUP) and can be applied only by Category 7D certified pesticide applicators.

- **Trapping** is effective in removing particularly problematic individuals. However, dispersing groundhogs from the surrounding area quickly occupy vacant territories in good habitat. Steel leg-hold traps are illegal in some states, so check with your state wildlife agency to determine what types of traps are legal. In most states, it is legal to use a medium-sized live trap baited with a variety of baits (*e.g.*, lettuce, apples, or fresh beans) that will attract a groundhog's attention if placed close to the burrow entrance. Traps should be placed at main entrances or along major travel corridors and checked at least once every 24 hours. Once captured, the groundhog may be killed humanely or released elsewhere on the owner's property. If the intent is to release the groundhog off-site, many states regulate where and how the live animal is to be handled. No releases are allowed on federal, state, county, or municipal land. **Relocation and release of a captured animal on property the trapper does not own is not permitted in some jurisdictions (*e.g.*, Virginia), so consult your local Wildlife Management Authority before attempting a capture.**

- **Shooting** groundhogs that are damaging crops or farmland is approved at any time of the year. Although groundhogs are considered a game species in some states (it is a "nuisance species" in VA), farmers do not need a valid hunting license to shoot nuisance groundhogs on their own property. Growers should verify with the state wildlife agency which weapons are legal for this purpose in your state.

## Rabbits

Rabbits can damage vegetation by clipping branches, stems, and buds. Damage may become especially pronounced during periods of heavy snow or in the spring when plants are emerging from the ground. Vegetation that has been clipped by rabbits is characterized by a cleanly snipped, 45° angle cut at the point of damage. Rabbit tracks and their pelleted scat often are found in areas of recent damage.

Strategies suitable to mitigate rabbit conflicts include the following items worthy of consideration:

- **Cultural practices** and **habitat modification** techniques mostly are used to reduce or eliminate brush piles, heavy vegetation, and other types of cover in and adjacent to areas where protection is desired and that serve as potential rabbit nesting and hiding sites. Please recognize that removal of such cover may be detrimental to other desirable wildlife species that also depend on these habitat features for protection or shelter and their numbers may be reduced as a result. However, using such habitat modification techniques that enhance the success of a rabbit's predators (*e.g.*, fox, coyote, and raptors) will allow them to help regulate rabbit numbers. Planting alternative food crops in adjacent tracts (a practice referred to as decoy cropping) has been suggested as a means to deter rabbits from high-value crops, but this approach typically attracts or supports higher numbers of rabbits rather than deter their presence.

- **Exclusion** of rabbits using fencing can be effective. A 2-foot tall fence consisting of 1-inch or smaller metal or wire mesh fencing material (note: rabbits will gnaw through plastic) will eliminate most rabbit damage. In regions susceptible to winter snow cover, and especially where woody nursery plants are being produced, consider increasing the height of the fence to accommodate the depth of persistent snow cover anticipated in your area; rabbits can gain access to plants if snow cover tops over the fence. In all cases, the bottom of the fence should be buried 8-12 inches below the ground surface and another 6-8 inches of fence material should extend outward away

from the crop at a 90° angle at that depth, forming an L-shape to the base. Alternatively, vulnerable areas often can be protected using 2 strands of electrified wire fencing. The lower strand should be set close to the ground (*i.e.*, within 2-3 inches), with the second strand 2-4 inches above the lower strand. Control of vegetation below these wires with herbicide treatment typically is necessary to prevent its contact with wires that cause a grounding of electric current and to maintain full voltage flow for effective deterrence. Finally, rabbit guards made of metal hardware cloth with ¼- to ¾-inch mesh may be effective in protecting individual high value specimens. Rabbit guards should maintain a minimum separation of 2 inches away from the stem of the plant to allow for future growth. Do not allow weedy growth or debris to accumulate inside these screen guards as this creates an ideal environment for borer infestation and may attract voles. All guards should be anchored to the ground to prevent rabbits from nudging or lifting the guard upward and gaining access under the barrier.

- **Miscellaneous methods:** Harassment techniques, such as dogs and motion detector activated water sprayers, may provide short-term protection. A number of taste-based repellent products are registered for use on rabbits, many of which are formulated with capsaicin, thiram, or ammonium soaps of fatty acids. Other repellent products derived from fox or coyote urines are designed and registered as odor-based area repellents. However, some rabbit repellents are not approved for use on foods grown for human consumption, so check the product label and active ingredients before use. Finally, because rabbits are classified as a game species in many states, they are subject to hunting regulations and imposed limits, which typically restrict “take” only to periods during the normal open seasons for rabbits. Trapping of rabbits similarly may be regulated or require special authorizations for capture outside the established trapping season. As was true for handling other captured wildlife, many states prohibit the individual performing that capture from removing, transporting/relocating, and releasing said animals on property they do not own. Consult your state Wildlife agency prior to implementing any hunting or trapping program to assure compliance with existing regulations.

### Voles

It is important to determine which species of vole occurs in your crop production sites. Vole species most associated with depredation issues in the Mid-Atlantic region are the **meadow vole** (*Microtus pennsylvanicus*) and the **woodland vole** (*Microtus pinetorum*). Meadow voles, also called meadow mice, are about 5½ to 7½ inches long, with fur that ranges from gray to yellow-brown with black-tipped hairs; they also display a bi-colored tail. Woodland voles are about 4-6 inches long, have red-brown fur, and a tail about the same length as the hind foot. Vole populations are cyclic, where populations periodically grow rapidly, remain high for approximately 1-2 years, but then crash abruptly. It is during these peak times where the potential for significant crop damage is greatest.

Because voles remain active year-round, the damage they cause to crops can occur at any time, depending upon the crop. In vegetable crops, damage usually occurs in spring, as young plants are emerging. Voles are generalist herbivores, so they feed on roots, shoots, tubers, leaves, and seeds of many different plants. Meadow voles spend more time above ground than do woodland voles, but both species inflict serious damage by feeding on the subsurface root systems of plants. Aboveground damage frequently consists of gnawing and girdling of the main stems of woody perennial plants or the sprouts and suckers that emerge from the base of such plants. Meadow voles construct surface runways (approx. 1½ to 2 inches wide) under or within the accumulated organic matter and duff layer that exists in fields; these runs often terminate at a 1-inch diameter wide hole that drops into an underground burrow network. In contrast, pine voles remain underground and inflict damage in the form of root girdling, which often goes unnoticed until severe damage already has occurred, and the affected plant is in rapid decline. Both species are known for constructing burrows that follow trickle irrigation lines or areas where the soil has been loosened by mechanical planters.

Managing vole damage is notoriously difficult and, when implemented, often will be a time-consuming endeavor. Nevertheless, there are strategies for dealing with voles to keep populations in check and to minimize the resulting damage to more tolerable levels.

- **Cultural practices and habitat modification measures** are helpful in deterring vole populations. Voles avoid areas where few food resources and little protective cover exist. Control of ground cover vegetation with herbicides, mowers, or disking is effective, although voles will travel under the protection of snow cover in vegetation-free areas. Herbicides are the preferred method to eliminate sod. Cultural practices that reduce the amount of organic litter around the base of plants are essential. All areas should be kept clear of debris or stored objects on the ground (such as bags, boxes, pruned branches) because these items provide protection to voles and can hinder mowing and proper bait placement. Plastic or synthetic weed barriers will encourage the establishment of vole populations

## A. General Production Recommendations

beneath these devices, so their use should be avoided. Frequent mowing between rows should be implemented to reduce habitat and cover for rodents and to enhance the effectiveness of natural predators. Erecting raised wooden perches from which avian predators can hunt may increase natural mortality among voles due to the presence and success of hawks and owls.

- **Exclusion methods** usually are feasible only at small scales and to protect individual high-value specimens. Hardware cloth or woven wire fence stem guards ( $\leq \frac{1}{2}$ -inch mesh) can be used to protect the stem of a woody plant, similar to the process described for rabbits. However, voles often can tunnel up from below and emerge inside a guard, thereby negating the effort. Products composed of sharp-edged crushed rock or pumice granules can be used to line the bottom and edges of a planting hole and may deter voles from digging into the root ball. These measures require significant time and handwork during installation, so an analysis of cost-effectiveness is necessary before considering such methods.
- **Repellents** that contain predator urine (coyote and fox) may have some limited effectiveness in reducing vole numbers, primarily through the effects of induced stress on reproduction. However, repellents are expensive and offer only short-term relief from damage. Also, many repellents are not approved for use on plants grown for human consumption, so read product labels carefully before making any selection.
- **Trapping** may be useful only where vole damage is localized ( $<1$  acre). Place 2 to 3 snap traps per runway and positioned perpendicular to the runway with triggers centered in the runway for greater success. All traps should be covered by a weighted box or pail to prevent non-target captures. Box traps and multiple-catch mouse traps also have been used to trap voles. Because these traps hold individuals alive, any non-target animals captured can be released unharmed. Bait traps with small grains, seeds, diced or sliced fruit, or peanut butter. If a trap is unsuccessful for 2 consecutive nights, move the trap to another location.
- **Toxicants** often are the preferred method used to control large vole populations. Most of the toxicants registered for use on voles are classified as Restricted Use Pesticides (RUPs), which requires applicators to acquire and possess advanced certification for vertebrate application (Category 7D). Because the potential hazard for non-target loss is so high with these toxins, current product labeling stipulates these materials must be disseminated using properly located bait stations or in-burrow placement; broadcast delivery over the ground surface is not allowed. Even when carefully following label instructions, there remains a strong likelihood that non-target predator species may be exposed to potential rodenticide poisoning due to the residual presence of toxic material in the gut of the live animals they capture or the carcasses on which they feed. To ensure the legality of using a particular toxicant in your state, contact your state's Pesticide Control Program.

## 12. Pollination

Seed and fruit production in many vegetable crops is dependent on pollen transfer within or between flowers. In most cases, pollen transfer is accomplished by insects such as bees or flies, and it is often beneficial to release pollinating insects into the crop during the flowering stage to achieve desirable fruit set and mature quality. Some crops like cucurbits require multiple pollination events for normal fruit development. The size and shape of a mature fruit is usually related to the number of seeds, and each seed is the result of a pollination event. Generally, as the number of bee visits increases there will be an increase in fruit set, number of seeds per fruit, fruit weight, and improved fruit shape. In strawberries, sufficient pollination also results in fruits with a longer shelf life and better color. Delay in pollination affects the timing of fruit set, while lack of adequate pollination usually results in small or misshapen fruit in addition to low yields. Even some crops that are capable of self-pollination (*e.g.*, eggplant, lima beans, okra, peppers) often benefit from pollen transfer by insects.

### Integrated Crop Pollination

Bees are the most important group of insects for crop pollination. Today's approach integrates managed and wild bee species. More information is available in several online resources, *e.g.*,

- "What is Integrated Crop Pollination" at <https://youtu.be/yMP5dTDRi6g>,
- "The Integrated Crop Pollination Project" at <https://icpbees.org/tools-for-growers/>
- "Integrated Crop Pollination for Squashes, Pumpkins and Gourds" at <https://icpbees.org/wp-content/uploads/2014/05/Integrated-Crop-Pollination-for-Cucurbita-crops.pdf>

## A. General Production Recommendations

European honey bees (*Apis mellifera*) and commercial bumble bees are most used for managed pollination services because they can be moved. Populations of wild bees can also be important for vegetable pollination. Wild bees include bumble bees (*Bombus* species), the squash bee (*Xenoglossa pruinosa*), orchard bees (*Osmia* species), and many species of solitary bees, most of which nest in soil. Surveys of wild bees reveals over 500 species in the Mid-Atlantic U.S., but not all will necessarily be visiting any given crop. The community of managed or wild bees visiting a crop varies among crops and can be influenced by other flowering plants competing for these same bees.

Activity of managed or wild bees on crop flowers at the correct time will enhance pollination. Individual cucurbit and strawberry flowers are usually open and attractive to bees for a day or less. The opening of the flower, release of pollen, and commencement of nectar secretion normally precede bee activity, and the timing is coordinated with receptivity of the stigma. Pumpkin, squash, and watermelon flowers normally open around daybreak and close by noon, whereas cucumber, strawberry, and muskmelon flowers generally remain open the entire day. Pollination usually takes place on the day the flowers open due to the short periods of pollen viability and stigmatic receptivity.

Activity and behavior vary with the species of pollinator. Bumble bees are active over a wide range of weather conditions and can tolerate foraging in cooler temperatures. Honey bee activity is determined primarily by weather and conditions outside the hive. Honey bees rarely leave the hive when the outside temperature is below 55°F (13°C). Flights seldom intensify until the temperature reaches 70°F (21°C). Wind speed more than 15 mph seriously impedes bee activity. Cool, cloudy weather and threatening storms reduce honey bee flights. Squash bees are active soon after sunrise in July and August. Most of the feeding of female squash bees is completed by midmorning (9 or 10 AM) after which they return to their nests in the soil. Male squash bees will continue to feed on flowers, often overnight.

Populations of wild bee species vary in abundance from year to year. Regular pesticide applications may reduce the abundance and diversity of these pollinators, and some agricultural practices such as tillage may destroy wild bees that nest in the soil. The implementation of conservation tillage practices may reduce nest damage.

Most Mid-Atlantic states have completed or are in the process of implementing **State Managed Pollinator Protection Programs (MP3s)**. Consult your State Apiarist (Table A-6) or the Department of Agriculture for your State requirements. **See also section D.6.3.1 Protection of Pollinators in this guide for state-specific initiatives and regulations on pollinator protection from pesticides, including MP3s and labels/regulations/laws related to applications of pesticides and protection of pollinators.**

**Table A-6. State Apiarist Contact Information**

The state regulatory agency, state apiarist, and the web link to the managed pollinator protection plan made available by each state agency. New Jersey has recognized pollinator protection measures that can be found on the State bee inspection site linked below.

<p><b>Delaware Department of Agriculture, Jennylyn Redner</b> (<a href="mailto:jennylynn.redner@delaware.gov">jennylynn.redner@delaware.gov</a>), <a href="https://agriculture.delaware.gov/pesticide-management/pollinator-protection-plan/">https://agriculture.delaware.gov/pesticide-management/pollinator-protection-plan/</a></p>
<p><b>Maryland Department of Agriculture, Cybil Preston</b> (<a href="mailto:cybil.preston@maryland.gov">cybil.preston@maryland.gov</a>), <a href="https://mda.maryland.gov/plants-pests/Documents/MP3-Pollinator-Plan.pdf">https://mda.maryland.gov/plants-pests/Documents/MP3-Pollinator-Plan.pdf</a></p>
<p><b>New Jersey Department of Agriculture, Meghan McConnell</b> (<a href="mailto:Meghan.McConnell@ag.nj.gov">Meghan.McConnell@ag.nj.gov</a>), <a href="https://www.nj.gov/agriculture/divisions/pi/prog/beeinspection.html">https://www.nj.gov/agriculture/divisions/pi/prog/beeinspection.html</a></p>
<p><b>Pennsylvania Department of Agriculture, Karen Roccasecca</b> (<a href="mailto:kroccasecca@pa.gov">kroccasecca@pa.gov</a>), <a href="https://pollinators.psu.edu/bee-health/pennsylvania-pollinator-protection-plan-p4#:~:text=The%20Pennsylvania%20Pollinator%20Protection%20Plan,the%20varied%20landscapes%20in%20Pennsylvania">https://pollinators.psu.edu/bee-health/pennsylvania-pollinator-protection-plan-p4#:~:text=The%20Pennsylvania%20Pollinator%20Protection%20Plan,the%20varied%20landscapes%20in%20Pennsylvania</a></p>
<p><b>Virginia Department of Agriculture and Consumer Services, Keith Tignor</b> (<a href="mailto:keith.tignor@vdacs.virginia.gov">keith.tignor@vdacs.virginia.gov</a>), <a href="https://www.vdacs.virginia.gov/pdf/BMP-plan.pdf">https://www.vdacs.virginia.gov/pdf/BMP-plan.pdf</a></p>
<p><b>West Virginia Department of Agriculture, Shanda King</b> (<a href="mailto:sking@wvda">sking@wvda</a>) <a href="https://agriculture.wv.gov/wp-content/uploads/WV-Managed-Pollinator-MP3-Final-Draft.pdf">https://agriculture.wv.gov/wp-content/uploads/WV-Managed-Pollinator-MP3-Final-Draft.pdf</a></p>

## A. General Production Recommendations

### Commercially Available Honey Bees

For crops readily visited by honey bees, the most reliable way to ensure pollination is to own or rent strong colonies of European honey bees from a reliable beekeeper. European honey bees are the primary managed pollinators because colonies with large populations can be easily moved to the field each year. With the arrival of parasitic honey bee mites (mainly *Varroa destructor*) along with likely impacts of pathogens, insecticides, and fungicides, the numbers of overwintering European honey bee colonies have significantly decreased in the last 20 years. Abundant colonies of feral honey bees (wild colonies nesting in trees or other cavities) are now uncommon to rare in most areas, and beekeepers are losing large numbers of colonies to mites, disease, and other stress factors. As a result, fewer beekeepers are providing honey bee colonies for pollination services, and some colonies may be of marginal quality for pollination.

The **Mid-Atlantic Apiculture Research and Extension Consortium** is a regional group focused on addressing the crisis facing the beekeeping industry (<https://canr.udel.edu/maarec/about-2/>). Additional relevant websites are the Center for Pollinator Research (<https://ento.psu.edu/pollinators>), the Honey Bee Health Coalition (<https://honeybeehealthcoalition.org/>) and Project *Apis m.* (<https://www.projectapism.org/>).

A brief introduction to best management practices for honey bees can be found at [https://honeybeehealthcoalition.org/wp-content/uploads/2019/01/HBHC\\_Hive\\_BMPs\\_v1.0\\_reduced.pdf](https://honeybeehealthcoalition.org/wp-content/uploads/2019/01/HBHC_Hive_BMPs_v1.0_reduced.pdf), and more guides at <https://www.projectapism.org/resources-2>. Best practices include: (1) Locating colonies in areas with sufficient flower forage and protected from exposure to sunlight (an east or southeast hive entrance encourages bee flights); (2) Elevating the colony to have the front entrance free of grass and weeds; and (3) Allowing a clean water supply within a quarter mile of the hive.

The number of colonies per acre for adequate pollination varies with location, attractiveness of the crop, density of flowers and length of blooming period, colony strength, and competitive plants in the area. In vine crops and strawberries, recommendations are to place 1 to 2 colonies per acre, with more hives required for higher density plantings.

### To ensure adequate quality and numbers of honey bee colonies, growers should:

- **Have a written and signed contract between the grower and the beekeeper.** This will ensure that enough pollinators are provided and that beekeepers are protected from pest control practices that may injure bees. The contract should specify the number and strength of colonies, rental fee, time of delivery, and distribution of bees in the field, as well as a plan to manage weeds that may act as competitive bloom. A sample contract is provided at [https://www.projectapism.org/s/2017\\_pollination-contract-template.pdf](https://www.projectapism.org/s/2017_pollination-contract-template.pdf). When entering into a pollination contract, there are some basic guidelines for beekeepers and growers alike (for more detailed information, see <https://honeybeehealthcoalition.org/growers-and-beekeepers-role/>).
- **Obtain an adequate number of colonies.** This varies among crops, location, attractiveness of the crop, density of the flowers and length of the blooming period, colony strength, and competitive plants in the area. A rule of thumb is to start with one colony per acre and adjust from there. Areas well populated with wild bees will not need as many rented honey bee hives.
- **Obtain bees at the appropriate time.** For melons, cucumbers, squash, and strawberries, honey bees should be moved in when the crop is flowering adequately to attract bees. Competing food sources from other flowers (e.g., dandelions), should be eliminated by mowing, cultivation, or herbicides (prior to bees being moved there).
- **Locate colonies for maximum effect.** Place colonies in groups of 4 to 8 in favorable locations throughout the farm or field to provide an even distribution of the bees. In large fields, pollination is effective if groups of 10 to 20 hives are distributed in sunny, wind-protected spots. Bales of straw or packing boxes stacked behind colonies offer wind protection. Be aware of the pollination requirements of your specific crop varieties. Some varieties may require “pollenizers” (sources of viable pollen for sterile varieties) to achieve adequate fruit set. Bees must be located so that they will encounter pollenizers and carry their viable pollen to the production varieties.
- **Rent honey bee colonies that are healthy and contain a large enough population to do the job.** Packaged bees (bees purchased through the mail) and small hives are inferior to strong, overwintered colonies. Two weak colonies are not equal to one strong colony. However, in some areas, colony loss has been so high that it may not be realistic to exclusively rely on overwintered colonies for pollination services. More information is available at <https://apiaryinspectors.org/US-beekeeping-survey-24-25>.

- **Contact beekeepers early.** Colonies may be in short supply. If you do not have a past relationship, make initial contact with the beekeeper the previous fall. Beekeepers usually assess the survival and strength of their colonies from mid-February to mid-March.
- **Requests for hive relocation should be given before application, ideally 48 hours or more. However, some states, such as New Jersey have notification regulations of 24 hours in advance, which is enforceable.** Some states have employed a hive registration program or an online voluntary registration tool that can help applicators locate and contact nearby beekeepers; contact your State Apiarist (see Table A-6).
- See **Section D. 6.3.1 Protection of Pollinators** for detailed information about FieldWatch online hive mapping services which has been sponsored by all Mid-Atlantic States. **Currently, the neonicotinoid pesticide labels prescribe a 48-hour notification in label-prescribed circumstances.**

## **Honey bee colony size and strength can be assessed in several ways:**

### **1. Inspect hives:**

This method is most time-consuming, but also most accurate. Colonies used for springtime pollination should have at least: a laying queen, 1½ or 2 stories (hive bodies or boxes), and 4 to 6 frames of brood; and enough adult bees to cover 6 to 8 frames. These are the minimum requirements. Stronger colonies with larger populations make superior pollination units and may command a higher price. As these stronger colonies are opened, bees will “boil out” or cover the tops of the frames. When smoked, however, the bees move down onto the frames and may not cover the frame tops. In this case, the frames themselves should be covered with bees. Note that there will be some variability in the quality of the colonies you rent. Generally, a group of colonies where 10% fall below the minimum standard is acceptable, only if also 10% are above the minimum standard. Also, for a variety of reasons, some colonies may become queen-less for a time; however, if these colonies meet all the other minimum requirements, they will still be effective pollination units.

You can request hives to be inspected. In most states, this is a general requirement for sale and interstate transport; see Table A-6 for your state apiarist contact information. Requests should be made as early as possible to facilitate scheduling. The beekeeper will be informed if an evaluation is requested by the grower. Colonies are inspected to determine the colony size (number of supers), the presence of a laying queen, the number of frames of brood and adult bees, and the presence of disease and parasites. At least 10% of the colonies in an apiary, or a minimum of 5 colonies, are selected at random for inspection. Inspected colonies are identified by stickers. If selected colonies are banded or stapled, these are not refastened by the inspector. A copy of the evaluation report is given to both the grower and the beekeeper.

### **2. Assess traffic at hive entrance:**

This method is less time-consuming but also less accurate. On a calm, warm (70-80°F, 21-27°C) day between 11 AM and 3 PM, bee traffic at hive entrances should be heavy. During a one-minute observation period, strong colonies should have 50-100 or more bees arriving and leaving the hive. Bees also should be seen arriving with pollen pellets on their back legs. In weak colonies, fewer than 40 bees will be seen arriving and leaving per minute. Colonies that are being used for summer pollination should have heavier traffic at the hive entrance.

Another crude way to assess colony strength is to observe entrances when temperatures are cool (55-60°F, 13-16°C). In strong colonies, flights will be observed at these cool temperatures, but in weaker colonies bees rarely fly when temperatures are below 60°F. Honey bees very rarely fly when the temperature is below 55°F.

### **3. Assess bee density on the crop:**

This method allows you to assess the contribution of feral or other honey bee colonies in the area in addition to rented bees. If you are using rented colonies, however, this method tells you little about the quality of the bees. We suggest that if you use this technique and find that the number of bees on the crop is small, you then use options (1) or (2) to assess colony strength before renting additional bees.

### **4 Additional information:**

Other sources of information for bee guides in your area are:

- State of NJ Department of Agriculture at <https://www.state.nj.us/agriculture/divisions/pi/>
- The Virginia Fruit Web site at <https://www.virginiafruit.ento.vt.edu/VAFS-bees.html>
- NCAT - ATTRA Sustainable Agriculture at <https://attra.ncat.org/>
- Farm Management for Native Bees, A Guide for Delaware at

## A. General Production Recommendations

<https://agriculture.delaware.gov/wp-content/uploads/sites/108/2017/12/FarmManagementforNativeBees-AGuideforDelaware.pdf>

- Managing Alternative Pollinators, A handbook for Beekeepers, Growers and Conservationists, SARE Outreach at <https://www.sare.org/resources/managing-alternative-pollinators/>

### Commercially Available Bumble Bees

Common Eastern bumble bee (*Bombus impatiens*) colonies may be purchased commercially to use as pollinators in vegetables and small fruits. The behavior, physiology and morphology of bumble bees make them ideal pollinators because of the speed at which they transfer pollen, the efficiency with which they gather pollen within various crops, and their ability to fly in adverse weather for longer periods of time. Bumble bees can also “buzz” pollinate, vibrating their wing muscles at a frequency that dislodges pollen from the flower, a technique not seen in honey bees. Due to their robust body size bumble bees begin foraging earlier and end later in the day and at lower temperatures. Bumble bees are effective in greenhouse and high tunnel settings to pollinate tomatoes and strawberries. They also have been successfully used for field pollination in blueberries and watermelon. However, in pumpkins, efforts to increase pollination by adding commercial bumble bee colonies are not always successful, perhaps due to the presence of adequate wild bee (wild bumble bee or squash bee) populations.

Place bumble bee colonies in the field after crops have begun to bloom. Bees that have found unintended forage in the beginning of the season are likely to continue to forage on this unintended source, especially if it is more favorable than the intended crop. Follow instructions provided by the bee supplier. Give the allotted time before opening up the colonies for the first time. Although bumble bees will need to excavate from natural enclosures to begin foraging, colonies should be given at least 30 minutes to settle after being handled during shipment and placement. Check each colony 2-3 hours later to ensure that the bees have successfully released and exited the nest. On occasion, bees are not released successfully and will need to be cut out.

Growers are urged to reduce each bumble bee colony entrance to one open hole at least two hours before each pesticide application. This will allow bumble bees to return to the hive and be kept in the colony to decrease exposure to pesticides. Bumble bees accumulate pesticides very easily within the wax and their bodies.

Place bumble bee colonies under shade to increase their productivity and longevity. Units placed in natural shade (along forest/field edges) or fitted with a shade structure last longer and are significantly more productive than those in full sunlight, especially during the warm summer months. Bumble bees constantly and actively strive to keep their colony temperature at around 86°F (30°C). Colonies exposed to direct sunlight use more energy for colony cooling.

Bumble bee colonies should be placed as far from honey bee hives as possible, especially when crops are not in bloom. When forage is low, colonies of pollinators should be more than 1 mile apart. Honey bees are very resourceful, and a bumble bee colony is a great source of pollen and nectar. If surrounding forage is low or not agreeable to honey bees, bumble bees will be susceptible to honey bee pollen theft resulting in weakened honey and bumble bee colonies. Bumble bees may be transferred to another field for additional pollination services throughout a season. Before moving, close the plastic opening tab to the one-hole open position. Allow forager bees at least two hours to return to the colony. The bumble bee colony may then be transferred to another site.

Follow the supplier’s recommendations for the number of hives to use in a particular crop. Commercial bumble bee hives live for 6 to 12 weeks (about 3 months) and must be replaced each year. Dispose of bumble bee colonies in a timely and humane fashion. There is a risk of commercial bees breeding with native populations. Commercial bumble bees are mass reared, and therefore have less genetic diversity than native bees. The genetic integrity of wild bees is important because it allows for adaptation to a wide variety of environmental conditions and various pathogens that they may encounter. Disposal of commercial colonies may also minimize potential transmission of pathogens.

**All bees are vulnerable to many chemicals used to control insects, pathogens, and weeds.**

#### **Related to pesticides and the use of commercial pollination services:**

- Avoid pre-bloom pesticides just before bees are brought onto a crop. If one is needed pre-bloom, select a material with lower bee toxicity and apply only when bees are not foraging, preferably late evening.
- Do not apply pesticides post-bloom until after managed colonies are removed.

- Honey bees need water for temperature regulation and brood production. Provide a clean water supply near the hives. Keep wheel ruts and areas around the sprayer fill point drained to eliminate a possible insecticide-laden water source.

## Wild Bees

Many wild bees, including squash bees (*Xenoglossa (Eucera) pruinosa*), multiple bumble bee species (*Bombus* sp., predominantly *Bombus impatiens*), orchard bees (*Osmia* sp.) and an assortment of other solitary bees (sweat bees, mining bees) are excellent crop pollinators. In the Mid-Atlantic regions, wild pollinators have provided sufficient pollination for small, diversified farms located in complex landscapes that include wood lots and unmanaged (fallow) lands in close proximity. The landscape can strongly influence bee populations through the availability of nesting substrates (open soil, fallen logs, abandoned rodent burrows). In diversified farmscapes with a history of growing cucurbits, bumble bees and/or squash bees have provided sufficient pollination to pumpkins regardless of whether managed commercial bees were present. Landscapes utilizing conservation tillage tend to have higher populations of squash bees, presumably due to less habitat disruption.

Availability of additional food resources in nearby wild lands or a diverse (flowering) cropping system can help support wild bee populations throughout the growing season. The USDA National Resources Conservation Service is building efforts to supplement farms with perennial plantings (pollinator strips) or cover cropping schemes designed to provide timely floral resources.

Wild bumble bees live in colonies founded by a queen. The workers, who are daughters of the queen, do the foraging, brood-rearing and defend the nest. New queen bumble bees (called “gynes”) emerge from their natal nest in late summer or autumn. Each gyne will mate, forage, and then hibernate through the winter in a small, insulated cavity. In the spring the gyne will emerge and search for a larger cavity to establish her nest in such as an old rodent nest or beneath clumps of bunchgrass. Colonies will increase in numbers over the spring and summer, reaching a peak of 250-450 individuals (in *Bombus impatiens*) before producing new gynes and males. These new reproductive individuals will disperse and start the cycle over, while their natal colony dies out, leaving the gynes as the only carry-overs to the next year.

Most native bees do not live in groups like honey and bumble bees. Each female solitary bee establishes her own nest which may be in the ground, an old beetle burrow in wood, or in a pithy stem (elderberry or brambles). Each female gathers pollen and nectar and feeds nest cells, making a pollen ball and laying a single egg in each cell. She repeats this process many times over the duration of her life and will die before her offspring mature. The offspring overwinter in the cell within the nest, emerging the following spring or summer. Female solitary bees are reliable pollinators, visiting many flowers in their lifetime.

Snags or brush piles, along with undisturbed tall grassy areas, provide nesting sites for tunnel-nesting bees and bumble bees. Hedgerows, shelterbelts, and windbreaks containing flowering trees and shrubs can provide nesting habitat for bees as well as food. Deep soil tillage can block or harm ground-nesting bees.

Bees can vary greatly in their foraging range depending on body size and resource availability. Large species like bumble bees can fly long distances, but probably forage within 1 to 3 miles from the colony. Most species stay closer to their nest, no farther than about 0.5 mile. When resources are plentiful, bees are more likely to forage over shorter distances. It may be advantageous to manage farmscapes with these pollinators in mind, reserving bee habitat to benefit the crops and surrounding landscape.

### Information for managing wild bees, along with the biology of relevant species can be downloaded at:

- Farm Management for Native Bees, A Guide for Delaware at <https://agriculture.delaware.gov/wp-content/uploads/sites/108/2017/12/FarmManagementforNativeBees-AGuideforDelaware.pdf>
- Using integrated crop pollination for pumpkins and squash: <https://icpbees.org/wp-content/uploads/2014/05/Integrated-Crop-Pollination-for-Cucurbita-crops.pdf> and <https://content.ces.ncsu.edu/squash-bees-in-the-home-garden>
- Squash Bee Biology, Biology and Pollination Services of the Squash Bee, *Eucera (Peponapis) pruinosa* at <https://lopezuribelab.com/squash-bee-biology/>, and <https://content.ces.ncsu.edu/squash-bees-in-the-home-garden>.
- Managing Alternative Pollinators, A handbook for Beekeepers, Growers and Conservationists, SARE Outreach at <https://www.sare.org/resources/managing-alternative-pollinators/>.

## A. General Production Recommendations

### Collections of resources are compiled at:

- The Integrated Crop Pollination Project, Resources for Growers, <https://icpbees.org/tools-for-growers/>
- The Center for Pollinator Research, <https://ento.psu.edu/pollinators/information-for-growers>.  
There is ongoing research to determine whether reliance on wild bees will be adequate for pollination of large acreages grown for commercial production.
- The Xerces Society provides guidelines for developing landscapes and farmscapes that encourage conservation of communities of pollinators at <https://www.xerces.org/pollinator-conservation/>.

#### NOTE:

**In this guide, this section addresses Pollination while Section D.6.3.1 Protection of Pollinators addresses specific practices for growers to follow to protect all pollinators from pesticide applications.**

**Section D.6.3.1 also includes an in-depth discussion of Product-Specific Labeling for the neonicotinoid class of pesticides, current state-specific pollinator protection activities, including MP3s, laws, and regulations.**

## 13. Food Safety Concerns

Reports of foodborne illness attributed to consumption of fresh fruits and vegetables have increased. Unlike processed foods, fresh fruits and vegetables are not heat-treated to eliminate potentially harmful microorganisms. Larger and more centralized farming and improved storage methods have resulted in the distribution of produce over vast geographic areas. Raw fruits and vegetables are also handled more frequently in the distribution chain. Cases of foodborne illness that once were limited to localized areas can now be spread over many states or countries. In addition, new minimal processing technologies have been brought to the marketplace, for example fruits and vegetables that have been washed, peeled, and cut into convenient ready-to-eat products. Since these products are subject to more handling and typically are not heat-processed to eliminate harmful bacteria, they are at a greater risk of becoming contaminated and subsequently leading to foodborne illness. Most fresh fruits and vegetables are grown, harvested, and packed under safe and sanitary conditions. However, several highly publicized cases of foodborne illness have been associated with consumption of lettuce, salad mixes, green onions, tomatoes, sprouts, cantaloupe, cabbage, cucumbers, herbs, and carrots. Implicated in most of these outbreaks have been the human pathogens: *Salmonella enterica*, *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Shigella* bacteria; *Cryptosporidium* and *Cyclospora* parasites; and Hepatitis A and Norovirus viruses.

In response to increasing concerns about the safety of fresh produce grown in the United States, the Food and Drug Administration (FDA) published “The Guide to Minimize Microbial Food Safety Hazards for Fresh Fruits and Vegetables” in 1998. This guide is intended to assist growers, packers, and shippers of unprocessed or minimally processed fresh fruits and vegetables by increasing awareness of potential food safety hazards and providing suggestions for practices to minimize those hazards. Many internet resources on food safety are also available that feature updated information from this guide and other sources.

In 2002, the United States Department of Agriculture (USDA) developed an audit/certification program known as “Good Agricultural Practices” (GAPs) to verify conformance to the 1998 guide. This is a voluntary program, although an increasing number of distribution networks are mandating GAPs or Harmonized Audit certification from each participating grower.

More recently, in 2011, the Food Safety Modernization Act (FSMA) was signed into law. There are seven sections to FSMA with the Produce Safety Rule applying to many growers and packers. The final Produce Safety Rule (under FSMA) was released November 2015, with the first compliance date in January 2018 with complete compliance required by 2020 except for sections under review, *i.e.*, water. The Produce Safety Rule (<https://www.fda.gov/food/food-safety-modernization-act-fsma/fsma-final-rule-produce-safety>) establishes mandatory practices operations must take to prevent microbial contamination of fresh produce. Whether a produce operation needs to comply with the Produce Safety Rule depends on whether it produces fresh fruits and vegetables and sales volume. To help operations prepare for an inspection, state departments of agriculture and cooperative

extension in each state are providing individual farm assessments (On Farm Readiness Review). These assessments are free, confidential and take on average 2 hours to complete.

**In the current food safety climate, increased record-keeping, and adherence to strict procedures of human hygiene are inevitable. All three resources (the 1998 guide, GAPs and FSMA Produce Safety Rule) identify potential hazards and discuss possible control methods in various aspects of pre-harvest, harvest, and post-harvest production, including: 1. Water, 2. Manure and Municipal Biosolids, 3. Wild and Domesticated Animals, 4. Worker Health, Hygiene and Training, 5. Sanitation, 6. Transportation, and 7. Product Trace-back. Each section is summarized below.**

### **1. Water:**

Water has the potential to be a source of microbial contamination. Growers and packers should be aware of the source and quality of water that contacts fresh produce and consider practices that will protect water quality. Growers should periodically test irrigation water for the quantity of fecal indicator organisms specifically generic *E. coli* (often represented by colony forming unit (CFU) or most probably number (MPN) of generic *E. coli* per 100 ml water). Frequency can be based on regulations and buyer requirements. For example, groundwater should be tested at least once per year and surface water three times per year under most Good Agricultural Practices programs). If fecal indicator organisms in irrigation water exceed the agricultural water standards, water treatment with effective disinfectants will be necessary before continuing to use the water source. SaniDate 5.0 or 12 and calcium hypochlorite tablets (Accutab) have been shown to be effective on the decontamination of bacterial foodborne pathogens. These products are approved by the Organic Materials Review Institute (OMRI) for use in irrigation water. Check the label to make sure the product can be used for this purpose; **the label is the law!**

Growers often irrigate field crops using surface water, water obtained from rivers, lakes, ponds, or irrigation ditches. However, surface water can become contaminated by upstream animal operations, sewage discharge, or runoff from fields. Drip, trickle, underground, or low volume spray irrigation techniques are ways to minimize irrigation water contact with harvestable portions of the crop. Groundwater is less likely to become contaminated than surface water, although wells should be maintained in good working condition including proper backflow devices and be constructed and protected so that surface water or runoff from manure storage areas cannot enter the system.

During harvest and post-harvest operations, when water is used in the field during the harvest process or in postharvest when water is used to cool, wash, or move produce, only water that meets the standard of no detectable generic *E. coli* based on a 100 ml water sample can be used. Untreated surface water cannot be used for harvest and post-harvest activities. Water in dump tanks and flume systems should be changed regularly to prevent the buildup of organic materials. Contact surfaces should be cleaned and sanitized to help prevent cross-contamination. Sanitizers, such as chlorine and peroxyacetic acid may be added to water but should be routinely monitored and recorded to ensure they are maintained at appropriate levels (*e.g.*, water should be monitored for proper chlorine efficacy; 100 to 150 ppm of free chlorine, and a pH in the range of 6.5 to 7.5).

### **2. Manure and Municipal Bio-solids:**

Manure may be contaminated with human pathogens and should be properly treated and stored before field application. Store manure and compost away from produce fields and packinghouses to protect the produce crop from seepage and runoff. Physical barriers such as ditches, mounds, grass/sod waterways, diversion berms, and vegetative buffer areas may also help to prevent runoff. Current recommendations are to maximize the time between application of manure to production areas and harvest. For non-composted or raw manure, the recommendation is to wait at least 120 days (4 months) between manure application and harvest and at a minimum two weeks before planting. Growers should be aware that the FSMA Produce Safety Rule regulations for manure of animal origin have not yet been finalized. The Food and Drug Administration allows the use of the National Organic Standards for manure application until the final regulations are written. The National Organic Standards are 1. Incorporated into the soil at least 120 days (about 4 months) before harvest when the edible portion of the crop has soil contact; OR 2. Incorporated into the soil at least 90 days (about 3 months) before harvesting all other food crops. Recommendations, guidance, regulations may change; growers are encouraged to consult relevant online resources or county extension offices about up-to-date manure recommendations and regulations.

Although municipal bio-solids (sewage sludge) are approved for certain agricultural uses, they are not recommended for application to soils used for vegetable production. This is due to the potential for human health issues. See “Sewage Sludge” in section B 4 Nutrient Management.

## A. General Production Recommendations

### **3. Wild and Domesticated Animals:**

Wild animals, although more difficult to control, should be discouraged from entering fields; especially where crops are destined for fresh markets. Wildlife prevention may include noise makers, decoys, hunting, fencing, or netting. However, the FDA does not authorize farms to take action(s) that would violate the Endangered Species Act or other federal, state, or local animal protection requirements (check with county extension on animal protection requirements). Domestic animals (including livestock and pets) may be a source of contamination and should be excluded from fields during the growing and harvesting season. Growers who use animals (such as horses) during production are advised to do a risk assessment of their operation and have a written plan in place to address possible sources of contamination.

### **4. Worker Health and Hygiene:**

Human pathogens can be transferred to produce by workers who harvest or pack fresh produce. Growers should provide sanitary facilities that are accessible, clean, and well equipped (bathrooms or portable toilets with an adequate supply of toilet paper; handwashing stations with basin, microbially safe water, soap, disposable paper towels or other appropriate hand drying devices, and a waste container). All employees (field workers to office administration) should be trained in good hygiene practices, such as toilet use and proper handwashing. Any worker who shows signs of an illness including diarrhea, coughing, fever, sneezing, sores, or infected wounds should not be allowed to handle produce.

### **5. Sanitation (includes field harvesting and packing facilities):**

Fresh produce can become contaminated through contact with soils, pests, equipment, and chemicals. Harvesters, knives, pruners, pallets, containers, or bins, among others, should be cleaned and sanitized before use. They should be replaced if they cannot be cleaned or are damaged (poor condition). Equipment, packing, and storage areas should be kept clean; empty or unused pallets, bins, or containers should be kept in a covered location to prevent contamination. Additionally, all equipment should be regularly serviced and inspected for general maintenance. Food contact surfaces should be cleaned and sanitized at the end of each day. A pest control program must be established to prevent or limit rodents, birds, and insects from nesting in equipment and entering the packing and storage facilities/areas. Sanitizers, such as chlorine or peroxyacetic acid, may be added to water to prevent cross-contamination of produce during washing or transporting in dump tanks and flumes. If using a sanitizer, monitor the concentration on a regular schedule. It is recommended the water be changed when it becomes saturated with organic material increasing its turbidity. For more information on sanitizers labeled for produce and the difference between cleaning and sanitizing visit the Produce Safety Alliance website at <https://cals.cornell.edu/produce-safety-alliance/resources#sanitation>.

### **6. Transportation:**

Fresh produce can become contaminated during loading, unloading, and shipping. Inspect transportation vehicles for cleanliness, pests, odors, and obvious dirt or debris before loading. Make sure that fresh produce is not shipped in trucks that have previously been used to transport animals, fish, chemicals, or waste. Refrigeration units in trucks should be turned on before loading to ensure that proper temperatures are maintained during loading and transport.

### **7. Trace-back:**

Traceability is defined as a procedure which tracks where a food product came from (for example farm, field, row, date harvested) to where a food product is going (market, distribution center, consumer). Usually adequate trace-back procedures require a grower to track one step backwards and forwards. Growers should be able to trace each lot with the date of harvest, farm identification, and who handled the produce from grower to receiver. The ability to trace the distribution history of food items from grower to consumer will not prevent a foodborne outbreak or recall from occurring; however, traceability procedures may limit the public health and economic impacts of an outbreak or recall.

**Additional information to help vegetable growers adopt Good Agricultural Practices on the farm and in the packinghouse; as well as information on the FSMA Produce Safety Rule can be obtained from extension offices or the governmental agriculture authority in your state.**

