



This is a section from the

2020/2021

Mid-Atlantic

Commercial Vegetable

Production Recommendations

The recommendations are **NOT** for home gardener use.

The **full manual**, containing recommendations specific to New Jersey, can be found on the Rutgers NJAES website in the Publications section: <http://njaes.rutgers.edu/pubs/publication.asp?pid=E001>.

This manual will be revised biennially. In January 2021, a **critical update** with important updates to the 2020/2021 manual will be communicated through local Extension Agents and Vegetable Specialists.

The **label** is a legally-binding contract between the user and the manufacturer. The user must follow all rates and restrictions as per label directions. The use of any pesticide inconsistent with the label directions is a violation of Federal law.

Cooperating Agencies: Rutgers, The State University of New Jersey, U.S. Department of Agriculture, and County Boards of Chosen Freeholders. Rutgers Cooperative Extension, a unit of the Rutgers New Jersey Agricultural Experiment Station, is an equal opportunity program provider and employer.

F. Commodity Recommendations

Pesticide Use Disclaimer

THE LABEL IS THE LAW

Before using a pesticide, check the label for up to date rates and restrictions.

Labels can be downloaded from: <http://www.cdms.net/>, <https://www.greenbook.net/> or <http://www.agrian.com/labelcenter/results.cfm>

For more information on Pesticide Safety and the Pesticide Label see chapter D.

Guide to the Recommended Pesticide Tables in the Following Crop Sections:

1. Pesticides are listed by group or code number based on chemical structure and mechanism of action, as classified by the Weed Science Society of America (WSSA) for herbicides, the Insecticide Resistance Action Committee (IRAC) for insecticides, and the Fungicide Resistance Action Committee (FRAC) for fungicides.

If the number is in bold font, the product may have resistance concerns.

2. For restricted use pesticides, the restricted active ingredients are labeled with a *. (See section D 3.2.1 “Restricted Use Classification Statement” for more information).

3. In addition to the pesticides listed below, other formulations or brands with the same active ingredient(s) may be available. ALWAYS CHECK THE LABEL:

- a) to ensure a pesticide is labeled for the same use,**
- b) to ensure the pesticide is labeled for the desired crop, and**
- c) for additional restrictions.**

4. All pesticide recommendations are made for spraying a broadcast area of 1 acre (43,560 square feet). Adjust the rate for banded applications (for more information, see section E 1.3 Calibrating Granular Applicators).

5. Check the label for the maximum amount of pesticide per application and the maximum number of applications per year.

6. Bee Toxicity Rating (Bee TR): N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.

Labeled Applications Sites for Pumpkins									
		Plastic mulch production					Bare-ground production		
		Soil-Applied		Postemergence					
Herbicides	WSSA group number	Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest	Soil-applied	POST	Post-harvest
Sandea	2		YES		YES		YES	YES	
Curbit	3		YES				YES		
Prefar	8	YES	YES				YES		
Command	13		YES				YES		
Strategy			YES				YES		
Reflex*	14		YES		YES		YES**		
Dual	15		YES				YES**		
Select	1			YES				YES	
Select Max	1			YES				YES	
Poast	1			YES				YES	
Gramoxone*	22				YES	YES			YES

*Special Local Needs Label 24(c), be sure it is registered for the specific state and for the intended use.

**Dual and Reflex are labeled for bareground only if the spray is directed to the row middles.

1. Soil Applied						
Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12
<p>-Plasticulture row middles application only: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v or include a non-selective herbicide.</p> <p>-Bareground: apply broadcast after seeding but before crop emergence or no sooner than 7 days before transplanting.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds. Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field.</p> <p>-Do not apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum number of Sandea applications per year is 2 and do not exceed 2 oz/A during the crop season.</p>						
3	Curbit 3EC	1 to 3 pt/A	ethalfuralin	0.38 to 1.13 lb/A	--	24
<p>-Plasticulture: row middles only: apply as a banded spray after crop emergence or transplanting. Do not soil incorporate.</p> <p>-Bareground: apply broadcast after direct-seeding but prior to crop emergence; do not use on transplanted pumpkins.</p> <p>-Controls annual grasses and certain annual broadleaf weeds, including carpetweed and pigweed sp.</p> <p>-Use lower rate for coarse-textured soils or soils with low organic matter.</p> <p>-Where overhead irrigation is available, activate Curbit with 0.5 inch of irrigation within 2 days after application; if no irrigation or rainfall occurs within 5 days of application, activity of Curbit can be reduced.</p> <p>-Available as a pre-mix herbicide Strategy. Strategy at 3 pt/A= Curbit at 26 fl oz (0.6 lb ai) and Command at 8 fl oz (0.188 lb ai)</p> <p>-Maximum applications per season: not specified</p>						
3 + 13	Strategy 2.1SC	1.5 to 6 pt/A	ethalfuralin plus clomazone	0.39 to 1.58 lb/A	45	24
<p>-Plasticulture: row middles application only.</p> <p>-Bareground: apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Strategy is a prepackage mixture of Curbit 3EC and Command 3ME. Refer to individual products for comments.</p> <p>-Clomazone spray or vapor drift may injure susceptible crops and other vegetation, refer to Command 3ME for comments.</p> <p>-Do not apply prior to planting crop. Do not soil incorporate.</p> <p>-Certain crop varieties may have the potential for injury or loss with this product. Consult qualified crop advisors for information pertaining to varieties in your area. -Maximum applications per season: not specified.</p>						
8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	--	12
<p>-Plasticulture: under plastic: apply in a band under the plastic, immediately before laying the mulch. Allow 7 day before making transplant holes to allow condensation to incorporate the herbicide. Plasticulture: row middles application is labeled.</p> <p>-Bareground: apply preemergence or preplant incorporated.</p> <p>-Preemergence applications should be followed by irrigation within 36 h (apply enough water to wet the soil at least 2 to 4 inches deep). Preplant incorporated applications should be incorporated 1 to 2 inches deep (deeper than 2 inches will result in reduced weed control).</p> <p>-Provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters.</p> <p>-Do not apply within 45 days of harvesting squash.</p> <p>-Do not apply more than 6 lb ai/A per season.</p>						

1. Soil Applied - continued on next page

F Pumpkins and Winter Squash

1. Soil Applied - continued

13	Command 3ME	0.67 to 2 pt/A	clomazone	0.025 to 0.75 lb/A	45	12
<p>-Labeled for winter squash and processing pumpkins; not labeled for jack-o-lantern pumpkins.</p> <p>-Plasticulture: row middles application only. -Bareground: apply broadcast just before planting but before crop emergence, or just before transplanting. Use the lower rate when used on coarse-textured soils low in organic matter, when weed pressure is light, or to minimize herbicide carryover that could affect subsequent crops.</p> <p>-Controls annual grasses and many broadleaf weeds including common lambsquarters, velvetleaf, spurred anoda, and jimsonweed. Carpetweed, morningglory sp., pigweed sp., and yellow nutsedge will not be controlled. Higher rates will improve control (or expand number of species controlled) such as common cocklebur, common ragweed, or jimsonweed (refer to label for specific weeds and rates).</p> <p>-WARNINGS: Command spray or vapor drift may injure sensitive crops and other vegetation up to several hundred yards from the point of application. Do not apply adjacent to sensitive crops (see label) or vegetation, or under unfavorable wind or weather conditions. Command may limit subsequent cropping options, see the label.</p> <p>-Available as a pre-mix herbicide Strategy: Strategy at 3 pt/A= Command at 8 fl oz (0.188 lb ai) and Curbit at 26 fl oz (0.6 lb ai)</p> <p>-Maximum number of Command applications per year: 1.</p>						
14	Reflex 2SL	Rates vary, refer to the specific label	fomesafen	0.13 to 0.38 lb/A	32	24
<p>-For pumpkins ONLY. A Special Local Needs Label 24(c) has been approved for the use of Reflex 2SL to control weeds in pumpkins in DE, MD, NJ, PA, and VA (expires 12/31/2020 in DE, MD, VA, 12/31/2022 in NJ, and 12/31/2023 in PA). The use of this product is legal ONLY if a waiver of liability has been completed (see https://www.syngenta-us.com/labels/indemnified-label-login).</p> <p>-Rates differ by states, soil types, and planting method. Rates as low as 10 fl oz/A can cause injury on coarse-textured soils.</p> <p>-Plasticulture: row middles application only. Bareground: apply broadcast within 24 h after direct-seeding and follow with 0.2 to 0.5 inches of overhead irrigation at least 36 hr before pumpkin begin to crack through the soil. For transplants, apply Reflex and then irrigate with 0.2 to 0.5 inches of water and then transplant. Do not prepare transplant holes until after Reflex application and irrigation.</p> <p>-Foliar application of Reflex will severely damage or kill pumpkin. The potential of crop injury is greater on lighter textured soils combined with intensive irrigation programs or high amounts of rainfall, therefore, adjust rates accordingly.</p> <p>-Reflex provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. Pumpkin varieties may vary in their response to Reflex. Treat small acreages first to determine tolerance, especially when applying to a new variety.</p> <p>-Reflex rates lower than 16 fl oz/A may not provide full-season control and should be used with other herbicides and/or other methods of weed control. -Consider rotational crops when applying fomesafen. If crop is replanted, do not re-apply Reflex. Refer to 24(c) label for specifics on rotational restrictions.</p> <p>-Maximum for Reflex application in DE, MD, NJ, PA, and VA: 24 fl oz/A IN ALTERNATE YEARS.</p>						
15	Dual Magnum 7.62E	1 to 1.33 pt/A	s-metolachlor	0.95 to 1.27 lb/A	30	24
<p>-For pumpkins ONLY. Plasticulture: row middles application only. Bareground: apply as an inter-row or inter-hill spray, leaving 1 ft of untreated area over the row. Do not use as an over the top application. Do not soil incorporate.</p> <p>-Suppresses or controls annual grasses, yellow nutsedge, and certain annual broadleaf weeds including nightshade species.</p> <p>-Dual Magnum will not control emerged weeds. Cultivate and/or hoe or tank-mix with Gramoxone 2SL to control emerged weeds before treatment. -Use the lower rate on fields with coarse-textured soils low in organic matter. Use the higher rates on fields with fine-textured soil and those with high organic matter. Maximum applications per season: not specified.</p>						

2. Postemergence

Group	Product Name	Product Rate	Active Ingredient (* = Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC	6 to 8 fl oz/A	clethodim	0.094 to 0.13 lb/A	14	24
	Select Max 0.97EC	12 to 16 fl oz/A				
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.19 to 0.28 lb/A	14	12
<p>-Select 2EC: use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). Select Max: use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). Poast: use COC at 1.0% v/v.</p> <p>-The use of COC may increase the risk of crop injury when hot or humid conditions prevail. To reduce the risk of crop injury, omit additives or switch to NIS when grasses are small and soil moisture is adequate.</p> <p>-Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.</p> <p>-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled.</p> <p>-Controls many annual and certain perennial grasses, including annual bluegrass, but Poast is preferred for goosegrass control. For best results, treat annual grasses when they are actively growing and before tillers are present. Control may be reduced if grasses are large or under hot or dry weather conditions.</p> <p>-Repeated applications may be necessary to control certain perennial grasses. If repeat applications are necessary, allow 14 days between applications. Rainfastness is 1 h.</p> <p>-Do not tank-mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses. Do not apply more than 8 fl oz of Select 2EC in a single application and do not exceed 32 fl oz/A for the season; do not apply more than 16 fl oz of Select Max in a single application and do not exceed 64 fl oz/A for the season.</p> <p>-Do not apply more than 1.5 pt/A Poast in single application and do not exceed 3 pt/A for the season.</p>						

2. Postemergence - continued on next page

2. Postemergence - continued

2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12
<p>-Plasticulture: row middles application only.</p> <p>-Bareground: broadcast for bareground. Apply Sandea after the crop has at least 3 to 5 true leaves but before first female flowers appear and no sooner than 14 days after transplanting. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v (1 qt/100 gal).</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf; control of weeds taller than 3 inches may not be adequate. Sandea will not control common lambsquarters or eastern black nightshade if applied postemergence; for row middle application, tankmix with a non-selective herbicide to increase spectrum of control.</p> <p>-Sandea provides both residual and postemergence control of susceptible weed species.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field.</p> <p>-Do not apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Rainfastness is 4 h. Maximum number of Sandea applications per year is 2 and do not exceed 2 oz/A during the crop season</p>						
22	Gramoxone 2SL	1.95 pt/A	paraquat*	0.49 lb/A	14	24
<p>-A Supplemental Label has been approved for the use of Gramoxone 2SL for postemergence weed control in DE, MD, NJ, PA, and VA. -Row middles as a shielded application. Apply as a directed spray in a minimum of 20 gal spray mix/A to control emerged weeds between the rows after crop establishment. Include a nonionic surfactant at 0.25% v/v.</p> <p>-Use shields or hoods to prevent spray contact with the crop and low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. See the label for additional information and warnings.</p> <p>-Rainfastness is 30 min. A maximum of 3 applications per year are allowed.</p> <p>-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

3. Postharvest

Group	Product Name	Product Rate	Active Ingredient (* = Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 2.0	2.25 to 3 pt/A	paraquat*	0.56 to 0.75 lb/A	--	24
<p>-A Special Local Needs Label 24(c) has been approved in VA (expires 12/31/2022) and a Supplemental Label in DE for the use of Gramoxone SL 2.0 for postharvest application to desiccate the crop. Apply after the last harvest for bareground or plasticulture.</p> <p>-Always include an adjuvant. Spray coverage is essential for optimum effectiveness.</p> <p>-See the label for additional information and warnings.</p> <p>-Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed.</p> <p>-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

4. Other Labeled Herbicides These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name	Active Ingredient (* = Restricted Use)
14	Aim	carfentrazone
14	Vida	pyraflufen

Insect Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Insecticides

Seed Corn Maggots

The use of neonicotinoid insecticides (Group 4A) at planting may help reduce seed corn maggot populations. See also Maggots in section E 3.1 Soil Pests - Detection and Control.

Aphids Note: Aphids transmit mosaic virus.

Apply one of the following formulations: Note: Thorough spray coverage beneath leaves is important. Treat seedlings every 5-7 days, or as needed.						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M

Aphids - continued on next page

F Pumpkins and Winter Squash

Aphids - continued

4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil	21	4	M
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
29	Beleaf 50SG	2.0 to 2.8 oz/A	flonicamid	0	12	L

Armyworms and Cabbage Loopers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
11A	Dipel DF, others (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
11A	XenTari (OMRI) (armyworms)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
11A	XenTari (OMRI) (cabbage loopers)	0.5 to 1.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel (armyworms)	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Exirel (cabbage loopers)	10.0 to 17.0 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
28 + 4A	Voliam Flexi (cabbage looper only)	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Cucumber Beetles

Young plants need to be protected from cucumber beetle feeding as the beetles can transmit the causal agent of bacterial wilt. Cucumber beetles also cause direct damage to pumpkin and winter squash rinds. Management of adult cucumber beetles early in the season may help reduce damage to rinds later in the season. Seeds pretreated with a neonicotinoid seed treatment such as Farmore DI400 should provide up to 14 days of control of cucumber beetle. Otherwise, apply one of the following formulations:

Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
28	Exirel	20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Cutworms See also section E 3.1. Soil Pests - Detection and Control.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					

Leafminers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	3	4	M
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	H
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Melonworms and Pickleworms

Apply one of the following formulations. When using foliar materials make one treatment prior to fruit set, and then treat weekly. For soil or drip applications check the label for instructions on treatment frequency.						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Sevin XLR Plus	0.5 to 1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	2.0 to 3.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole	30	12	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Mites

Mite infestations generally begin around field margins and grassy areas. **DO NOT** mow or maintain these areas after midsummer to prevent mites from moving into the crop. Localized infestations can be spot-treated. Begin treatment when 10-15% of the crown leaves are infested early in the season.

Apply one of the following formulations. Note: Continuous use of carbaryl or pyrethroids may result in mite outbreaks. Addition of crop oils or organosilicon spray additives will increase miticide effectiveness.						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L
20B	Kanemite 15SC	31.0 fl oz/A	acequinocyl	1	12	L
21 A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M

F Pumpkins and Winter Squash

Rindworms

For Lepidopteran rindworms, apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L

Squash Bugs

Begin treatments if more than one egg mass per plant is present. Sprays should target nymphal stages.

Apply one of the following formulations: Note: Under-leaf spray coverage is essential.						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	10.5 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M

Squash Vine Borers

When vines begin to run, apply to bases of plants 4 times at 7-day intervals. Pheromone traps for squash vine borer are commercially available. These traps can be used to indicate when moth activity begins. Note: Use of spinosad or spinetoram for Cabbage Looper control will reduce squash vine borer populations.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					

Thrips

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	3	4	M
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Whiteflies

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil	21	4	M
7C	Knack	8.0 to 10.0 fl oz/A	pyriproxyfen	7	12	L
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	14.0 fl oz/A	afidopyropen	0	12	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
29	Beleaf 50SG	2.8 oz/A	flonicamid	0	12	L

Group 3A Pyrethroid Insecticides Registered for Use on Pumpkins and Winter Squash					
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):					
Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H
Baythroid XL	0.8 to 2.8 fl oz/A	beta-cyfluthrin*	0	12	H
Bifenthrin 2EC, others	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H
Danitol 2.4EC	10.67 to 16.0 fl oz/A	fenpropathrin*	7	24	H
Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H
Lambda-Cy 1EC, others	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
Mustang Maxx	1.28 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H
Permethrin 3.2EC, others	4.0 to 8.0 fl oz/A	permethrin*	0	12	H
Tombstone, others	0.8 to 2.8 fl oz/A	cyfluthrin*	0	12	H
Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
Combo products containing a pyrethroid					
Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam (Group 4A)	1	24	H
Gladiator	19.0 fl oz/A	zeta-cypermethrin* + abamectin* (Group 6)	7	12	H
Besiege	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole (Group 28)	1	24	H

Group 4A Neonicotinoid Insecticides Registered for Use on Pumpkins and Winter Squash					
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):					
Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
Assail 30SG	2.5 to 5.3 oz/A	acetamiprid	0	12	M
Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - soil/drip	21	12	H
Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - foliar (note: PHI: do not make application after 4 th true leaf has unfolded)	see note	12	H
Actara 25WDG	1.5 to 5.5 oz/A	thiamethoxam	0	12	H
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H
Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil/drip	21	12	H
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
Venom 70SG	5.0 to 7.5 oz/A	dinotefuran - soil/drip	21	12	H
Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
Combo products containing a neonicotinoid					
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole (Group 28)	1	12	H
Endigo ZC	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin* (Group 3A)	1	24	H

Disease Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Fungicides

Nematodes - See also sections E 1.5 Soil Fumigation and E 1.6 Nematode Control. Use fumigants listed in section E 1.5, or nematicides listed below. Consult the label.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Vydate L	0.5 to 1.0 gal/A incorporate into top 2-4 inches of soil, <i>OR</i> 2.0 to 4.0 pt/A apply 2 w after planting and repeat 2-3 w later.	oxamyl*	1	48	H
7	Velum Prime 4.16SC	6.5 to 6.84 fl oz/A	fluopyram	0	12	--
--	Nimitz 4EC	3.5 to 5.0 pt/A Incorporate or drip-apply 7 d before planting	fluensulfone	n/a	12	N

Seed Treatment

Check with your seed company if seed has been treated with an insecticide and fungicide. If it has not been treated, use a mixture of thiram 480DP (4.5 fl oz/100 lb seed) and an approved commercially available insecticide.

Damping-Off caused by Phytophthora, Pythium, and Rhizoctonia

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply one of the following at-planting (see label for application timing, methods, and restrictions):						
Phytophthora and Pythium root rot						
4	Ridomil Gold 4SL ¹	0.5 to 1.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E ¹	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4	MetaStar 2E AG ¹	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
Phytophthora, Pythium, and Rhizoctonia root rot						
4 + 11	Uniform 3.66SE	0.34 fl oz/1000 ft row. Avoid direct seed contact, which may cause delayed emergence.	mefenoxam + azoxystrobin	AP	0	N
Rhizoctonia root rot						
11	azoxystrobin 2.08F	0.40 to 0.80 fl oz/1000 ft row	azoxystrobin	AP	4	N
Pythium root rot only						
28	Previcur Flex 6F	1.2 pt/A in transplant water, drip irrigation, or direct spray at base of plant and soil	propamocarb HCl	2	12	N

¹To determine the amount of Ridomil Gold, Ultra Flourish, or MetaStar needed per acre, use the following calibration formula for changing from broadcast to band application: [Band width (ft) / row spacing (ft)] x broadcast rate (lb/A) = Amount needed lb/A. ²Applied at planting.

Bacterial and Fungal Diseases**Angular Leaf Spot/Bacterial Leaf Spot**

Both diseases can produce foliar symptoms that are often overlooked. Early detection is important, since control of the foliar phase can reduce infections in developing fruit. Infected fruit will become unmarketable. Both diseases are seedborne and can survive on infested debris for at least one year or until the debris decomposes. Rotate away from fields with a history of bacterial problems. Incorporate the following into a standard disease management program when leaf spot is first detected and repeat every 7 to 10 days: fixed copper at labeled rates plus mancozeb.

Anthracnose - see Gummy Stem Blight (Black Rot) and Anthracnose below.

Bacterial Wilt

Controlling striped and spotted cucumber beetles is essential for preventing bacterial wilt. See "Cucumber Beetles" in the Cucumber Insect Control section for specific recommendations. Insecticide applications made at planting may not prevent beetle damage season-long; additional foliar insecticide applications may be necessary.

Choanophora fruit rot

This disease occurs during warm wet weather and develops predominantly on flowers or fruit near the ground. Management is difficult because disease development is rapid and weather dependent. Fungicide sprays are not effective because flowers, which open daily, must be protected immediately. Practices that reduce soil moisture or reduce flower-soil contact, such as raised beds and plastic mulch, may be beneficial.

Downy Mildew

Scout fields for disease incidence on a regular basis. Begin targeted sprays when downy mildew is predicted for the region. For current status of the disease, check the Cucurbit Downy Mildew forecasting website at <http://cdm.ipmpipe.org/>. Strains of downy mildew that infect one cucurbit crop may not affect pumpkin or winter squash. Unnecessary fungicide application can be avoided by not spraying until disease is predicted in the region on watermelon. Preventative applications are much more effective than applications made after disease is detected. Materials with different modes of action (FRAC codes) should always be alternated to reduce the chances for fungicide resistance development.

Code	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
Sprays should be applied on a 7-d schedule when disease is forecast or present in the region. Under severe disease conditions and conducive weather, spray interval may be reduced IF the label allows. TANK-MIX one of these products WITH a protectant fungicide such as chlorothalonil 6F or Gavel 75DF:						
49+40	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
49+M05	Orondis Opti 3.37SC	1.75 to 2.5 pt/A	oxathiapiprolin + chlorothalonil	0	12	--
21	Ranman 400SC	2.10 to 2.75 fl oz/A (plus a non-ionic or organosilicon surfactant; do not apply with copper; see label)	cyazofamid	0	12	L

Downy Mildew - continued on next page

For Immediate Medical Attention

Call 911

**For a Pesticide Exposure Poisoning
Emergency Call**



For All States

This number will automatically connect you to the poison center nearest to you.

Anyone with a poisoning emergency can call the toll-free telephone number for help. Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

For Pesticide Spills

Small Spills: See the product label for cleanup advice.

Large spills: Call the National Response Center at 1-800-424-8802 or CHEMTREC at 800-424-9300 (24 hours) - Industry assistance with emergency response cleanup procedures for large, dangerous spills.

Be aware of your responsibility to report spills to the proper state agency.