



This is a section from the

2020/2021

Mid-Atlantic

Commercial Vegetable

Production Recommendations

The recommendations are **NOT** for home gardener use.

The **full manual**, containing recommendations specific to New Jersey, can be found on the Rutgers NJAES website in the Publications section: <http://njaes.rutgers.edu/pubs/publication.asp?pid=E001>.

This manual will be revised biennially. In January 2021, a **critical update** with important updates to the 2020/2021 manual will be communicated through local Extension Agents and Vegetable Specialists.

The **label** is a legally-binding contract between the user and the manufacturer. The user must follow all rates and restrictions as per label directions. The use of any pesticide inconsistent with the label directions is a violation of Federal law.

Cooperating Agencies: Rutgers, The State University of New Jersey, U.S. Department of Agriculture, and County Boards of Chosen Freeholders. Rutgers Cooperative Extension, a unit of the Rutgers New Jersey Agricultural Experiment Station, is an equal opportunity program provider and employer.

F. Commodity Recommendations

Pesticide Use Disclaimer

THE LABEL IS THE LAW

Before using a pesticide, check the label for up to date rates and restrictions.

Labels can be downloaded from: <http://www.cdms.net/>, <https://www.greenbook.net/> or <http://www.agrian.com/labelcenter/results.cfm>

For more information on Pesticide Safety and the Pesticide Label see chapter D.

Guide to the Recommended Pesticide Tables in the Following Crop Sections:

1. Pesticides are listed by group or code number based on chemical structure and mechanism of action, as classified by the Weed Science Society of America (WSSA) for herbicides, the Insecticide Resistance Action Committee (IRAC) for insecticides, and the Fungicide Resistance Action Committee (FRAC) for fungicides.

If the number is in bold font, the product may have resistance concerns.

2. For restricted use pesticides, the restricted active ingredients are labeled with a *. (See section D 3.2.1 “Restricted Use Classification Statement” for more information).

3. In addition to the pesticides listed below, other formulations or brands with the same active ingredient(s) may be available. ALWAYS CHECK THE LABEL:

- a) to ensure a pesticide is labeled for the same use,**
- b) to ensure the pesticide is labeled for the desired crop, and**
- c) for additional restrictions.**

4. All pesticide recommendations are made for spraying a broadcast area of 1 acre (43,560 square feet). Adjust the rate for banded applications (for more information, see section E 1.3 Calibrating Granular Applicators).

5. Check the label for the maximum amount of pesticide per application and the maximum number of applications per year.

6. Bee Toxicity Rating (Bee TR): N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.

Splitting during handling occurs in fruit under excessive water pressure as a result of excess irrigation or rainfall. **Sunscald** occurs when fruit are exposed to direct sunlight, especially on extremely hot days. Under these conditions, rind surfaces can reach temperatures exceeding 140°F (60°C), killing cells and resulting in sunburn spots. Fruit with little or no foliar cover are at most risk. Sunscald or sunburn first appears as a gray or white area on the exposed upper surface of the fruit. Fruit with dark rinds are more susceptible to sunscald than those with light colored rinds. Sunscald severity is related directly to fertility regime and foliage cover. Proper fertility and soil management promotes adequate vine growth and coverage of fruit. Sunscald severity is also associated with diseases that reduce foliage cover, such as anthracnose, alternaria, gummy stem blight and downy mildew. Recommendations for managing these diseases may be found in the Disease Control section below.

Water soaking occurs where excess water accumulates at the bottom of the fruit resulting in a water soaked appearance of internal flesh. Water accumulates during cloudy weather when transpiration from vines is low. Water soaking sometimes appears in fruits where foliage has deteriorated since excess water cannot be transpired.

Weed Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Herbicides

1. Identify the weeds in each field and select recommended herbicides. More information is available in the “Herbicide Effectiveness on Common Weeds in Vegetables” (Table E-2) in chapter E Pest Management.
2. Minimize herbicide resistance development. Identify the herbicide site of action group number and follow recommended good management practices; **bolded group numbers in tables below are herbicides at higher risk for selecting resistant weed populations.** Include non-chemical weed control whenever possible.

Labeled Applications Sites for Watermelon									
Herbicides	WSSA group number	Plastic mulch production					Bare-ground production		
		Soil-Applied		Postemergence			Soil-applied	POST	Post-harvest
		Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest			
Sandea	2	YES	YES		YES		YES		
Curbit	3		YES				YES		
Prowl H2O	3		YES						
Treflan	3		YES						
Sinbar	5	YES	YES				YES		
Prefar	8	YES	YES				YES		
Command	13		YES				YES		
Strategy	3 + 13		YES				YES		
Reflex*	14	YES	YES		YES		YES		
Dual*	15		YES						
Poast	1			YES				YES	
Select	1			YES				YES	
SelectMax	1			YES				YES	
Gramoxone*	22				YES	YES			YES

*Special Local Needs Label 24(c), be sure it is registered for the specific state and for the intended use.

1. Soil-Applied

Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	57	12
<p>-Plasticulture: can be applied in a band under the plastic, immediately before laying the mulch; delay seeding or transplanting for 7 days after application. Plasticulture row middles: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v or include a non-selective herbicide.</p> <p>-Bareground: apply broadcast after seeding but before crop emergence or no sooner than 7 days before transplanting.</p> <p>-Maximum rate for application in seeded or transplanted row is 0.75 oz/A, and up to 1 oz/A for row middle application.</p> <p>-Limit movement of treated soil into transplant hole during transplanting.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds. Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. -Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field.</p>						

1. Soil-Applied, Sandea - continued on next page

F Watermelons

1. Soil-Applied, Sandea - continued

<p>- Do not apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum Sandea applications per year is 2 and do not exceed 1 oz/A during the crop season.</p>						
3	Curbit 3EC	1 to 3 pt/A	ethalfluralin	0.38 to 1.13 lb/A	--	24
<p>-Plasticulture, row middles only: apply as a banded spray after crop emergence or after transplanting. Do not soil incorporate.</p> <p>-Bareground: apply broadcast after direct-seeding but prior to crop emergence; do not use on transplanted melons.</p> <p>-Controls annual grasses and certain annual broadleaf weeds, including carpetweed and pigweed sp.</p> <p>-Use lower rate for coarse-textured soils or soils with low organic matter.</p> <p>-Where overhead irrigation is available, activate Curbit with 0.5 inch of irrigation within 2 days after application; if no irrigation or rainfall occurs within 5 days of application, activity of Curbit can be reduced.</p> <p>-Available as a pre-mix herbicide Strategy. Strategy at 3 pt/A= Curbit at 26 fl oz (0.6 lb ai) and Command at 8 fl oz (0.188 lb ai)</p> <p>-Maximum applications per season: not specified</p>						
3	Prowl H2O 3.8CS	2.1 pt/A	pendimethalin	1 lb/A	35	24
<p>-Plasticulture: row middles only: apply as a banded spray before seeded crop has emerged or before transplanting.</p> <p>-Bareground: apply with shielded sprayer band between rows, leaving 6 inches of untreated area on both sides of the seeded or transplanted row. Apply before seeded crop emerges or before transplanting.</p> <p>-Where overhead irrigation is available, activate Prowl with 0.5 inch of rainfall or sprinkler irrigation within 48 hr of application; if no irrigation or rainfall occurs within 5 days of application, activity of Prowl can be reduced</p> <p>-A second application at the same rate may be applied to row middles as a banded spray postemergence a minimum of 21 days after the first application, but before the vines begin to run. Do not apply over the top of the crop, or severe injury may occur.</p> <p>-Maximum Prowl H2O applications per season is 2 and do not exceed 4.2 pt/A during the crop season.</p>						
3	Treflan 4EC	1 to 2 pt/A	trifluralin	0.5 to 1 lb/A	60	12
<p>-Plasticulture: row middles only: apply as a directed spray after emergence when plants have reached the 3 to 4 true leaf stage.</p> <p>-Not labeled for bareground production. Primarily controls annual grasses with a few broadleaf weeds.</p> <p>-Do not use (or reduce the rate) when cold, wet soil conditions are expected, or crop injury may result.</p> <p>-Maximum applications per season: not specified.</p>						
3 + 13	Strategy 2.1SC	1.5 to 6 pt/A	ethalfluralin plus clomazone	0.39 to 1.58 lb/A	45	24
<p>-Plasticulture: row middles application. -Bareground: apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Strategy is a prepackage mixture of Curbit 3EC and Command 3ME.</p> <p>-Clomazone spray or vapor drift may injure susceptible crops and other vegetation, refer to Command 3ME for comments.</p> <p>-Do not apply prior to planting crop. Do not soil incorporate.</p> <p>-Refer to individual products for comments. Maximum applications per season: not specified.</p>						
5	Sinbar 80WDG	2 to 4 oz/A	terbacil	0.1 to 0.2 lb/A	70	12
<p>-Plasticulture: can be applied in a band under the plastic, immediately before laying the mulch. Sinbar can be broadcast over the plastic before transplanting or before holes are made in the plastic; but must be washed off with a minimum of 0.5 inches for rainfall or irrigation before transplanting. Plasticulture row middles: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged include a non-selective herbicide.</p> <p>-Bareground: apply broadcast after seeding but before crop emergence. -Do not apply over the top of the crop or allow spray to contact crop foliage, or injury may result. -Controls many annual broadleaf weeds, but may be weak on pigweed species. Use the lower rate on coarse-textured soils low in organic matter and higher rates on fine-textured soils and on soils with high organic matter.</p> <p>-Maximum Sinbar applications per year is 2 and do not exceed 4 oz/A during the crop season</p>						
8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	--	12
<p>-Plasticulture: under plastic: apply in a band under the plastic, immediately before laying the mulch. Allow 7 day before making transplant holes to allow condensation to incorporate the herbicide. Plasticulture: row middles application is labeled.</p> <p>-Bareground: apply preemergence or preplant incorporated.</p> <p>-Preemergence applications should be followed by irrigation within 36 h (apply enough water to wet the soil at least 2 to 4 inches deep). Preplant incorporated applications should be incorporated 1 to 2 inches deep (deeper than 2 inches will result in reduced weed control).</p> <p>-Prefar provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters. -Do not apply more than 6 lb ai/A per season.</p>						
13	Command 3ME	0.4 to 0.67 pt/A	clomazone	0.15 to 0.25 lb/A	--	12
<p>-Plasticulture: row middles application only.</p> <p>-Bareground: apply broadcast just before planting or after planting but before crop emergence. Use the lower rate when used on coarse-textured soils low in organic matter, when weed pressure is light, or to minimize herbicide carryover that could affect subsequent crops.</p> <p>-Controls annual grasses and many broadleaf weeds including common lambsquarters, velvetleaf, spurred anoda, and jimsonweed. Carpetweed, morningglory sp., pigweed sp., and yellow nutsedge will not be controlled. Higher rates will improve control (or expand number of species controlled) such as common cocklebur, common ragweed, or jimsonweed (refer to label for specific weeds and rates).</p> <p>-WARNINGS: Command spray or vapor drift may injure sensitive crops and other vegetation up to several hundred yards from the point of application. Do not apply adjacent to sensitive crops (see label) or vegetation, or under unfavorable wind or weather conditions. Command may limit subsequent cropping options, see the label.</p> <p>-Available as a pre-mix herbicide Strategy: Strategy at 3 pt/A= Command at 8 fl oz (0.188 lb ai) and Curbit at 26 fl oz (0.6 lb ai)</p> <p>-Maximum Command applications per year is 1.</p>						

1. Soil-Applied - continued on next page

1. Soil-Applied - continued

14	Reflex 2SL	Rates vary, refer to the specific label	fomesafen	0.16 to 0.25 lb/A	35	24
<p>-A Special Local Needs Label 24(c) has been approved for the use of Reflex 2SL to control weeds in watermelon in DE, MD, NJ and VA (expires 12/31/2020 for DE, MD, VA, and 12/31/2022 in NJ). The use of this product is legal ONLY if a waiver of liability has been completed (see https://www.syngenta-us.com/labels/indemnified-label-login).</p> <p>-Rates vary by state and application method; refer to label to determine correct rates.</p> <p>-Plasticulture: can be applied in a band under the plastic at 10 to 12 fl oz, immediately before laying the mulch.</p> <p>-Plasticulture: Reflex at 10 to 12 fl oz can be broadcast over the plastic before transplanting or before holes are made in the plastic; but must be washed off with a minimum of 0.5 inches for rainfall or irrigation before transplanting.</p> <p>-Plasticulture row middles: before emergence of seeded crop or before transplanting; apply up to 12 fl oz in VA or up to 16 fl oz in DE and MD. Plasticulture row middles with shielded/hood sprayers after transplanting; apply 16 to 24 fl oz in DE and MD prior to vines "running" off the plastic. Severe crop injury can occur if spray comes in contact with crop foliage.</p> <p>-Bareground direct-seeded: apply broadcast within 24 h after seeding followed by 0.2 to 0.5 inch of overhead irrigation at least 36 h before watermelon crack the soil surface.</p> <p>-Bareground transplants: apply as broadcast spray followed by irrigation of 0.2 to 0.5 inches. Then prepare holes and transplant; avoid moving herbicide-treated soil into transplant holes.</p> <p>-Reflex provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. -Watermelon varieties may vary in their response to Reflex. Treat small acreages first to determine crop tolerance, especially when applying to a new variety. -Consider rotational crops when applying fomesafen. If crop is replanted do not re-apply Reflex. Rotational restrictions are dependent on whether fomesafen was applied under the plastic, bare ground, or over plastic mulch, refer to 24(c) label for specifics. -Maximum Reflex application in DE, MD, NJ, and VA: 24 fl oz/A IN ALTERNATE YEARS</p>						
15	Dual Magnum 7.62E	0.67 to 1.27 pt/A	s-metolachlor	0.64 to 1.21 lb/A	60	24
<p>-A Special Local Needs Label 24(c) has been approved for the use of Dual Magnum 7.62E to control weeds between the rows of plastic mulch in watermelon in DE and VA (expires 2/24/2021 for DE; 12/31/2021 for VA). The use of this product is legal ONLY if a waiver of liability is completed (see https://www.syngenta-us.com/labels/indemnified-label-login).</p> <p>-Plasticulture: row middle application only.</p> <p>-Do not apply Dual Magnum to the plastic mulch, or allow the spray to contact watermelon foliage. Do not soil incorporate.</p> <p>-Suppresses or controls annual grasses, yellow nutsedge, and certain annual broadleaf weeds including nightshade species. Use the lower rate on fields with coarse-textured soils low in organic matter. Use the higher rates on fields with fine-textured soil and those with high organic matter.</p> <p>-Maximum number of Dual Magnum applications per year is one and do not exceed 1.27 pt/A during the crop season.</p>						

2. Postemergence

Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC	6 to 8 fl oz/A	clethodim	0.094 to 0.13 lb/A	14	24
	Select Max 0.97EC	12 to 16 fl oz/A				
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.19 to 0.28 lb/A	14	12
<p>-Select 2EC: use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). Select Max: use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). Poast: use COC at 1.0% v/v.</p> <p>-The use of COC may increase the risk of crop injury when hot or humid conditions prevail. To reduce the risk of crop injury, omit additives or switch to NIS when grasses are small and soil moisture is adequate.</p> <p>-Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.</p> <p>-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled. Controls many annual and certain perennial grasses, including annual bluegrass, but Poast is preferred for goosegrass control. For best results, treat annual grasses when they are actively growing and before tillers are present. Control may be reduced if grasses are large or under hot or dry weather conditions.</p> <p>-Repeated applications may be necessary to control certain perennial grasses. If repeat applications are necessary, allow 14 days between applications. Rainfastness is 1 h.</p> <p>-Do not tank-mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses. Do not apply more than 8 fl oz of Select 2EC in a single application and do not exceed 32 fl oz/A for the season; do not apply more than 16 fl oz of Select Max in a single application and do not exceed 64 fl oz/A for the season.</p> <p>-Do not apply more than 1.5 pt/A Poast in single application and do not exceed 3 pt/A for the season.</p>						
14	Reflex 2SL	Rates vary, refer to the specific label	fomesafen	0.16 to 0.25 lb/A	35	24
<p>-A Special Local Needs Label 24(c) has been approved for the use of Reflex 2SL for Post-transplant control of weeds in watermelon in DE, MD, NJ, and VA (expires 12/31/2020 for DE, MD, VA, and 12/31/2020 for NJ). The use of this product is legal ONLY if a waiver of liability has been completed (see https://www.syngenta-us.com/labels/indemnified-label-login).</p> <p>-Rates vary by state and application method; refer to label to determine correct rates.</p> <p>-See soil applied section for application prior to planting or transplanting.</p> <p>-Plasticulture row middles with shielded/hood sprayers after transplanting; apply prior to vines "running" off the plastic. Severe crop injury can occur if spray comes in contact with crop foliage. Foliar application of Reflex will severely damage or kill watermelon.</p> <p>-Watermelon varieties may vary in their response to Reflex. Treat small acreages first to determine crop tolerance, especially when</p>						

2. Postemergence, Reflex - continued on next page

F Watermelons

2. Postemergence, Reflex - continued

<p>applying to a new variety. Treat small acreages first to determine crop tolerance, especially when applying to a new variety.</p> <p>-Reflex provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. Consider rotational crops when applying fomesafen. Rotational restrictions are dependent on whether fomesafen was applied under the plastic, bare ground, or over plastic mulch, refer to 24(c) label for specifics.</p> <p>-Consider rotational crops when applying fomesafen. If crop is replanted do not re-apply Reflex. Rotational restrictions are dependent on whether fomesafen was applied under the plastic, bare ground, or over plastic mulch, refer to 24(c) label for specifics.</p> <p>-Maximum Reflex application in DE, MD, NJ, and VA: 24 fl oz/A IN ALTERNATE YEARS</p>						
22	Gramoxone SL 2.0	1.95 pt/A	paraquat*	0.49 lb/A	14	24
<p>-A Supplemental Label has been approved for the use of Gramoxone 2SL for postemergence weed control in DE, MD, NJ, PA, and VA. Row middles as a shielded application. Apply as a directed spray in a minimum of 20 gal spray mix/A to control emerged weeds between the rows after crop establishment. Include a nonionic surfactant at 0.25% v/v. Use shields or hoods to prevent spray contact with the crop and low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. See the label for additional information and warnings.</p> <p>-Rainfastness is 30 min. A maximum of 3 applications per year are allowed.</p> <p>-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

3. Postharvest

Group	Product Name	Product Rate	Active Ingredient (*=Restricted Use)	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 2.0	2.25 to 3 pt/A	paraquat*	0.56 to 0.75 lb/A	--	24
<p>-A Special Local Needs Label 24(c) has been approved in VA (expires 12/31/2022) and a Supplemental Label in DE for the use of Gramoxone SL 2.0 for postharvest application to desiccate the crop.</p> <p>-Apply after the last harvest for bareground or plasticulture. Always include an adjuvant.</p> <p>-Spray coverage is essential for optimum effectiveness. See the label for additional information and warnings.</p> <p>-Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed.</p> <p>-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

4. Other Labeled Herbicides

These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name	Active Ingredient (*=Restricted Use)
2	League	imazosulfuron
3	Dacthal	DCPA
14	Aim	carfentrazone
14	Vida	pyraflufen

Insect Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Insecticides

Seed Corn Maggots

See also [Maggots](#) in section E 3.1 Soil Pests - Detection and Control.

Maggot problems can occur in the field and in transplant bedding trays in the greenhouse. An application of a soil-incorporated insecticide may be needed immediately before planting. The use of neonicotinoid insecticides (Group 4A) at planting may help to reduce seed corn maggot populations.

Aphids Note: Cultivars that are resistant to multiple aphid-transmitted viruses are available.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*=Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl* - melon aphid only	1-3	48	H
1B	Dimethoate 400	0.5 to 1.0 pt/A	dimethoate*	3	48	H
4A	Neonicotinoid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil/drip	21	4	M
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M

Aphids - continued on next page

Aphids - continued

9B	Fulfil 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
29	Beleaf 50SG	2.0 to 2.8 oz/A	flonicamid	0	12	L

Armyworms and Cabbage Loopers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
3A	Pyrethroid insecticides registered for use on Watermelons : see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
11A	Dipel DF, others (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
11A	XenTari (OMRI) (armyworms)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
11A	XenTari (OMRI) (cabbage loopers)	0.5 to 1.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel (armyworms)	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Exirel (cabbage loopers)	10.0 to 17.0 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Voliam Flexi (cabbage looper only)	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Cucumber Beetles

Watermelons are resistant to bacterial wilt; however, control may be needed to prevent feeding damage to seedlings. Seeds pretreated with a neonicotinoid seed treatment such Farmore DI-400 should provide up to 14 days of control of cucumber beetle. Otherwise, treat when an average of 2 beetles per plant is found. Management of adult cucumber beetles early in the season may help reduce damage to rinds later in the season.

Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
28	Exirel	20.5 fl oz/A	cyantraniliprole	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Cutworms See also section E 3.1. Soil Pests - Detection and Control.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV (variegated cutworm)	1.5 pt/A	methomyl*	1	48	H
1A	Lannate LV (granulate cutworm)	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					

F Watermelons

Leafminers

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
1B	Dimethoate 400	0.5 to 1.0 pt/A	dimethoate*	3	48	H
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	3	4	M
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	H
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Mites Mite infestations generally begin around field margins and grassy areas. **DO NOT mow or maintain these areas after midsummer** as this forces mites into the crop. Localized infestations can be spot treated. Begin treatment when 10-15 % of the crown leaves are infested early in the season, or when 50% of the terminal leaves are infested later in the season. Note: Continuous use of Sevin, or the pyrethroids may result in mite outbreaks.

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
6	Agri-Mek SC	1.75 to 3.5 fl oz/A	abamectin*	7	12	H
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L
20B	Kanemite 15SC	31.0 fl oz/A	acequinocyl	1	12	L
21 A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
21A	Portal XLO	2.0 pt/A	fenpyroximate	3	12	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M

Melonworms and Pickleworms

Apply one of the following formulations. If foliar materials are used, make one treatment prior to fruit set, and then treat weekly. If soil or drip applications are used, check the label for instructions on treatment frequency.						
Group	Product Name	Product Rate	Active Ingredient(s) (*= Restricted Use)	PHI (d)	REI (h)	Bee TR
1A	Lannate LV	1.5 to 3.0 pt/A	methomyl*	1-3	48	H
1A	Sevin XLR Plus	0.5 to 1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
3A + 4A	Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam	1	24	H
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	2.0 to 3.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole	30	12	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin*	7	12	H

Rindworms

For Lepidopteran rindworms, use one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L

Thrips

Apply one of the following formulations:						
Group	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Watermelons: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	3	4	M
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Group 3A Pyrethroid Insecticides Registered for Use on Watermelons

Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):

Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
Asana XL	5.8 to 9.6 fl oz/A	esfenvalerate*	3	12	H
Baythroid XL	0.8 to 2.8 fl oz/A	beta-cyfluthrin*	0	12	H
Bifenthrin 2EC, others	2.6 to 6.4 fl oz/A	bifenthrin*	3	12	H
Danitol 2.4EC	10.67 to 16.00 fl oz/A	fenpropathrin*	7	24	H
Hero EC	4.0 to 10.3 fl oz/A	zeta-cypermethrin* + bifenthrin*	3	12	H
Lambda-Cy 1EC, others	2.56 to 3.84 fl oz/A	lambda-cyhalothrin*	1	24	H
Mustang Maxx	1.28 to 4.0 fl oz/A	zeta-cypermethrin*	1	12	H
Permethrin 3.2EC, others	4.0 to 8.0 fl oz/A	permethrin*	0	12	H
Tombstone, others	0.8 to 2.8 fl oz/A	cyfluthrin*	0	12	H
Warrior II	1.28 to 1.92 fl oz/A	lambda-cyhalothrin*	1	24	H
Combo products containing a pyrethroid					
Endigo ZC	4.0 to 4.5 fl oz/A	lambda-cyhalothrin* + thiamethoxam (Group 4A)	1	24	H
Gladiator	19.0 fl oz/A	zeta-cypermethrin* + abamectin* (Group 6)	7	12	H
Besiege	6.0 to 9.0 fl oz/A	lambda-cyhalothrin* + chlorantraniliprole (Group 28)	1	24	H

Group 4A Neonicotinoid Insecticides Registered for Use on Watermelons

Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):

Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H
Assail 30SG	2.5 to 5.3 oz/A	acetamiprid	0	12	M
Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - soil/drip	21	12	H
Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - foliar (note: PHI: do not make application after 4 th true leaf has unfolded)	see note	12	H
Actara 25WDG	1.5 to 5.5 oz/A	thiamethoxam	0	12	H
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H
Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil/drip	21	12	H
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H
Venom 70SG	5.0 to 7.5 oz/A	dinotefuran - soil/drip	21	12	H
Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H
Combo products containing a neonicotinoid					
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole (Group 28)	1	12	H
Endigo ZC	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin* (Group 3A)	1	24	H

Angular Leaf Spot

At first sign of disease, apply the labeled rates of fixed copper plus mancozeb. Repeat every 7 d. To minimize the spread of disease, avoid working in field while foliage is wet.

Anthracnose

Excellent resistance is available in some varieties and those should be used when possible. Begin fungicide applications when vines run or earlier if symptoms are detected. **If resistance to FRAC code 11 (strobilurin) fungicides has been detected in the area, do not use Quadris, Quadris Top, Tanos or Cabrio.**

Code	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
Under LIGHT or MODERATE disease pressure, ALTERNATE:						
M05	chlorothalonil 6F	2.0 to 3.0 pt/A (low rate early in the season)	chlorothalonil	0	12	N
WITH a TANK MIX the following fungicide PLUS mancozeb 80 DF 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A:						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N
Under HIGH disease pressure, TANK-MIX one of the following fungicides WITH chlorothalonil 6F 2.0 to 3.0 pt/A:						
3 + 11	Quadris Top 1.67SC	12.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	0	12	--
3 + 11	Topguard 4.29SC	10.0 to 14.0 fl oz/A	flutriafol + azoxystrobin	1	12	--
7 + 11	Merivon 2.09SC	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	0	12	N
7 + 11	Pristine 38WG	18.5 oz/A	boscalid + pyraclostrobin	0	12	--
11	azoxystrobin 2.08F	11.0 to 15.5 fl oz/A	azoxystrobin	0	4	N
11	Cabrio 20EG	12.0 to 16.0 fl oz/A	pyraclostrobin	0	12	N
AND ROTATE with a TANK MIX of the following fungicide PLUS mancozeb 75DF 2.0 to 3.0 lb/A OR chlorothalonil 6F 2.0 to 3.0 pt/A every 7 days:						
1	thiophanate-methyl 70WP	0.5 lb/A	thiophanate-methyl	1	12	N

Bacterial Fruit Blotch (BFB)

Obtain seed or seedlings that were tested and found to have “no evidence” of the pathogen, which will reduce the risk of BFB development. Practice good sanitation during transplant production. Segregate different seed lots in the transplant house to reduce the chance of cross contamination. Scout seedlings daily, have suspect plants tested and destroy all diseased plants. Use only transplants from houses in which there were no seedling symptoms of BFB. If BFB is detected after transplanting, always work infested fields at the end of the day. Rotate to allow 2 years between watermelon plantings and control volunteers during those years.

Code	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
Apply one of the following fungicide schedules beginning before the first flower is open and continuing until 3 weeks after flowering. Subsequent fruit sets must also be protected.						
M01	copper (OMRI)	at labeled rates	copper	0	see label	N
P01	Actigard 50WG (must apply 1 or 2 weeks prior to flowering to be effective)	0.5 to 1.0 oz/A	acibenzolar-S-methyl	0	12	N

Downy Mildew

Scout fields for disease incidence regularly. Begin targeted sprays when disease occurrence is predicted for the region (check the Cucurbit Downy Mildew Forecasting website at <http://cdm.ipmpipe.org>). Strains of downy mildew that infect one cucurbit crop may not affect watermelon. Unnecessary fungicide application can be avoided by not spraying until disease is predicted in the region on watermelon. **Preventative applications are much more effective than applications made after detection. Materials with different Modes of Action (FRAC codes) should be alternated.** The following are the most effective products.

Code	Product Name	Product Rate	Active Ingredient(s) (* = Restricted Use)	PHI (d)	REI (h)	Bee TR
Sprays should be applied on a 7-day schedule when disease is forecast or present in the region. Under severe disease conditions and conducive weather, spray interval may be reduced IF the label allows.						
TANK-MIX one of these products WITH a protectant fungicide such as chlorothalonil 1.5 to 2.0 pt 6F/A:						
49+40	Orondis Ultra 2.33SC	5.5 to 8 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
49+M05	Orondis Opti 3.37SC	1.75 to 2.5 pt/A	oxathiapiprolin + chlorothalonil	0	12	--
21	Ranman 400SC (Do not apply with copper ; see label for details)	2.10 to 2.75 fl oz/A	cyazofamid	0	12	L

Downy Mildew - continued on next page

For Immediate Medical Attention

Call 911

**For a Pesticide Exposure Poisoning
Emergency Call**



For All States

This number will automatically connect you to the poison center nearest to you.

Anyone with a poisoning emergency can call the toll-free telephone number for help. Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

For Pesticide Spills

Small Spills: See the product label for cleanup advice.

Large spills: Call the National Response Center at 1-800-424-8802 or CHEMTREC at 800-424-9300 (24 hours) - Industry assistance with emergency response cleanup procedures for large, dangerous spills.

Be aware of your responsibility to report spills to the proper state agency.